



## Distribution of Microorganisms and Fractionation of Sulphur in Anthropogenic Wetlands under Long-term Elevated CO<sub>2</sub> Soil

Nyamath Syed <sup>a</sup>, Subburamu Karthikeyan <sup>a,b\*</sup>, Balachandar Dhananjayan <sup>a</sup>,  
Meena Suresh <sup>c</sup> and Sara Parwin Banu Kamaludeen <sup>d</sup>

<sup>a</sup> Department of Agricultural Microbiology, Directorate of Natural Resource Management, Agricultural College and Research Institute, Coimbatore-641 003, India.

<sup>b</sup> Centre for Post-Harvest Technology, Agricultural Engineering College and Research Institute, Coimbatore-641 003, India.

<sup>c</sup> Department of Soil Science and Agricultural Chemistry, Agricultural College and Research Institute, Trichy-620009, India.

<sup>d</sup> Department of Environmental Sciences, Tamil Nadu Agricultural University, Coimbatore-641 003, Tamil Nadu, India.

### Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

### Article Information

DOI: 10.9734/IJPSS/2022/v34i232536

### Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/93644>

Original Research Article

Received 06 September 2022  
Accepted 12 November 2022  
Published 16 November 2022

### ABSTRACT

The fate of extra carbon accessing soil under elevated CO<sub>2</sub> levels, as well as the repercussions for plant nutrition, is primarily determined by soil microbe activity. However, most increased CO<sub>2</sub> research has reported changes usually increases in soil organic carbon and reduction in the pH of the soil which is merely the first step in understanding how soil processes are altered. We analyzed these variables by assessing enzyme activities and identifying the individual components impacted by high CO<sub>2</sub> and those that reflect changes in soil organic matter pools. The majority of the microbial variables studied showed a significant increase under eCO<sub>2</sub> conditions, The rise in dehydrogenase activity suggests that the increased biomass of bacteria coincided with an increase in their activity. The rise in phosphatase activities implies that organic matter breakdown is being stimulated overall. The sulphur fractions had a significantly increased number of substrates consumed by soil microorganisms under increasing CO<sub>2</sub>. Moreover, direct examination of data from these perspective steep shifts in soil biological activity points to possible areas of investigation.

\*Corresponding author: E-mail: skarthy@tnau.ac.in;

**Keywords:** *Elevated CO<sub>2</sub>; soil organic matter; sulphur fractions; microbial distribution; soil dehydrogenase.*

## 1. INTRODUCTION

The world population is predicted to reach 9.1 billion by 2050, requiring a 60-110 % increase in the food supply [1]. Which would demand agricultural productivity enhancement given the restricted scope of arable land growth. Food security is a serious concern in nations with an increasing population and shrinking cultivable land space. Rice (*Oryza sativa* L.) is one of the most important staple crops, feeding almost half of the world population. Rice supplies more than half of the dietary requirements for 1.3 billion people, while rice provides 25-50 percent of the overall food requirements for 400 million individuals [2]. It is also a key source of calorie intake and nutrition, accounting for more than one-fifth of the calories ingested by humans globally. Rice is India's most significant staple food, consumed by around 65 percent of the population [3]. It is grown in almost all states, but West Bengal (13.79 percent), Uttar Pradesh (13.34 percent), Andhra Pradesh including Telangana (12.84 percent), Punjab (11.01 percent), Odisha (6.28 percent), Chhattisgarh (5.61 percent), Tamil Nadu (5.54 percent), Bihar (5.19 percent), Assam (4.41 percent), Haryana (3.88 percent), and Madhya Pradesh are the major rice producing states (3.86 percent).and it provides two-thirds of a person's calorie supply and one-half of total protein consumption [4]. Rice is special in that it can be produced in damp conditions when many other crops cannot. Such damp areas are common across Asia. However, soil fertility and fertilizer control are critical to a strong rice crop. Carbon and nutrient dynamics, which vary throughout agricultural ecosystems, are critical in soil fertility and crop yield [5]. As global temperatures rise due to greenhouse gas (GHG) emissions, sustainable agricultural output will be a major challenge in the future [6]. CO<sub>2</sub> is a significant GHG, accounting for approximately 60% of the entire greenhouse impact [7].

CO<sub>2</sub> emissions have increased in recent decades, elevating the atmospheric CO<sub>2</sub> level from 280 ppm at the start of the industrial revolution to 410 ppm now [8]. Human actions like fossil fuel combustion, indiscriminate land use, land cover, and so on are to blame for rising GHG concentrations in the atmosphere. In addition to water vapor and solar irradiance, well-mixed GHGs such as CO<sub>2</sub>, nitrous oxide (N<sub>2</sub>O), methane (CH<sub>4</sub>), and halocarbons contribute to

global temperature rises and climate change [9]. Global surface temperature is rising significantly, and crops may be unable to be cultivated in the future if such events continue [10]. According to an international team of 17 academics from 17 countries, temperature rises due to global warming could subsequently be double what climate models assess. Sustained CO<sub>2</sub> increases in the atmosphere would have an effect on soil and plant nutrition [11].

Sulphur, one of the most essential components for all plants and animals, is regarded as the fourth most important nutrient for agricultural crop development after nitrogen, phosphorus, and potassium. The relevance of sulphur in Indian agriculture is growing due to the recognition of its function in improving crop yield, not only for oil seeds, pulses, legumes, and forages but also for numerous kinds of cereal [12]. The sulphur deficit in crops is becoming more common as a result of the continued use of sulphur-free fertilizers, high-yielding crop types, intense multiple cropping systems, and increased productivity. Sulphur deficiency affects around 41% of India's farmed land. In India, crops remove around 1.26 million tons of sulphur, while fertilizers restore approximately 0.76 million tonnes [13]. Furthermore, the recovery of additional sulphur from external sources is approximately 8-10%. The continued depletion of local sulphur deposits throughout the post-green revolution period has resulted in a sulphur deficit in many sections of the country [14]. Sulphur exists in both inorganic and organic forms in soils, and the proportion of inorganic to organic sulphur varies greatly depending on the composition of the soil, its depth, and the management variables to which the soil is subjected. In most arable land top soils, total soil sulphur, which includes inorganic and organic binding forms, varies between 250 and 2500 kg/ha. Water soluble and adsorbed SO<sub>4</sub><sup>2-</sup> composes inorganic " S "which is widely assumed to be the direct source of plants. In general, it accounts for less than 5% of total soil S. SO<sub>4</sub><sup>2-</sup> is present in soil solutions only present in trace amounts that vary continually, and its concentration at any one moment is determined by the balance of plant uptake, and fertilizer input, mineralization, and immobilization. Understanding the S transformations in connection to crop availability in soil is required to utilize S fertilizers more efficiently.

Recent studies reveal that if temperatures rise by more than 0.5°C, the effects will be disastrous for millions of the world's poorest people due to reduced food output. As eCO<sub>2</sub> causes, an increase in global temperature in the future, soil health, nutrient availability, and crop output are all expected to suffer. Adaptation studies are being conducted all around the world, although they are often insufficient. India is far behind in terms of studies on the effects of climate change on agriculture. As a result, the current study was carried out to investigate the impacts of eCO<sub>2</sub> on the dynamics of microbes and sulphur fractions in the soil.

## 2. MATERIALS AND METHODS

### 2.1 Site Description

The study was conducted in the A1 field in the Wetland Farms, Tamil Nadu Agricultural University, Coimbatore, India (11°005'13" N, 76° 93'045" E) which is under long-term elevated (eCO<sub>2</sub>) 550±30 for a decade. Soil sampling was conducted on a randomized quadrant basis during the year 2018. A total of three samples with three replications have been collected from (i) Long-term elevated CO<sub>2</sub> open-top chamber (OTC) (eCO<sub>2</sub>) (ii) OTC without any CO<sub>2</sub>, ambient (aCO<sub>2</sub>), and (iii) Soil samples from a normal low-land rice ecosystem in the ambient bulk soil (ABS).

### 2.2 Enumeration of Functional Guilds

Enumeration of different functional bacterial groups. The media employed for the isolation of sulphur-oxidizing bacteria is mineral salt thiosulphate medium [15]. For the enumeration of methanogens, the [16] method was followed. For the enumeration of sulphur-reducing bacteria by [17] method. Methanotrophs enumerated by NMS medium. Diazotrophs and heterotrophs were enumerated by the method given by [18].

### 2.3 Soil Enzyme Analysis

All enzyme activities can be carried out by processing the soil size to < 2mm and following the colorimetric method for the analysis of the dehydrogenase [19], acid and alkaline phosphatase [20] beta-glucosidase [21] in the soil was estimated by the standard methods -, aryl sulphatase by [22]. and florescent diacetate hydrolysis was carried out by the standard method [23].

## 2.4 Sulphur Fractions in Soil

### 2.4.1 Fractionation of sulphur

Less than 2 mm of soil samples were used in the sequential extraction of soil sulphur fractions [24]. The smaller fractions (water soluble) were extracted with demineralized water for the first time at (w/v) 1:10. After 30 minutes of shaking, each sample was analyzed. The sample was spun at 10,000 rpm for 10 minutes, and the homogenate was assessed using the turbidity method. Exchangeable sulphur fractions were extracted at a 1:10 ratio with 0.032M NaH<sub>2</sub>PO<sub>4</sub> (w/v) proportion after 30 minutes of shaking, each sample was centrifuged at 10,000 rpm for 10 minutes and the supernatant was analyzed by turbidity method. For precipitated sulphur, the remainder of the soil sample was extracted by 1M HCl. The ratio of extraction was 1:20 (w/v). After 60 minutes of shaking, the samples were ready centrifuged at 10,000 rpm for 10 minutes, and analyzed by turbidity method.

### 2.5 Available Sulphur

Five grams of soil was taken in a polythene shaking bottle and put it in a shaker container. 25 ml 0.15 percent CaCl<sub>2</sub> solution, shake for 30 minutes, filter with Whatman no.42 filter paper, pipette out 5 ml extract in 25 ml volumetric flask, then add 10 ml sodium acetate and acetic acid buffer after that, add 1 gram of barium chloride crystals and mix for 1 minute. After that, add 1 ml of 0.25 percent gum acacia solution and dilute to 25 ml with distilled water and shake until the barium chloride crystals dissolve 2-8 minutes after 420nm absorption. Calculate the concentration of accessible sulphur in the soil sample by plotting the graph with a standard curve using K<sub>2</sub>SO<sub>4</sub>.

### 2.6 Statistical Analysis

All the data sets recorded were analyzed by SPSS version 16.0. means of values compared by the least significant difference (LSD) at a 5 % level of probability.

## 3. RESULTS AND DISCUSSION

### 3.1 Effect of eCO<sub>2</sub> Concentrations on Soil Chemical Properties. Soil pH, Electrical Conductivity, and Organic Carbon in Soil

The pH of the soil was decreased as CO<sub>2</sub> concentration rise (Table 1). The lowest pH value

(7.9) was noticed with 550 ppm eCO<sub>2</sub>, while the highest pH value (8.2) was observed in ambient bulk soil. According to [25], eCO<sub>2</sub> is anticipated to raise CO<sub>2</sub> concentration in soil via the generation of H<sub>2</sub>CO<sub>3</sub>, and hence pH in the soil is likely to drop. The majority of plant nutrients become accessible when soil pH varies from 6.5 to 8.0. When submerged, the pH of the soil normally reaches 7.0. According to [26], an increase in CO<sub>2</sub> concentration affects soil pH. However, investigations have revealed that when a large quantity of CO<sub>2</sub> was added to the soil system, the pH changed insignificantly. It is also insignificant when compared to the depth of the soil. This may be due to the solubility of carbon dioxide (CO<sub>2</sub>) in water, which results in the formation of undissociated carbonic acid. Acidic root exudates from plants and other microorganisms, on the other hand, lower pH and alter the solubility relationship of many plant nutrients. Acetic acid is the most common anaerobic respiration intermediate. During the decomposition process, all organic matter, including fats, amino acids, and carbohydrates, is converted to acetic acid. [27–29]. An increase in the eCO<sub>2</sub> increases the soil's organic carbon. The soil organic carbon was found higher in the eCO<sub>2</sub> range of 6.9 mg/kg while the lowest was observed in the ambient bulk soil range of 5.7 mg/kg. These findings support [30]. The suggestion is that physical fractionation techniques can aid in identifying changes in the soil carbon pool under higher CO<sub>2</sub> conditions, even in trials with high intrinsic variability and beginning C content, as in our experiment. The increased sensitivity of SOC to changes in C dynamics with rising CO<sub>2</sub> than total C is most likely due to considerably quicker turnover in SOM pools (Tables 1 & 3). The greater proportion of new SOC relative to total new soil versus the proportion of total SOC related to total new soil C indicates a quicker turnover of SOC. SOC content primarily through boosting photosynthate synthesis and allocation to the rhizosphere, hence increasing C intake into the soil [31]. Soil organic matter is a key factor that affects all physical, chemical, and biological activity in soils. It is soil's essence that enhances soil health, agricultural productivity, and soil aggregates. More C sequestration in soils equals lower CO<sub>2</sub> levels in the atmosphere, reducing the potential for global warming [32].

### 3.2 Microbial Distribution

The development of bacteria has shown an increasing tendency as CO<sub>2</sub> levels rise (Fig. 1).

However, insignificant increase in microbial populations of bacteria, fungi, actinomycetes, heterotrophs, and diazotrophs. There was a significant difference observed in methanotrophs, Sulphur oxidizers, methanogens, and phosphorous solubilizing bacteria at ( $p < 0.01$ ) and ( $p < 0.05$ ) levels of significance among the three soils. Insignificant decrease of SRB under eCO<sub>2</sub> is observed this may be due to the proliferation of methanogens by utilizing CO<sub>2</sub> as a substrate and limiting the substrates required for the sulphidogenesis. The highest microbial populations were seen in soil treated with eCO<sub>2</sub> followed by ambient CO<sub>2</sub> soil, respectively. Variations in the microbial population may have altered the breakdown of organic molecules. Furthermore, bacteria account for around one-quarter of all living biomass on the planet, and their biomass is employed as an early indication of changing physical and chemical features of soils. Increased microbial growth was associated with higher levels of plant-derived C in the rhizosphere under eCO<sub>2</sub>. Which is consistent with previous research using other plant species [33–35]. Increase in the eCO<sub>2</sub> increases the soil temperature and thus influences the total the microbial communities of bacteria, diazotrophs, fungi, actinobacteria, methanotrophs, and diazotrophs (Lou et al., 2006). In present soil hike in total anaerobic population under eCO<sub>2</sub> and decrease in methanotrophic population compared bulk soil as shown in Fig. 1. Similar results were obtained by [36]. Increases in the enzyme activities in soils boost up the carbon pools and diazotrophic population which was observed in our study (Fig. 1 & Table 2). Similar results were observed by [37]. Significantly higher microbial ratios under enhanced CO<sub>2</sub> treatments than under ambient CO<sub>2</sub> imply that CO<sub>2</sub> amplification of C inputs may still play a significant role in defining community structure in N-rich agroecosystems, as seen in many forests and grasslands [38]. Organic matter concentration, water availability, and soil physical and chemical qualities all have an impact on microbial biomass.


### 3.3 Enzyme Activities in Soil

Soil enzymes profit from biological and metabolic activities in soil, which form the foundation of the terrestrial ecosystem. Elevated atmospheric CO<sub>2</sub> levels clearly have an effect on the soil enzyme activity that is influenced by the physical and chemical properties of soil, soil biological community, and plants either directly or indirectly. All the values are calculated by using a

**Table 1. Soil physical properties from three different soils elevated, ambient, and ambient bulk soil**

| Soil parameters                  | Elevated (eCO <sub>2</sub> ) | Ambient                | Ambient bulk soil     |
|----------------------------------|------------------------------|------------------------|-----------------------|
| pH                               | 7.9±0.1 <sup>a</sup>         | 8.1±0.3 <sup>b</sup>   | 8.2±0.4 <sup>b</sup>  |
| EC                               | 0.76±0.03 <sup>b</sup>       | 0.75±0.02 <sup>b</sup> | 0.8±0.01 <sup>a</sup> |
| Bulk density(mg/m <sup>3</sup> ) | 1.49 ±0.1 <sup>a</sup>       | 1.49±0.1 <sup>a</sup>  | 1.49±0.2 <sup>a</sup> |
| SOC (g/kg)                       | 6.9±0.1 <sup>a</sup>         | 5.8±0.4 <sup>b</sup>   | 5.7±0.2 <sup>b</sup>  |

All the values are means of ±SD



**Fig. 1. Distribution of different groups of microbes. Bacteria, fungi, actinomycetes, heterotrophs, diazotrophs, methanotrophs, sulphur-oxidizing bacteria, sulphur reducing bacteria, methanogens, and phosphate solubilizing bacteria from three different soils elevated(eCO<sub>2</sub>), ambient(A), and ambient bulk soil (ABS)**

standard graph with an ( $R^2 > 0.90$ ). There were significant changes observed in all enzyme activity in the soils at ( $p < 0.01$ ) and ( $p < 0.05$ ) levels of significance respectively. Where the highest values for all the samples were observed in the soil treated under long-term elevated CO<sub>2</sub> as shown in Table 2. Similar results were also obtained by [39]. The stimulation in enzyme activities by increasing CO<sub>2</sub> demonstrated that increased microbial biomass was accompanied by increased activity [40]. It is recognized that increased CO<sub>2</sub> levels can boost carbon input into the soil by improving the allocation of fixed carbon below ground [41]. This procedure can provide some information on future enzyme activity or the sum of active enzymes. However, because enzyme activities (particularly immobilized enzymes) are affected by edaphic circumstances, actual activity may differ from potential activity [42].

### 3.4 Sulphur Fractions

The content of S and sulphur fractions in the soil increased significantly due to eCO<sub>2</sub> in OTC, the

available sulphur ranging from 13.49 to 17.13 mg kg<sup>-1</sup> (Fig. 2). The largest available S content was discovered in the 550 ppm CO<sub>2</sub> treated OTC 17.13 mg kg<sup>-1</sup> followed by ambient soil 14.48 mg kg<sup>-1</sup>) and the lowest value was reported in ambient bulk soil 13.49 mg kg<sup>-1</sup>. Fractions of sulphur in the soils follow the order (AS ≥ WSS < ES < PS) Shown in Fig. 2. The capacity of soils to adsorb sulfate varies greatly due to the clay composition of soils containing different proportions of hydrous oxides of Al and Fe. However, [43] found that eCO<sub>2</sub> had little effect on soil S content. According to [44], diminished pH enhances sulphate adhesion in soil systems.

### 3.5 Total Carbon, Nitrogen, Sulphur, and Available NPK in Soil

The quantity of CO<sub>2</sub> within the OTC had a detrimental influence on available soil N, P, and K content, which increased as CO<sub>2</sub> concentration increased significantly at 1% and 5% levels of significance (Table 3). The maximum soil NPK

**Table 2. Soil enzyme activities from three different soils elevated, ambient, and outside ambient**


| Soil parameters  | Soil Dehydrogenase ( $\mu\text{g TPF g}^{-1}$ soil $24 \text{ hr}^{-1}$ ) | Beta Glucosidase ( $\mu\text{g pNP g}^{-1}$ soil) | Fluorescent Diacetate ( $\mu\text{g fluorescein released g}^{-1}$ soil) | Alkaline phosphatase ( $\mu\text{g pNP g}^{-1}$ soil) | Acid phosphatase ( $\mu\text{g pNP g}^{-1}$ soil) | Urease ( $\mu\text{g NH}_3 \text{ g}^{-1}$ soil) | Aryl Sulphatase ( $\text{nmol g}^{-1} \text{min}^{-1}$ ) |
|------------------|---|---|---|---|---|--|--|
| eCO <sub>2</sub> | 10.02±0.23 <sup>a</sup>   | 132.35±1.29 <sup>a</sup>                          | 4.04±0.04 <sup>a</sup>  | 57.69±0.21 <sup>a</sup>                               | 36.04±0.12 <sup>a</sup>                           | 48.16±0.39 <sup>a</sup>                          | 12.20±0.11 <sup>a</sup>                                  |
| A                | 5.38±0.27 <sup>c</sup>  | 130.38±1.54 <sup>a</sup>                          | 3.92±0.08 <sup>b</sup>  | 49.01±0.15 <sup>b</sup>                               | 31.41±0.19 <sup>a</sup>                           | 41.36±0.34 <sup>b</sup>                          | 9.55±0.12 <sup>b</sup>                                   |
| ABS              | 8.22±0.26 <sup>b</sup>  | 118.92±1.34 <sup>b</sup>                          | 2.96±0.09 <sup>c</sup>  | 45.93±0.18 <sup>b</sup>                               | 36.42±0.26 <sup>a</sup>                           | 34.85±0.26 <sup>c</sup>                          | 8.94±0.16 <sup>c</sup>                                   |

*All the values are means of ±SD. Significant change in all the enzyme activities at 1% and 5% level of significance*

**Table 3. Distribution of carbon, nitrogen, and sulphur in three different soils elevated, ambient, and ambient bulk soil**

| Soil parameters   | Elevated CO <sub>2</sub>  | Ambient soil             | Ambient bulk soil        |
|-------------------|---------------------------|--------------------------|--------------------------|
| Available N kg/ha | 285.6±2.2 <sup>a</sup>    | 268.8±3.6 <sup>a</sup>   | 246.4±3.9 <sup>b</sup>   |
| Available P Kk/ha | 37.01±0.56 <sup>a</sup>   | 32.48±0.94 <sup>b</sup>  | 28.95±0.85 <sup>c</sup>  |
| Available K kg/ha | 490±12 <sup>a</sup>       | 470±14 <sup>b</sup>      | 454±18 <sup>c</sup>      |
| Total N (%)       | 0.0784±0.005 <sup>a</sup> | 0.0795±0.09 <sup>a</sup> | 0.0756±0.06 <sup>a</sup> |
| Total S (%)       | 0.42±0.08 <sup>a</sup>    | 0.41±0.09 <sup>a</sup>   | 0.41±0.06 <sup>a</sup>   |
| Total C (%)       | 0.88±0.02 <sup>a</sup>    | 0.87±0.02 <sup>a</sup>   | 0.87±0.01 <sup>a</sup>   |

*All the values are means of ±SD. Significant difference found among available N, P, K at (p<0.01) and (p<0.05) level of significance*



**Fig. 2. Distribution of sulphur fractions Available Sulphur (AS), Water-Soluble Sulphur (WSS), Exchangeable Sulphur (ES), and Precipitated Sulphur (PS) from three different soils elevated, ambient, and ambient bulk soil**

concentration, was found with eCO<sub>2</sub>, followed by ambient and ambient bulk soil respectively as shown in Table 3, Plant requirement for nitrogen (N) is predicted to grow as atmospheric CO<sub>2</sub> levels rise (eCO<sub>2</sub>). Nitrogen supplementation improves the impact of CO<sub>2</sub> on plant production. Furthermore, eCO<sub>2</sub> greatly improved N absorption [45,46]. As a result, the amount of the response in plant productivity is primarily determined by how well plant N absorption can keep up with eCO<sub>2</sub>-induced stimulation of carbohydrate formation and growth. Plants may positively control a number of physiological processes, such as enzyme production and root development, to boost plant nutrient acquisition capacity for optimum eCO<sub>2</sub> adaptation [47]. Significant increases in available P requirement by plants are predicted under increased CO<sub>2</sub> levels due to photosynthetic stimulation and subsequent growth responses. Elevated CO<sub>2</sub> affects P acquisition by altering root shape and increasing rooting depth. Furthermore, with high CO<sub>2</sub>, the amount and content of root exudates are anticipated to alter due to changes in carbon fluxes through the glycolytic pathway and the tricarboxylic acid cycle. As a result, root exudates may induce P mobilization via alkylation of P from low solubility P complexes, shifts in the biochemical environment, and alters indigenous soil microbial activity.[48]. Significant increases in the available potassium in the soil due to lowering of pH in soil due to the formation of carbonic acid and solubility of the ions increases in soil. Increases in the available potassium under elevated CO<sub>2</sub> have been reported by [49]. In terms of total

carbon nitrogen and sulphur, no significant difference was found among the three different treatments observed.

#### 4. CONCLUSION

Elevated CO<sub>2</sub> levels are responsible for rising global temperatures, which may have a significant influence on soil health, crop output, and plant nutrient usage efficiency. In the current experiment, the findings showed that the responses of soil microbes and their community structure to elevated CO<sub>2</sub> changed significantly in the rice low land ecosystem which is under long-term elevated. S fractionation in rice field is a powerful tool for understanding S dynamics and biogeochemistry in wetland ecosystems. eCO<sub>2</sub> greatly influences the fractionation of sulphur. The study found that all sulphur fractions, including available S, water soluble S, and exchangeable S, precipitated S and total S were significantly influenced. Precipitated sulphur was discovered to be the most dominant among the observed fractions. These changes may be largely related to CO<sub>2</sub>-induced changes in soil C, N and S availability.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Wani SH, Sah SK. Biotechnology and abiotic stress tolerance in rice. *J Rice Res.* 2014;2:105.

2. Lavakush, Yadav J, Verma JP, et al. Evaluation of PGPR and different concentration of phosphorus level on plant growth, yield and nutrient content of rice (*Oryza sativa*). Ecol Eng. 2014;62: 123–128.  
Available:<https://doi.org/10.1016/j.ecoleng.2013.10.013>
3. Singh KM, Singh D. Challenges of ensuring food and nutritional security in Bihar. Challenges Ensuring Food Nutr Secur Bihar; 2020.
4. Alam MA, Rahman MM, Khan MAR, et al. Modulation of yield and yield attributes of transplanted aman rice through organic and inorganic fertilization in terrace soil. Eur J Agric Food Sci. 2021;3:97–105.
5. Alam MA, Rahman MM, Biswas JC, et al. Nitrogen transformation and carbon sequestration in wetland paddy field of Bangladesh. Paddy Water Environ. 2019; 17:677–688.  
Available:<https://doi.org/10.1007/s10333-019-00693-7>
6. Rahman MM. Nutrient-use and carbon-sequestration efficiencies in soils from different organic wastes in rice and tomato cultivation. Commun Soil Sci Plant Anal. 2013;44:1457–1471.
7. Rastogi M, Singh S, Pathak H. Emission of carbon dioxide from soil. Curr Sci. 2002;82:510–517.
8. Kahn H. On escalation: Metaphors and scenarios. 2009;1.
9. Mackay A. Climate change 2007: Impacts, adaptation and vulnerability. Contribution of working group ii to the fourth assessment report of the intergovernmental panel on climate change. J Environ Qual. 2008;37:2407.
10. Risbey JS. The new climate discourse: Alarmist or alarming? Glob Environ Chang. 2008;18:26–37.
11. Razzaque MA, Haque MM, Khaliq QA, Solaiman ARM. The effect of different nitrogen levels and enrichment CO<sub>2</sub> on the nutrient contents of rice cultivars. Bangladesh. J Sci Ind Res. 2009;44:241–246.
12. Singh AK, Amgain LP, Sharma SK. Root characteristics, soil physical properties and yield of rice (*Oryza sativa*) as influenced by integrated nutrient management in rice-wheat (*Triticum aestivum*) system. Indian J Agron. 2000;45:217–222.
13. Tiwari KN, Gupta BR. Sulphur for sustainable high yield agriculture in Uttar Pradesh. Indian J Fertil. 2006;1:37.
14. Sharma U, Subehia SK, Rana SS, et al. Soil sulphur fractions and their relationship with soil properties and rice (*Oryza sativa* L.) yield under long-term integrated nutrient management in an acid alfisol. Res Crop. 2014;15.
15. Vidyalakshmi R, Sridar R. Isolation and characterization of sulphur oxidizing bacteria. J Cult Collect. 2007;5:73–77.
16. Hungate RE. Chapter IV A roll tube method for cultivation of strict anaerobes. In: methods in microbiology. Elsevier. 1969;117–132.
17. Postgate JR. Versatile medium for the enumeration of sulfate-reducing bacteria. Appl Microbiol. 1963;11:265–267.
18. Döbereiner J. Isolation and identification of root associated diazotrophs in: Nitrogen fixation with non-legumes. Springer. 1989; 103–108.
19. Casida Jr. LE, Klein DA, Santoro T. Soil dehydrogenase activity. Soil Sci.1964;98:371–376.
20. Tabatabai MA, Bremner JM. Use of p-nitrophenyl phosphate for assay of soil phosphatase activity. Soil Biol Biochem. 1969;1:301–307.
21. Eivazi F, Tabatabai MA. Factors affecting glucosidase and galactosidase activities in soils. Soil Biol Biochem. 1990;22:891–897.
22. Tabatabai MA, Bremner JM. Arylsulfatase activity of soils. Soil Sci Soc Am J. 1970; 34:225–229.
23. Schnierer J, Rosswall T. Fluorescein diacetate hydrolysis as a measure of total microbial activity in soil and litter. Appl Environ Microbiol. 1982;43:1256–1261.
24. Morche L. S-fluxes and spatial alterations of inorganic and organic sulphur fractions in soil as well as their accumulation and depletion in the rhizosphere of agricultural crops by partial use of the radioisotope <sup>35</sup>S. Bonn Rheinischen Friedrich-Wilhelms-Universität Bonn (in Germany); 2008.
25. Carrillo Y, Dijkstra FA, Pendall E, et al. Plant rhizosphere influence on microbial C metabolism: The role of elevated CO<sub>2</sub>, N



- availability and root stoichiometry. *Biogeochemistry*. 2014;117:229–240.
26. Celia MA, Peters CA, Bachu S. Geologic storage of CO<sub>2</sub>: Leakage pathways and environmental risks. In: AGU. Spring Meeting Abstracts. 2002;GC32A-03.
  27. Cardon ZG. Influence of rhizodeposition under elevated CO<sub>2</sub> on plant nutrition and soil organic matter. *Plant Soil*. 1995; 187:277–288.
  28. Lindsay WL. Chemical equilibria in soils. John Wiley and Sons Ltd; 1979.
  29. Patil RH, Colls JJ, Steven MD. Effects of CO<sub>2</sub> gas as leaks from geological storage sites on agro-ecosystems. *Energy*. 2010; 35:4587–4591.
  30. Canadell JG, Pitelka LF, Ingram JSI. The effects of elevated [CO<sub>2</sub>] on plant-soil carbon below-ground: A summary and synthesis. *Plant Soil*. 1995;187:391–400.
  31. Satapathy SS, Swain DK, Pasupalak S, Bhadoria PBS. Effect of elevated [CO<sub>2</sub>] and nutrient management on wet and dry season rice production in subtropical India. *Crop J*. 2015;3:468–480.
  32. Six J, Carpentier A, van Kessel C, et al. Impact of elevated CO<sub>2</sub> on soil organic matter dynamics as related to changes in aggregate turnover and residue quality. *Plant Soil*. 2001;234:27–36.
  33. Drigo B, Kowalchuk GA, Knapp BA, et al. Impacts of 3 years of elevated atmospheric CO<sub>2</sub> on rhizosphere carbon flow and microbial community dynamics. *Glob Chang Biol*. 2013;19:621–636.
  34. Da Costa DP, Dias ACF, Cotta SR, et al. (2018) Changes of bacterial communities in the rhizosphere of sugarcane under elevated concentration of atmospheric CO<sub>2</sub>. *GCB Bioenergy*. 2018;10:137–145.
  35. Wang YH, Yu ZH, Li YS, et al. Elevated CO<sub>2</sub> alters the structure of the bacterial community assimilating plant-derived carbon in the rhizosphere of soya bean. *Eur J Soil Sci*. 2019;70:1212–1220.
  36. Liu D, Tago K, Hayatsu M, et al. Effect of elevated CO<sub>2</sub> concentration, elevated temperature and no nitrogen fertilization on methanogenic archaeal and methane-oxidizing bacterial community structures in paddy soil. *Microbes Environ*. 2016;31:349–356.
  37. Kumar U, Nayak AK, Sahoo S, et al. Combined effects of elevated CO<sub>2</sub>, N fertilizer and water deficit stress on diazotrophic community in sub-humid tropical paddy soil. *Appl Soil Ecol*. 2020; 155:103682.
  38. Feng X, Simpson AJ, Schlesinger WH, Simpson MJ. Altered microbial community structure and organic matter composition under elevated CO<sub>2</sub> and N fertilization in the duke forest. *Glob Chang Biol*. 2010; 16:2104–2116.
  39. Dhillon SS, Roy J, Abrams M. Assessing the impact of elevated CO<sub>2</sub> on soil microbial activity in a Mediterranean model ecosystem. *Plant Soil*. 1995;187: 333–342.
  40. Ebersberger D, Niklaus PA, Kandeler E. Long term CO<sub>2</sub> enrichment stimulates N-mineralisation and enzyme activities in calcareous grassland. *Soil Biol Biochem*. 2003;35:965–972.
  41. Jones TH, Thompson LJ, Lawton JH, et al. Impacts of rising atmospheric carbon dioxide on model terrestrial ecosystems. *Science*. 1998;280(80):441–443.
  42. Freeman C, Liska G, Ostle NJ, et al. The use of fluorogenic substrates for measuring enzyme activity in peatlands. *Plant Soil*. 1995;175:147–152.
  43. Fangmeier A, De Temmerman L, Black C, et al. Effects of elevated CO<sub>2</sub> and/or ozone on nutrient concentrations and nutrient uptake of potatoes. *Eur J Agron*. 2002; 17:353–368.
  44. Shainberg I, Sumner ME, Miller WP, et al. Use of gypsum on soils: A review. *Adv soil Sci*. 1989;1–111.
  45. Ainsworth EA, Long SP. What have we learned from 15 years of free-air CO<sub>2</sub> enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO<sub>2</sub>. *New Phytol*. 2005;165:351–372.
  46. Langlely JA, Megonigal JP. Ecosystem response to elevated CO<sub>2</sub> levels limited by nitrogen-induced plant species shift. *Nature*. 2010;466:96–99.
  47. Li Y, Yu Z, Liu X, et al. Elevated CO<sub>2</sub> increases nitrogen fixation at the reproductive phase contributing to various yield responses of soybean cultivars. *Front Plant Sci*. 2017;8:1546.
  48. Jin J, Tang C, Sale P. The impact of elevated carbon dioxide on the phosphorus

- nutrition of plants: A review. Ann Bot. 2015;116:987–999.
49. Reddy KR, Zhao D. Interactive effects of elevated CO<sub>2</sub> and potassium deficiency on photosynthesis, growth, and biomass partitioning of cotton. F Crop Res. 2005; 94: 201–213.

---

© 2022 Syed et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/93644>