



Role of Condensed Tannins in the Antioxidant Activity of Seeds of *Cola nitida* (Vent.) Schott & Endl. (*Malvaceae*)

Abdou Sarr ^{a*}, Serigne Ibra Mbacké Dieng ^a,
Madièye Sene ^b, Charlot Diatta ^b, Kady Diatta-Badji ^a,
William Diatta ^a and Alioune Dior Fall ^a

^a Laboratory of Pharmacognosy and Botany, Faculty of Medicine, Pharmacy and Odontology, Cheikh Anta Diop University, Dakar, Senegal.

^b Laboratory of Pharmacology, Faculty of Medicine, Pharmacy and Odontology, Cheikh Anta Diop University, Dakar, Senegal.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/EJMP/2022/v33i121111

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/93755>

Original Research Article

Received: 17/09/2022
Accepted: 19/11/2022
Published: 22/11/2022

ABSTRACT

Background: Native to West Africa, *Cola nitida* is a tropical tree of about 8-12 in height that grows in lowland rainforest. This plant is best known in Africa for its seeds, used in phytotherapy but for their socio-cultural importance.

Aims/Objective: This study investigated the antioxidant activity of condensed tannins of *Cola nitida* seeds by carrying out two antioxidant tests (DPPH and FRAP).

*Corresponding author: E-mail: abdou.sarr@ucad.edu.sn;

Methods: From a hydro-ethanolic extract of *Cola nitida* seeds, two samples were made. One treated with casein (EC) and another one without treatment (EWC). The research of condensed tannins were carried out by precipitation with Stiasny reagent. The total polyphenol and tannin contents were evaluated by the Folin-Denis method and the antioxidant power by DPPH and FRAP tests.

Results: Extract without treatment (EWC) showed more antioxidant activity than the extract treated with casein (EC). Thus, the IC_{50} of EWC which contains condensed tannins was $5.54 \pm 0.005 \mu\text{g/ml}$, while that of EC (without condensed tannins) reached $61.92 \pm 0.165 \mu\text{g/ml}$.

Conclusion: *Cola nitida* seeds are rich in condensed tannins that play an important role in the antioxidant activity.

Keywords: *Cola nitida*; seeds; polyphenols; tannins; antioxidant activity.

1. INTRODUCTION

Natural products, especially those of plant origin, have always been an important source of bioactive molecules. Currently, about 25-30% of drugs available for the treatment of diseases are derived from natural products (plants, animals, bacteria and fungi) [1]. According to the World Health Organization [2], traditional medicine is very important but often underestimated in healthcare. It exists in almost every country in the world, and the demand for services in this area is growing.

Nowadays, numerous studies carried out in the field of ethnopharmacology had shown that most of the plants used in traditional medicine and tested in laboratories contained secondary metabolites that are active in pharmacological models.

Among these metabolites, polyphenols occupy a prominent place. They represent a large chemical family including tannins, flavonoids, anthocyanins, etc. Several biological activities have been attributed to polyphenols. These compounds had shown anti-carcinogenic, anti-inflammatory, antiatherogenic, anti-thrombotic, analgesic, antibacterial, antiviral, anticancer [3], anti-allergenic, vasodilator [4] and antioxidant activities [5].

The plant used for this study was *Cola nitida* (Vent) Schott. and Endl. The seeds were chosen by their several uses in Africa traditional medicine against many diseases such as dysentery, cough, diarrhea, vomiting [6], and by the fact that they have potential for pharmaceutical and food development.

The aim of this study was to evaluate the role of condensed tannins in the antioxidant activity of *Cola nitida* seeds.

2. MATERIALS AND METHODS

2.1 Plant Material

The plant material consists of seeds of *Cola nitida* (Vent) Schott. and Endl. (*Malvaceae*) which were purchased at the Sandaga market in Dakar (Senegal). The drug was identified at the Laboratory of Pharmacognosy and Botany of the Faculty of Medicine, Pharmacy and Odontostomatology (Cheikh Anta DIOP University of Dakar). *Cola nitida* is registered in the herbarium of the same institute under the number of 95. The seeds were crushed into small pieces and dried at room temperature before being ground into a powder using an electric grinder fitted with a medium-sized sieve. The powder was stored in a sachet protected from light.

2.2 Methods

2.2.1 Preparation of plant extracts

For extraction, 30 g of *Cola nitida* seeds powder were introduced into a cartridge and extracted with 700 ml of the ethanol-water mixture (80/20; v/v), for 24 hours in a Soxhlet apparatus [7]. The solution obtained was evaporated using a rotary evaporator at 65°C . The residue resulting from this operation was dried in an oven for 48 hours at 45°C to obtain the dry hydro-ethanolic extract (EC).

From the dry hydro-ethanolic extract, 60 ml of an aqueous solution at $50 \mu\text{g/ml}$ were prepared. The solution was then divided into two 100 ml Erlenmeyer flasks at the rate of 30 ml for each container. In one of the Erlenmeyer flasks, 3 grams of casein are added and the whole was submitted to magnetic stirring for 3 hours in order to fix the condensed tannins [8]. The Stiasny reaction was used to check whether all the condensed tannins have been fixed by the

casein. This process led to the extract without condensed tannins (EC). The two solutions thus obtained (EWC and EC) were used for the tests of antioxidant activity and for the determination of total polyphenols.

2.2.2 Polyphenols and condensed tannins contents

The method used is that described by Joslyn [9], slightly modified. In a haemolysis tube, 2500 µl of sample at 50 µg/ml (EWC or EC) and 500 µl of Folin Denis reagent were mixed. After 3 min incubation, 500 µl of 25% sodium carbonate was added before centrifuging the tube at 4000 rpm for 4 min. The absorbance was measured at 760 nm against a blank (methanol) using a Bio System BTS/340 UV-visible spectrophotometer. The dosage was done in triplicate (n = 3). A calibration range carried out with gallic acid at different concentrations (0; 5 ;10 ;15 ;20 µg/ml) was treated in the same way as the samples.

Polyphenols contents were expressed in mg gallic acid equivalent per gram of dry extract (mg GAE/g) according to the formula: $T = C.V/M$.

- T : Total polyphenol content expressed in mg EAG/g,
- C : Concentration of polyphenols established from the equation of the calibration line in mg/l,
- V : Volume of extract in l,
- M : Mass of the extract studied per g of extract.

Tannins contents represented the difference between total polyphenols (in EWC) and residual polyphenols (in EC).

2.2.3 Antioxidant activity

DPPH assay: The determination of the DPPH free radical scavenging activity of samples was done using the described method [10]. An ethanol solution of DPPH was prepared by dissolving 4 mg of DPPH in 100 ml of ethanol, followed by incubation between 4-8° for at least 16 hours. An aliquot of each sample (0.2 ml) at appropriate concentration was added to 0.8 ml of ethanol solution of DPPH. The two extracts (EWC and EC) were tested at different concentrations. The absorbance was measured at 517 nm after 30 min. Each experiment was done in triplicate.

The antioxidant activity related to the DPPH free radical scavenging effect was expressed as percentage of inhibition (IP) according to the following formula:

$$IP (\%) = 100 \times (A0 - A1) / A0.$$

- A0: absorbance of the control
- A1: absorbance of the sample

The IC₅₀ values (concentration of sample required to scavenge 50% of free radicals) IC₅₀ values were determined by plotting inhibition percentages versus concentrations IC₅₀ values were determined by plotting a percent inhibition versus concentration curve using the software Statgraphics Plus 5.0.

FRAP assay: The ferric reducing antioxidant power was determined according to the described method [11]. An aliquot of 0.20 ml of each sample (EWC and EC) at appropriate concentration was mixed with 0.5 ml of phosphate buffered saline (0.2 M; pH: 6.6) and 0.5 ml of 1% potassium ferricyanide (K₃Fe(CN)₆). The mixture was incubated at 50°C for 30 min and 0.5 ml of 10% trichloroacetic acid was added. After centrifugation for 10 minutes at 3000 rpm, the supernatant (0.5 ml) was mixed with distilled water (0.5 ml) and 0.1% ferric chloride (0.1 ml). The assays were performed three times with triplicate samples. Absorbance was measured at 700 nm. The reducing power (RP) expressed as percentage was obtained with the following formula:

$$RP (\%) = 100 \times (A0 - A1) / A0$$

- A0: absorbance of the control
- A1: absorbance of the sample

2.3 Statistical Analysis

The statistical analysis was done by Statview 4.5 software using the Fischer test. The difference was considered significant when $p < 0.05$ compared to the negative control.

3. RESULTS

3.1 Extraction

After extraction of 30 g of *C. nitida* seeds powder, a dry extract weighing 3.78 g was obtained, giving a yield of 12.6%.

3.2 Polyphenols and Condensed Tannins Contents

The total polyphenol content as well as the residual polyphenol content after precipitation of the tannins from the samples were calculated using the equation $y = 0.0341x + 0.0367$ ($R^2 = 0.9994$) from the calibration line obtained with gallic acid shown in Fig. 1. As shown in Fig. 2,

EWC had a total polyphenol content of 64.72 ± 0.63 mg GAE/g. After treatment with casein (EC), tannins were eliminated and the residual polyphenol content was 37.53 ± 0.52 mg EAG/g. Thus, condensed tannins content was 27.19 ± 0.11 mg GAE/g.

3.3 Antioxidant Activity

3.3.1 DPPH assay

The two samples (EWC and EC) of *Cola nitida* seeds significantly inhibit the DPPH radical ($p < 0.05$ versus negative control) at all concentrations tested and, in a dose dependent manner as shown Fig. 3.

The IC₅₀ values had confirmed that the extract not treated with casein ((IC₅₀: 5.54 ± 0.005 µg/ml) had shown better ability to scavenge the DPPH free radical than the extract treated with casein (IC₅₀: 61.92 ± 0.165 µg/ml) as illustrated in Fig. 4.

3.3.2 FRAP assay

The reducing activity of the hydro-ethanolic extract not treated with casein (EWC) and that of the hydro-ethanolic extract treated with casein (EC) are illustrated in Fig. 5. The results of the FRAP test confirmed those of the DPPH test. Indeed, as EWC had a greater reducing power than EC at all tested concentrations.




Fig. 1. Calibration line obtained with gallic acid




Fig. 2. Total polyphenol (in EWC), Residual polyphenol (in EC) and condensed tannins contents

EWC: Extract not treated with casein; EC: Extract treated with casein




Fig. 3. Inhibition percentage of samples on DPPH test

EWC: Extract not treated with casein; EC: Extract treated with casein; AA: ascorbic acid




Fig. 4. IC₅₀ of samples on DPPH test

EWC: Extract not treated with casein; EC: Extract treated with casein; AA: ascorbic acid




Fig. 5. Reducing percentage of different samples in FRAP test

EWC: Extract not treated with casein; EC: Extract treated with casein; AA: ascorbic acid

4. DISCUSSION

According to Odebode [12], *Cola nitida* seeds are rich in phenolic compounds such as condensed tannins and flavonoids. The extraction of polyphenolic compounds is a crucial step for the valorization of these active ingredients recognized for their high antiradical power [13]. It depends on the method used and the appropriate choice of solvent preserving the biological properties of these phenolic compounds [14].

As part of our study, we first performed a Soxhlet extraction of *Cola nitida* seeds powder using an ethanol-water mixture (80/20; v/v) as solvent. These two polar solvents are likely to extract heterosides compounds such as the polyphenols contained in the seeds of *Cola nitida*. The yield of this extraction was 12.6%. Adesanwo and al. [15] had found a higher yield (17.6 %) using methanol as solvent. With dichloromethane, they obtained a yield of 0.77%. These results suggest that the most represented active principles in *Cola nitida* seeds are mainly polar compounds.

The main objective of our study was to highlight the antioxidant activity of the tannins of the hydro-ethanolic extract of *Cola nitida* seeds by the DPPH and FRAP methods.

The DPPH method is based on the ability of an antiradical to stabilize the purple DPPH free radical by transforming it into pale yellow DPPH, H⁺, by trapping a proton [16]. As for the FRAP

method, it is based on the ability of a compound to reduce the ferric ion (Fe³⁺) to the ferrous one (Fe²⁺). The reducing capacity of ferric ion seems to be related to the degree of hydroxylation and extend to the conjugation in phenolic compounds [17].

The results obtained showed that EWC significantly inhibited the DPPH free radical and reduced ferric iron at all tested concentrations ($p < 0.05$). After treatment with casein (fixation and elimination of tannins), the antioxidant activity drops significantly for both the DPPH method and the FRAP method. Thus, for the DPPH test, at the highest concentration tested (10 µg/ml), the inhibition percentage (IP) of EWC was $71.63 \pm 0.06\%$. After treatment with casein (EC), the IP decreased to $17.21 \pm 0.05\%$ or 4.16 times less at the same concentration. The determination of the 50% inhibitory concentrations (IC₅₀) confirmed these results. Indeed, the IC₅₀ of EWC was only 5.54 ± 0.005 µg/ml. After treatment with casein, it increased 11.15 times, with an IC₅₀ value of 61.92 ± 0.165 µg/ml. The results of the FRAP test corroborate those of the DPPH method, with reducing powers of $90.06 \pm 0.18\%$ and $53.95 \pm 0.06\%$ for EWC and EC respectively. This means that the reducing power of EWC was 1.66 time greater than that of EC at the highest concentration tested (5,87 µg/ml).

The dosage of total polyphenols and residual polyphenols made it possible to deduce the

tannin content of the EWC. Thus, the content of total polyphenols in this extract was 64.72 ± 0.6 mg GAE/g. After fixation of the tannins condensed by casein, it was estimated at 37.53 ± 0.52 mg GAE/g. The content of condensed tannins is then 27.19 ± 0.26 mg EAG/g, ie 42.2% of the total polyphenols.

These results showed that *Cola nitida* seeds are rich in condensed tannins that play an important role in the antioxidant activity noticed for this drug. These results agree with those of Dieng and al. [18], who had proved that the condensed tannins of *Piliostigma reticulatum* (DC) Hochst, had a clear antioxidant activity. So condensed tannins could be used against oxidative stress to prevent several diseases. In fatty foods, sorghum condensed tannins are already used as antioxidant additives for their conservation [19]. The use of natural antioxidants in food could be a good alternative to chemical antioxidants. Indeed, although authorized, these chemical additives would have long-term toxic effects. Certain nanoparticles of chemical antioxidants could cross the placental barrier and cause teratogenic effects [20]. However, condensed tannins have the disadvantage of binding to proteins, carbohydrates and mineral elements, thus reducing nutritional value of certain nutrients [21].

5. CONCLUSION

These studies showed that the hydro-ethanolic extract of *Cola nitida* seeds (EWC) had good antioxidant activity. Rid of condensed tannins (EC), this activity decreased sharply. Condensed tannins contribute significantly to the antioxidant activity of *Cola nitida* seeds.

CONSENT AND ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Muanda FN. Identification de polyphenols, évaluation de leur activité antioxydante et étude de leurs propriétés biologiques. *Thèse Doct. Pharm., Metz*. 2010;239.
2. WHO. WHO Strategy for Traditional Medicine. 2013;2014-2023. ISBN: 978 92 4 250609 9 (classification NLM : WB 55)(www.who.int).
3. Babar A, Hahne J, Paek KV. Methyl jasmonate and salicylic acid induced oxidative stress and accumulation of phenolics in panax ginseng bioreactor root suspension cultures. *Molecules*. 2007;12(3):607–621.
4. Falleh H, Ksouri R, Chaieb K, Karray-Bouraoui N, Trabelsi N, Boulaaba M, Abdelly C. Phenolic composition of *Cynara cardunculus* L. organs and their biological activities. *C. R. Biologies*. 2008;331:372-379.
5. Gómez-Caravaca AM, Gómez-Romero M, Arráez-Román D, Segura-Carretero A, Fernández-Gutiérrez A. Advances in the analysis of phenolic compound in products derived from bees. *J Pharm Biomed Anal*. 2006;41(4):1220-34.
6. Ayensu ES. Medicinal plants of West Africa. 1st Edition. Algonac, Michigan: Reference Publications Inc. 1978;58–61:130, 201, 222.
7. Redfern J, Kinninmonth M, Burdass D, Verran J. Using soxhlet ethanol extraction to produce and test plant material (essential oils) for their antimicrobial properties. *J Microbiol Biol Educ*. 2014; 15(1):45–46.
8. Sarr A, Dieng SIM, Badji KD, Mbaye AI, Diatta W, Ka A, Fall AD, Bassène E. Phytochemical screening and determination of polyphenols in the hydro-ethanolic extract of trunkbark and its fractions of *Stereospermum kunthianum* cham (*Bignoniaceae*). *Asian Plant Research Journal* 2021;7(1):1-9.
9. Joslyn MA. Ash content and ashing procedures. In: Joslyn, M.A., Ed., methods in food analysis. Physical, Chemical and Instrumental Methods of Analysis, 2nd Edition. Academic Press, New York. 1970;109-140.
10. Nunes PX, Silva SF, Guedes JR, Tolentino J, Augusto L, Quintans-Junior LJ, Filho JMB. Biological oxidations and antioxidant activity of natural products. *Phytochemicals as nutraceuticals-Global Approaches to Their Role in Nutrition and Health*. 2012;278. DOI: 10.5772/2375
11. Bassène E. Initiation à la recherche sur les substances naturelles : Extraction, analyses, essais biologiques. Presses Universitaires, Dakar; 2012.

12. Odebode AC. Phenolic compounds in the kola nut (*Cola nitida* and *Cola acuminata*) (*Sterculiaceae*) in Africa. *Rev. Biol. Trop.* 1996;44(2):513-515.
13. Bonnaillie C, Salacs M, Vassilova E, Saykova I. Etude de l'extraction de composés phénoliques à partir de pellicules d'arachide (*Arachis hypogea* L.). *Revue de Génie Industriel.* 2012;7:35-45.
14. Mahmoudi S, Khali M, Mahmoudi N. Etude de l'extraction des composés phénoliques de différentes parties de la fleur d'artichaut (*Cynara scolymus* L.). *Nature & Technologie.* 2013;9: 35-40.
15. Adesanwo JK, Ogundele SB, Akinpelu DA, McDonald AG. Chemical analyses, antimicrobial and antioxidant activities of extracts from *Cola nitida* See. *Journal of Exploratory Research in Pharmacology.* 2017;2(3):67-77.
DOI: 10.14218/JERP.2017.00015
16. Prior RL, Wu X, Schaich K. Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements. *Journal of Agriculture and Food Chemistry.* 2005;53 (10):4290-303.
17. Benzie IF, Choi SW. Antioxidants in food: Content, measurement, significance, action, cautions, caveats, and research needs. *Advances in Food and Nutrition Research.* 2014;71:1-53.
18. Dieng SIM, Mathieu C, Sarr A, Diatta-Badji K, Fall AD. Condensed tannins content and their influence on the antioxidant activity of bark hydroethanol extract of *Piliostigma reticulatum* (Dc) hochst and its fractions. *Pharmacognosy Journal.* 2020; 12(2):361-368.
DOI: 10.5530/pj.2020.12.57
19. Sikwese FE, Duodu KG. Antioxidant effect of crude phenolic extract from sorghum bran in sunflower oil in the presence of ferric ions. *Food Chem.* 2007;104:324-331.
20. Guillard A, Gaultier E, Cartier C, Devoille L, Noireaux J, Chevalier L, Morin M, Grandin F, Lacroix MZ, Coméra C, Cazanave A, de Place A, Gayraud V, Bach V, Chardon K, Bekhti N, Adel-Patient K, Vayssière C, Fisticaro P, Feltin N, de la Farge F, Picard-Hagen N, Lamas B, Houdeau E. Basal Ti level in the human placenta and meconium and evidence of a materno-foetal transfer of food-grade TiO₂ nanoparticles in an ex vivo placental perfusion model. *Part. Fibre Toxicol.* 2020;1751.
Available:https://doi.org/10.1186/s12989-020-00381-z
21. Ojo MA. Tannins in foods: Nutritional implications and processing effects of hydrothermal techniques on underutilized hard-to-cook legume seeds—a review. *Prev Nutr Food Sci.* 2022; 27(1):14–19.

© 2022 Sarr et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/93755>