



Effect of Irrigation Level and Nitrogen Levels on Economics, Quality and Water Productivity of Garden Cress (*Lepidium sativum* L.)

Nupur Sharma^{a++*}, Vinod Kumar Yadav^{a#}
and Manisha Jangir^{bt}

^a Krishi Vigyan kendra, Sawaimadhopur, Agriculture University, Kota, India.
^b SKN Agriculture University, Jobner, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i22751

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/96783>

Original Research Article

Received: 23/10/2022
Accepted: 30/12/2022
Published: 06/02/2023

ABSTRACT

A field experiment was conducted at krishi vigyan kendra, Sawaimadhopur, Rajasthan in two consecutive years of 2018-19 and 2019-20 on "Effect of irrigation level and nitrogen levels on economics, quality and water productivity of chandrasur (*L. sativum*)". The highest economics net return was found with three irrigations [25, 50 & 75 days after sowing (DAS)] I₃ (57517 Rs ha⁻¹) which was at par with I₂ with two irrigations (25 & 50 DAS) in the pooled data of both years. Highest benefit-cost (B/C) ratio was also found under three irrigations (25, 50 & 75 DAS) I₃ (2.60) which was at par with (I₂) two irrigations (25 & 50 DAS) in the pooled data of both years. Highest net return and

⁺⁺ Subject Matter Specialist (Agronomy);

[#] Assistant Professor (Soil Science);

[†] P.H.D. Research Scholar;

*Corresponding author: E-mail: nupurmahershi@gmail.com;

B/C ratio was found with application of 80 Kg N ha⁻¹ (57628 Rs ha⁻¹) and (2.65) which was at par with 60 Kg N ha⁻¹. In terms of oil yield Kg ha⁻¹ highest oil yield was found with (I₂) two irrigations (25 & 50 DAS) and 60 Kg N ha⁻¹. Water productivity (kg m⁻³) was found lowest in the treatment (25, 50 & 75DAS) (I₃) and 80 Kg N ha⁻¹ which was at par with (I₂) two irrigations (25 & 50 DAS) and 60 Kg N ha⁻¹.

Keywords: Garden cress; Irrigation level; *Lepidium sativum*; nitrogen levels; oil yield; water productivity.

1. INTRODUCTION

The “chandrasur” *Lepidium sativum* L., Brassicaceae) is also known with the common name “Garden cress”. It is an annual herbaceous plant native of Asia and north Africa, but distributed. It is very nutritious which can use as green leafy vegetables as salad purpose. It can also use to increase traditionally height of childrens and milk in lactating women. It is used to treat liver diseases and infectious diseases, as immune system enhancer, antibacterial, etc. [1]. Due to its properties it can be used as insect repellent and to treat insect bites. In India it is cultivated mostly Madhya Pradesh, Rajasthan, Uttar Pradesh etc. In Rajasthan it is cultivated as fodder for animals. Chandrasur is a crop withlow water requirements. However, irrigation scheduling at specific time is very important for the maximum yield of crop.

Fertilization is very important for the mostly agricultural crop production. When we applied in injudiciously is can make loss and pollution of water also. Excessive application losses results in emission of green house gases nitrogen oxides [2].

Nitrogen fertilization plays a very important role in structural and yield component of the plants. In plants many enzymatic activities and chlorophyll and other structural components depends upon nitrogen [3]. Chandrasuris considered a species of high nitrate accumulating (Cavarianni et al. 2008). Vegetable plants require high amount of nitrogen because its vegetation period is short [4]. That's why there is a need to find out proper irrigation schedule and nitrogen level for the chandrasur for zone III B of Rajasthan.

2. MATERIALS AND METHODS

The study was conducted at the agronomic farm of Krishi Vigyan Kendra, Sawai madhopur, Agriculture University, Kota. The soil of experimental site was sandy loam soil,

moderately fertile, low in available organic carbon 0.30 %, low in available nitrogen 243 kg ha⁻¹, medium in available phosphorus 20.52 kg ha⁻¹, and high in available potash 279 kg ha⁻¹. Sawai madhopur consist in Zone III-B of Rajasthan (Flood Prone Eastern Plain) and Zone-V of Rajasthan (Sub Humid South- Eastern Plain) of Rajasthan. In the zone III-B the mean annual rainfall is 623 mm. Thechandrasur sowing was done in the second week of October during both the years. The experiment was laid out in split plot design in which three treatments of irrigation levels were done (1) one irrigation at 25 days after sowing (DAS) (I₁), (2) two irrigations at 25 & 50 DAS) (I₂), (3)three irrigations (25, 50 & 75 DAS) (I₃) and were placed in main plot. Four nitrogen levels (20kg N ha⁻¹, 40kg N ha⁻¹, 60kg N ha⁻¹, 80kg N ha⁻¹) were placed in sub plots. Sowing was done at spacing of 30×10 cm and 6 kg ha⁻¹ seed rate was used during experiment. Main irrigation channel was 1.5 m wide and sub irrigation channel was 1.0 m wide. Different nitrogen levels were applied through urea in two split doses first basal application and second after first irrigation 25 DAS. Full dose of phosphorus was applied as basal in the form of SSP. For irrigation was used rain water. Net return and benefit-cost ratio (B/C ratio) was calculated after the harvesting of crop on the basis of following formula:

Net return = Gross return-Cost of Cultivation

B/C ratio = Net return / Cost of cultivation

Oil content (kg ha⁻¹) was analysed in laboratory of Agriculture University, Kota.

Water productivity was measured by following formula:

Water Productivity (kg m⁻³)= Grain yield (kg ha⁻¹)/ total water applied (m⁻³)

The statistical calculation were done by as per analysis of variance described by Gomez and Gomez 1984 to analysis the test of significant of treatments. Gomez KA. and Gomez AA [5].

3. RESULTS AND DISCUSSION

3.1 Oil Yield (Kg ha⁻¹)

Data presented in Table 1 revealed that the highest yield was obtained with three irrigations (25, 50 & 75 DAS) (I₃) 353.91 kg ha⁻¹ which was at par with two irrigation (25 & 50 DAS) (I₂). Application of 80 kg N ha⁻¹ found highest Oil yield 344.20 kg ha⁻¹ which was found statistically at par with 60 kg N ha⁻¹ 328.92 kg ha⁻¹. With increasing the number of irrigations the oil content in chandrasur will increase significantly up to a certain level. Similar results was found by [6] explained that by increasing the irrigation interval the oil percentage of chandrasur will increase to its maximum values. The result of this study also stated that increasing level of nitrogen the oil yield in seeds of chandrasur also increase. These results are in agreement with those establishing that fertilization increases the fatty acid presence, percentages and the fatty acids ratio in canola seed oil [7,8].

3.2 Water Productivity (kg m⁻³)

The results presented in Table 1 that during both years the highest water productivity (kg m⁻³) was found with three irrigations (25, 50 & 75 DAS) (I₃) 1.06 kg m⁻³ which was at par with two Irrigation (25 & 50 DAS) (I₂) 1.47 kg m⁻³. Same trend was found during each year. These results are in close conformity with Shivran et al. [9] who reported that increasing no. of irrigations up to three consumption of water was also increases. Increased no. of irrigation would trend to higher consumption of water due to higher seed yield.

Table 1 shows that the highest water productivity was found with application of 80 kg N ha⁻¹ 1.86 kg m⁻³ which was significantly higher over 20 kg N ha⁻¹ and 40 kg N ha⁻¹. The results confirm the findings of Pradhan et al. [10] who reported the water productivity significantly increases with the level of Nitrogen. The higher water productivity was due to higher grain yield with efficient utilization of water per drop of water. Similarly Tadayon et al. [11] reported that highest water productivity was obtained with application of 100 Kg N ha⁻¹ which was significantly higher over 0 and 50 kg N ha⁻¹.

3.3 Nitrogen Content in Grains (%)

The results presented in Table 2 show that highest content of nitrogen in grains was found under three irrigations (25,50 & 75 DAS) (I₃), 1.42% which was significantly higher over one irrigations (25 DAS) and at par with two irrigation (I₂) (25 & 50 DAS) 1.37 %. This might be due to adequate moisture content helps to increase the nutrient content and uptake in plants. These results are in close conformity with Umale et al. [12].

In terms of application of nitrogen fertilization highest nitrogen content in grains was found with application of 80 kg N ha⁻¹ which was at par with 60 kg N ha⁻¹ and significantly higher over 20 & 40 Kg N ha⁻¹. These results are in close conformity with Inne et al. [13] who stated that the highest content was found in application of highest amount of nitrogen 150 kg ha⁻¹.

Table 1. Effect of Irrigation and Nitrogen levels on Oil yield (Kg ha⁻¹) and Water productivity (kg m⁻³) of Chandrasur (*Lepidium sativum*)

Treatments	Oil yield (Kg ha ⁻¹)			Water productivity (kg m ⁻³)		
	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled
Irrigation levels						
Irrigation (One at 25 DAS) (I ₁)	263.53	271.83	267.68	2.49	2.57	2.53
two Irrigation (25 & 50 DAS) (I ₂)	320.77	324.92	322.85	1.41	1.53	1.47
Three irrigation (25, 50 & 75 DAS) (I ₃)	351.82	355.99	353.91	1.06	1.07	1.06
SEm±	10.23	9.14	6.86	0.09	0.08	0.06
CD (P=0.05)	35.40	31.63	21.14	0.33	0.29	0.20
Nitrogen levels						
20kg N ha ⁻¹ (N ₁)	280.42	290.48	285.45	1.50	1.58	1.54
40 kg N ha ⁻¹ (N ₂)	299.42	301.93	300.68	1.56	1.66	1.61
60 kg N ha ⁻¹ (N ₃)	327.75	330.10	328.92	1.73	1.75	1.74
80 kg N ha ⁻¹ (N ₄)	340.57	347.83	344.20	1.83	1.90	1.86
SEm±	6.66	8.10	5.24	0.06	0.06	0.04
CD (P=0.05)	19.33	23.51	14.87	0.16	0.17	0.11

3.4 Nitrogen Uptake in Grains (Kg ha⁻¹)

Results in Table 2 represented that highest nitrogen uptake in grains (kg ha⁻¹) was found in three irrigations (25, 50 & 75 DAS) (I³) 22.71 kg ha⁻¹ which was at par with two irrigations (25 & 50 DAS) (I²) 20.18 kg ha⁻¹ and significantly higher over one Irrigation (25 DAS). Razmjoo et al. [14] reported that scarcity of water is the main factor in low yield and productivity of medicinal plants. Deficiency of moisture induces many different physiological and metabolic responses like stomata closure, decrease in growth and photosynthesis. This might also leads to decrease in nutrient uptake and finally affects yield of crops. These results are in close conformity with Jibrin et al. [15].

In different nitrogen levels the highest nitrogen uptake 23.67 kg ha⁻¹ was found with application

of 80 kg N ha⁻¹ which was significantly higher over 20 and 40 kg N ha⁻¹ and at par with 60 kg N ha⁻¹ (20.62 kg ha⁻¹). These results are closely related to those reported by Raghuvanshi et al. [16] who stated that with higher nitrogen fertilization uptake of nitrogen by seed and strover was also increases in mustard [*Brassica juncea* (L.) Czern.]. At higher levels of fertilizers assures higher accessibility of nutrient in adequate amount for the plant to uptake [17].

3.5 Net Return (Rs ha⁻¹)

Data presented in Table 3 elaborated that in pooled data of both the years highest Net return (Rs ha⁻¹) and B/C ratio were found with three irrigations (25, 50 & 75 DAS) (I³) 57517 Rs ha⁻¹ and 2.60 which was at par with two irrigations (25 & 50 DAS) (I²) 51699 Rs ha⁻¹ and 2.38 respectively.

Table 2. Effect of Irrigation and Nitrogen levels on Nitrogen content in grains and Nitrogen uptake in grains (kg ha⁻¹) in Chandrasur (*Lepidium sativum*)

Treatments	Nitrogen content in grains (%)			Nitrogen uptake in grains (kg ha ⁻¹)		
	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled
Irrigation levels						
Irrigation (One at 25 DAS) (I ₁)	1.23	1.31	1.27	15.44	17.00	16.22
two Irrigation (25 & 50 DAS) (I ₂)	1.33	1.41	1.37	18.72	21.64	20.18
Three irrigation (25, 50 & 75 DAS) (I ₃)	1.37	1.47	1.42	21.84	23.58	22.71
SEm±	0.02	0.02	0.02	0.63	0.52	0.41
CD (P=0.05)	0.07	0.08	0.05	2.20	1.78	1.26
Nitrogen levels						
20kg N ha ⁻¹ (N ₁)	1.18	1.26	1.22	15.32	17.28	16.30
40 kg N ha ⁻¹ (N ₂)	1.28	1.36	1.32	17.10	19.34	18.22
60 kg N ha ⁻¹ (N ₃)	1.34	1.43	1.38	19.71	21.53	20.62
80 kg N ha ⁻¹ (N ₄)	1.44	1.54	1.49	22.53	24.82	23.67
SEm±	0.04	0.04	0.03	0.72	0.83	0.55
CD (P=0.05)	0.10	0.11	0.07	2.09	2.40	1.55

Table 3. Effect of Irrigation and Nitrogen levels on Net returns (Rs ha⁻¹) and B/C ratio of chandrasur (*Lepidium sativum*)

Treatments	Net return (Rs ha ⁻¹)			B/C ratio		
	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled
Irrigation levels						
Irrigation (One at 25 DAS) (I ₁)	42713	43657	43185	2.17	2.11	2.14
two Irrigation (25 & 50 DAS) (I ₂)	49272	54125	51699	2.33	2.44	2.38
Three irrigation (25, 50 & 75 DAS) (I ₃)	57547	57487	57517	2.66	2.54	2.60
SEm±	2621	2213	2341	0.13	0.10	0.12
CD (P=0.05)	9069	7657	9357	0.45	0.36	0.41
Nitrogen levels						
20kg N ha ⁻¹ (N ₁)	43910	46575	45242	2.14	2.16	2.15
40 kg N ha ⁻¹ (N ₂)	45970	48826	47398	2.21	2.24	2.23
60 kg N ha ⁻¹ (N ₃)	52613	53251	52932	2.51	2.42	2.46
80 kg N ha ⁻¹ (N ₄)	56882	58375	57628	2.68	2.63	2.65
SEm±	2212	2023	2176	0.11	0.09	0.07
CD (P=0.05)	6419	5872	6745	0.31	0.27	0.20

In terms of application of different nitrogen levels the highest net return and B/C ratio were found with application of 80 Kg N ha⁻¹ (57628 Rs ha⁻¹ and 2.65) which was at par with 60 kg N ha⁻¹ (52932 Rs ha⁻¹ and 2.46) and significantly higher over 20 & 40 Kg N ha⁻¹.

Highest net return and B/C ratio was obtained with two irrigation and 60 kg N ha⁻¹. This might be due to highest seed yield and optimum use of nutrient applied through fertilization. With two irrigations (25 & 50 DAS) availability of moisture was sufficient to crop growth and it gave higher yield which was at par with three irrigations. Availability of moisture also helps to enhance the accessibility and uptake of nutrient by the plants. Ultimate results was higher yield. These results are in close conformity with [18,17].

4. CONCLUSION

On the basis of the results emanated from present investigation conducted during Rabi 2018-19 and 2019-20, Based on two year study, results revealed that application of two irrigations (25 & 50 DAS) with application of 60 Kg N ha⁻¹ gives higher net return, BC ratio. Higher water productivity was also found under two irrigations and application of 60 kg N ha⁻¹.

ACKNOWLEDGMENT

I acknowledge my sincere thanks to Dr. Vinod kumar Yadav (Assistant Professor, soil science) and Mrs. Yamini Tak (Assistant Professor, Biochemistry, Agriculture university, Kota) for helping me in soil and plant sample analysis and guidance and support during my experiment.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Amin GH. Medicinal Plants of Iran, Tehran University Publication. 2005;106.
2. Güler S. Dünya'da ve Türkiye'de Gübre Tüketiminde Yaşanan Gelişmeler. In: Karaman MR, Brohi AR. (eds). Türkiye 3. Ulusal Gübre Kongresi, Tarım-Sanayi-Çevre, 11-13 Ekim, Tokat. 2004; 47-54.
3. Laza MR. Re: Calculation of Water Efficiency and Productivity?; 2017. Available: https://www.researchgate.net/post/Calculation_of_water_efficiency_and_productivity/58f55f08dc332dd26542b67a/citation/download
4. Maynard DN, Barker AY, Minotti PL, Peck NH. Nitrate accumulation in vegetables. Analysis in Agronomy. 1976;28:71.
5. Gomez KA, Gomez AA. Statistical Procedures for Agricultural Research. A Willey Inter science Publication, John Willy & Sons, New York. 1984:317-356.
6. Khalil SE, El-Noemani AA. Effect of irrigation intervals and exogenous proline application in improving tolerance of garden cress plant (*Lepidium sativum* L.) to water stress. Journal of Applied Sciences Research. 2012;8(1):157-167.
7. Gao J, Thelen KD, Min D, Smith S, Hao X, Gehl R. Effects of manure and fertilizer applications on canola oil content and fatty acid composition. Agron J. 2010;102:790–797.
8. Mohamed EAA, Muddathir AM, Abdalla AH. Effects of organic and inorganic fertilization on growth, yield, seed fixed oil content, and fatty acids profile of garden cress (*Lepidium sativum* L.). SN Applied Sciences. 2020;2:1753.
9. Shivran H, Kumar S, Tomar R, Chauhan GV. Effect of irrigation schedules on productivity and water use efficiency in Indian mustard (*Brassica juncea* L.). International Journal of Chemical Studies. 2018;6(4):15-17.
10. Pradhan S, Chopra UK, Bandyopadhyay KK, Singh R, Jain AK, Chand I. Effect of water and nitrogen management on water productivity and nitrogen use efficiency of wheat in a semi-arid environment. International Journal of Agriculture Food Science Technology. 2013;4(7):727-32.
11. Tadayon MR, Ebrahimi R, Tadayyon A. Increased water productivity of wheat under supplemental irrigation and nitrogen application in a semi-arid region. Journal of Agriculture Science Technology. 2012;14:995-1003.
12. Umale AA, Patel JJ, Kumar M, Ahmad S, Rabari KV. Influence of dates of sowing and irrigation scheduling (IW:CPE ratios) on growth, yield attributes and seed yield of Cress (*Lepidium sativum* L.) . Ecology. Environment & Conservation. 2015;S17-S21.
13. Inne A, Kul R, Ekinçi M, Turan M, Yildirim E. Nitrogen Fertilization Affects Growth, Yield, Nitrate and Mineral Content of

- Garden Cress (*Lepidium sativum* L.). Yuzuncu Yil University Journal of Agricultural Science. 2021;31(1).
14. Razmjoo K, Heydarizadeh P, Mohammed R. Effect of salinity and drought stresses on growth parameters and essential oil content of *Matricaria chamomile*. International Journal of Agriculture and Biology. 2008;4:451 – 454.
 15. Jibrin DM, Namakka A, Ibrahim DA. Response of garden cress (*Lepidium sativum* L.) To sowing methods, irrigation interval and fertilizer rates in northern Guinea savannah of Nigeria. Journal of Agriculture and Environment. 2018;14 (1):93-98.
 16. Raghuvanshi N, Kumar V, Dev J. Effect of Nitrogen Levels on Mustard (*Brassica juncea* (L.) Czern and Coss.) Varieties under Late Sown Condition. Current Journal of Applied Science and Technology. 2018;30(2): 1-8.
 17. Nayak H, Bohra JS, Yadav SP. Growth and nutrient uptake of indian mustard [*Brassica juncea* (L.) Czern and Coss.] genotypes as influenced by nitrogen and sulphur fertilization under irrigated condition. *Ecology. Environment and Conservation*. 2020:S79-S83.
 18. Saraswathi G, Vidya KM, Hedge L, Chavan ML, Kumar BMV. Effect of dates of sowing and fertilizer levels on nutrient uptake and economics of garden cress (*lepidium sativum* L.). Plant Archives. 2014;14 (1):327-330.
 19. Choudhary MS, Choudhary M, Choudhary R, Choudhary S, Choudhary P, Dhayal S. Effect of Fertility Levels on Growth and Yield of Chandrasur (*Lepidium sativum* L.) Varieties under Southern Rajasthan Condition. Annals of Agricultural Research. 2022;43(3):340-347.

© 2023 Sharma et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/96783>