



Negative Appendectomy Rates - Current Prospective- A Review

H. R. Kumar ^{a++*} and M. Soma ^a

^a *Department of Surgery, Taylor's University School of Medicine and Health Science, Subang Jaya, Selangor- 47500, Malaysia.*

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJMAH/2023/v21i10873

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/102421>

Systematic Review Article

Received: 05/05/2023

Accepted: 06/07/2023

Published: 15/07/2023

ABSTRACT

The negative appendectomy rate is generally accepted in the diagnosis of acute appendicitis to prevent complications like perforation. This rate varies according to countries in western and eastern regions of the world. Clinical scoring systems, inflammatory markers and imaging have been used to reduce the negative appendectomy rate. The introduction of computerized tomography has improved the diagnostic accuracy of acute appendicitis. As this rate has been progressively decreasing over the past few years, a review article was done to look at the role of scoring systems, inflammatory markers, and imaging in diagnosing acute appendicitis and hence reduce the negative appendectomy rate.

Keywords: Negative appendectomy; white appendix; clinical scoring system; appendectomy.

1. INTRODUCTION

Acute appendicitis is a common cause of patients presenting to the emergency department

for symptoms of right lower abdominal pain and its incidence is 5.7 to 50 patients per 100,000 in developed countries [1].

⁺⁺ Associate Professor;

*Corresponding author: E-mail: kharirajah@yahoo.com.my;

The global incidence of acute appendicitis worldwide is 233 patients per 100,000 population with a lifetime risk of 6.7% to 8.6% [2].

Negative appendectomy is defined as the final pathological report after appendectomy which shows a normal, congested appendix without inflammation. The negative appendectomy rate ranges from 20 -25% but with the use of per-operative imaging the rate has reduced to 10%. The factors that increase the negative appendectomy rate are female patients, patients who are younger than 40 years and patients with a history of diarrhea. Negative that decreases the negative appendectomy rate are presence of leukocytosis and positive appendicitis, on ultrasound and computerized tomography [3].

Negative appendectomy is often an indicator of the quality of management of acute appendicitis. As the main aim is to prevent complications like perforated appendicitis, there is a trend for diagnosing acute appendicitis early and proceeding with appendectomy and this has resulted in a higher negative appendectomy rate [4].

A review article was done to look at the current incidences of negative appendectomy, the factors that affect negative appendectomy and reduce it. A literature review was made on PubMed, Cochrane database of clinical reviews and google scholar to look for original articles, clinical trials, observational and cohort studies, clinical reviews, and review articles from 1996 to 2023. The following keywords were used, "negative appendectomy", "white appendix", "acute appendicitis", and "appendectomy". All articles were in English language and adults and children were included in the study. Case write ups and commentaries were excluded from this review. Pregnant patients who were presented with acute appendicitis were also excluded from this review.

2. NEGATIVE APPENDECTOMY DEFINITION AND RATE

There is no widely accepted term for the definition of negative appendectomy, the most common definition includes a proportion of macroscopically and or histologically normal appendix with no pathological infiltration of the mucosa by polymorphonuclear leucocytes or lymphocytes. This definition is used for cases that have undergone appendectomy [5].

Another definition of negative appendectomy is the presence of a normal looking appendix during laparoscopic appendectomy for patients who present with right lower abdominal pain. This is still the standard practice in some hospitals [6].

The negative appendectomy rates vary according to the region with a range of 18-25%. In certain regions like in Africa, the negative appendectomy rate can vary from 16% to 35% depending on the region [7-9].

In Asia the reported negative appendectomy rates also vary according to the region, like in India where the rate 23.7%. Where else in Iran the negative appendectomy rate is about 20%. Certain countries like Oman in central Asia have reported a negative appendectomy rate of 12.23%. In the East Asia the negative appendectomy rate is about 18.2% as reported in Hong Kong [10-13].

The negative appendectomy rates in western countries also vary with the rate being 19% to 33% in the United Kingdom [14,15].

The Right Iliac Fossa Treatment (RIFT) study noted that the negative appendectomy rate in the United Kingdom was 28.2% in females and 12.1% in males. The rate among both males and females in other European countries like Italy, Portugal and the republic of Ireland were less than 10% [16].

The Dutch prospective nationwide outcome audit of surgery of suspected acute appendicitis showed that the negative appendectomy rate in the Netherlands was 2.3% [17].

The conclusion from these studies is that the negative appendectomy rates vary according to various regions of the world. There are multiple factors that affect the negative appendectomy rate.

3. CLINICAL SCORING SYSTEMS AND NEGATIVE APPENDECTOMY RATE

The Alvarado score is one of the most common scoring systems that has been used to reduce the negative appendectomy rate. A score of 7 or more can reduce the negative appendectomy rate and this was retrospectively assessed by Tekeli et al. [18].

The Alvarado score was also prospectively evaluated by bouali et al, and they noted that an Alvarado score of 7 or more was associated with

a better diagnosis of acute appendicitis and a negative appendectomy rate of 4.8%. A cross sectional study by memon et al also confirmed this but the negative appendectomy rates were reduced in males and not female patients [19,20].

The Modified Alvarado score and the Raja Isteri Pengiran anak Saleha appendicitis (RIPASA) score were compared in a retrospective study [21]. A score of 7 and above for the modified Alvarado score and 7.5 and above for the Raja isteri Pengiran anak Saleha appendicitis score was associated with a reduction in the negative appendectomy rate. A prospective study also concluded that the use of both the modified Alvarado score and the RIPASA score was associated with a reduction in the negative appendectomy rate [22].

The raja isteri Pengiran anak Saleha appendicitis (RIPASA) score was prospectively evaluated and a cut of score of 7.5 and above in patients with symptoms of acute appendicitis were associated with a better diagnosis and a reduced negative appendectomy rate [23].

The Adult Appendicitis Score (AAS) was prospectively evaluated and a score of 10 and above was associated with a reduction of the negative appendectomy rate to 8.7% [24].

The Appendicitis Inflammatory Response Score (AIR) was evaluated in a cross-sectional study and a score of 8 and above was associated with a better diagnosis of acute appendicitis and a reduction of the negative appendectomy rate [25,26].

In the pediatric population a prospective study was conducted on the clinical scoring systems used in reducing the negative appendectomy rates, and the conclusion was that all the scoring systems were effective in reducing this rate. The negative appendectomy rate was 14.8% in this study [27].

The conclusion from these studies were that clinical scoring systems were effective in the triage of patients who present with suspected acute appendicitis and were effective in reducing the negative appendectomy rate.

4. INFLAMMATORY MARKERS AND NEGATIVE APPENDECTOMY RATE

The common inflammatory markers include the total white cell count, C-reactive protein,

neutrophil to lymphocyte ratio and platelet to lymphocyte ratio were retrospectively assessed by yazar et al who concluded that when these markers were used with clinical examination, they improved the diagnostic accuracy of acute appendicitis and hence reduced the negative appendectomy rate [28].

Most inflammatory markers are useful in the diagnosis of acute appendicitis but are not superior to clinical examination and the duration of symptoms are important to establish the diagnosis and reduce the negative appendectomy rate [29,30].

A systemic review by acharya et al on the biomarkers of acute appendicitis showed that the use of biomarkers in conjunction with clinical examination will enhance the diagnostic accuracy of acute appendicitis and hence reduce negative appendectomy rate [31].

Most newer biomarkers like the platelet to lymphocyte ratio, the monocyte to lymphocyte ratio is showing promise in diagnosing acute appendicitis but further studies may be needed to see if it can reduce the negative appendectomy rate [32,33].

The conclusion from all these studies is that biomarkers are useful in aiding the diagnosis of acute appendicitis when it is combined with clinical examination and imaging. When it is used on its own it will not reduce the negative appendectomy rate.

5. IMAGING AND NEGATIVE APPENDECTOMY

The use of imaging modalities like ultrasound and computerized tomography have improved the diagnostic accuracy and reduced the negative appendectomy rate [34]. The use of ultrasound to diagnose acute appendicitis is useful as it involves any use of radiation, and it can be done in the emergency department. This retrospective study showed its sensitivity was 89.6% and its specificity was 93.8% [35].

A meta-analysis by Jian Fu et al on the diagnostic accuracy of abdominal ultrasound in the diagnosis of acute appendicitis concluded that the sensitivity was 77.2% and specificity was 60%. Ultrasound was cheap, non-invasive and does not involve the use of ionizing radiation and hence it is useful in diagnosing acute appendicitis [36].

Table 1. The various negative appendectomy rates in various countries

Study	Country	N=numbers	Negative appendectomy rate (NAR)	Study type
Grossberg S et al.	South Africa	1217	19%	Retrospective study
Malekpour et al.	Iran	1454	20.4%	Retrospective study
Sharma et al.	India	118	23.72%	Retrospective study
Chaochankit et al.	Thailand	892	10%	Retrospective study
Nyamuryekurg E et al.	Tanzania	91	38.5%	Retrospective study
Bangu et al.	United Kingdom	5345	12.2%-28.2%	RIFT study-retrospective
Van rossem et al.	Netherlands	1975	3.3%	Prospective-observational audit

A retrospective study on The use of ultrasound in the diagnosis of acute appendicitis in the pediatric population concluded that the sensitivity was 82% and its specificity was 97% and it should be the imaging of choice in children regardless of gender [37].

However, in cases of indeterminate examination of patients with suspected appendicitis, following ultrasound, computerized tomography was the investigation of choice to diagnose acute appendicitis. The use of computerized tomography has seen a reduction in the negative appendectomy rate [38-40].

Computerized tomography is the best imaging modality to diagnose acute appendicitis and a retrospective cross- sectional study showed that the negative appendectomy rate of 2.66% after imaging against a rate of 13.16% of patients who were diagnosed clinically [41].

A retrospective study on the role of preoperative computerized tomography in patients with acute appendicitis found that the negative appendectomy rate was 4.7% against 12.7% against those who underwent no imaging. This study confirmed that the negative appendectomy rate was the same in both sexes [42].

A prospective study on the use of computerized tomography in patients with suspected appendicitis was associated with a negative appendectomy rate of 6.4% against 19% in patients who did not undergo any imaging. This showed the sensitivity of computerized tomography in diagnosing acute appendicitis [43].

Further multi center cross sectional study on the use of computerized tomography in the diagnosis of acute appendicitis, showed that the negative appendectomy rate was 4.1% and this showed

the effective diagnostic capability of this imaging modality [44].

The conclusion from these studies was that ultrasound and computerized tomography were very effective in reducing the negative appendectomy rates in cases of suspected acute appendicitis.

6. CONCLUSION

As the negative appendectomy varies between 10% to 25%, and with better use of clinical scoring systems, blood investigations and imaging has led to improved rates. The use of imaging like Computerized tomography has increased the diagnostic accuracy of acute appendicitis but its limiting factor is its cost, use of ionizing radiation and availability of radiologist to report these images. In Asian countries where cost is an issue, the use of clinical scoring systems may be effectively used to triage patients who present with suspected appendicitis and decide which patients will require imaging. This may be the best method to decrease the negative appendectomy rate.

CONSENT AND ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Di Saverio S, Podda M, De Simone B, Ceresoli M, Augustin G, Gori A, et al. Diagnosis and treatment of acute appendicitis: 2020 update of the WSES Jerusalem guidelines. World Journal of

- Emergency Surgery. BioMed Central Ltd. 2020;15.
DOI-10.1186/s13017-020-00306-3
2. Guan L, Liu Z, Pan G, Zhang B, Wu Y, Gan T et al. The global, regional, and national burden of appendicitis in 204 countries and territories, 1990–2019: A systematic analysis from the global burden of disease study 2019. *BMC Gastroenterol.* 2023 Dec 1;23(1). DOI-10.1186/s12876-023-02678-7
 3. Chaochankit W, Boochoa A, Samphao S. Negative appendectomy rate in patients diagnosed with acute appendicitis. *BMC Surg.* 2022 Dec 1;22(1). DOI-10.1186/s12893-022-01852-0
 4. Li J. Revisiting delayed appendectomy in patients with acute appendicitis. *World Journal of Clinical Cases.* Baishideng Publishing Group Co. 2021;9:5372–90. DOI-10.12998/wjcc.v9.i20.5372
 5. Noureldin K, Hatim Ali AA, Issa M, Shah H, Ayantunde B, Ayantunde A. Negative appendectomy rate: Incidence and predictors. *Cureus.* 2022 Jan 22. DOI-10.7759/cureus.21489
 6. Lee M, Paavana T, Mazari F, Wilson TR. The morbidity of negative appendectomy. *Ann R Coll Surg Engl.* 2014;96(7):517–20. DOI:10.1308/003588414X13946184903801
 7. Pooria A, Pourya A, Gheini A. Appendicitis: Clinical implications in negative appendectomy. *International Journal of Surgery Open.* 2021 Feb 1;29:45–9. DOI-10.1016/j.ijso.2021.01.004
 8. Osime O, Ajayi P. Incidence of negative appendectomy: Experience from a company hospital in Nigeria. *Cal J Emerg Med.* 2005 Oct;6(4):69-73. PMID: 20847869; PMCID: PMC2906999.
 9. Nyamuryekung'e MK, Athar A, Patel MR, Njau A, Sherman O, Jusabani A et al. Negative appendectomy rate in urban referral hospitals in Tanzania: A cross-sectional analysis of associated factors. *Annals of African Surgery.* 2021 Apr 1;18(2):109–14. DOI-10.4314/AAS.V18I2.9
 10. Sharma R, Kasliwal DK, Sharma RG. Evaluation of negative appendectomy rate in cases of suspected acute appendicitis and to study the usefulness of ultrasonography in improving the diagnostic accuracy. *Indian J Surg.* 2007 Oct;69(5):194-7. DOI: 10.1007/s12262-007-0020-6. Epub 2008 Jul 30. PMID: 23132981; PMCID: PMC3452587.
 11. Malekpour N, Basharat S, Bakhshi T. Negative appendectomy rate during 5 years in modarres hospital of Tehran, Iran, and correlation with imaging. *Journal of Analytical Research in Clinical Medicine.* 2018 Jun 10;6(2):98–101. DOI-10.15171/jarcm.2018.015
 12. Malik KA. Negative appendectomy rate in Sultan Qaboos University Hospital, Oman. *Journal of Surgery.* 2013;1(3):43. DOI-10.11648/j.js20130103.11
 13. Ma KW, Chia NH, Yeung HW, Cheung MT. If not appendicitis, then what else can it be? A retrospective review of 1492 appendectomies. *Hong Kong Med J.* 2010 Feb;16(1):12-7. PMID: 20124568.
 14. Mariadason JG, Wang WN, Wallack MK, Belmonte A, Matari H. Negative appendectomy rate as a quality metric in the management of appendicitis: Impact of computed tomography, Alvarado score and the definition of negative appendectomy. *Ann R Coll Surg Engl.* 2012 Sep;94(6): 395–401. DOI:10.1308/003588412X13171221592131. PMID: 22943328; PMCID: PMC3954319.
 15. Lim J, Pang Q, Alexander R. One year negative appendectomy rates at a district general hospital: A retrospective cohort study. *International Journal of Surgery.* 2016 Jul 1;31:1–4. DOI:10.1016/j.ijso.2016.05.030. Epub 2016 May 10. PMID: 27174508.
 16. Bhangu A, Nepogodiev D, Matthews JH, Morley GL, Naumann DN, Ball A, et al. Evaluation of appendicitis risk prediction models in adults with suspected appendicitis. *British Journal of Surgery.* 2020;73–86. DOI: 10.1002/bjs.11440. Epub 2019 Dec 3. PMID: 31797357; PMCID: PMC6972511.
 17. van Rossem CC, Bolmers MDM, Schreinemaker MHF, van Geloven AAW, Bemelman WA, van Acker GJD, et al. Prospective nationwide outcome audit of surgery for suspected acute appendicitis. *British Journal of Surgery.* 2016 Jan 1;103(1):144–51. doi: 10.1002/bjs.9964. Epub 2015 Oct 28. PMID: 26509648.

18. Tekeli MT, Ilhan E, Ureyen O, Senlikci A, Yeldan E, Ozturk M, et al. How much reliable is Alvarado scoring system in reducing negative appendectomy? Indian Journal of Surgery. 2017 Apr 1;79(2): 106–10.
DOI-10.1007/s12262-015-1433-2
19. Bouali M, El Berni Y, Moufakkir A, El Bakouri A, El Hattabi K, Bensardi F, et al. Value of Alvarado scoring system in diagnosis of acute appendicitis. Annals of Medicine and Surgery. 2022 May 1;77: 103642.
DOI: 10.1016/j.amsu.2022.103642.
PMID: 35637993; PMCID: PMC9142662.
20. Memon Z, Irfan S, Fatima K, Iqbal MS, Sami W. Acute appendicitis: Diagnostic accuracy of Alvarado scoring system. Asian J Surg. 2013 Oct;36(4):144–9.
DOI-10.1016/j.asjsur.2013.04.004
21. Shuaib A, Shuaib A, Fakhra Z, Marafi B, Alsharaf K, Behbehani A. Evaluation of modified Alvarado scoring system and RIPASA scoring system as diagnostic tools of acute appendicitis. World J Emerg Med. 2017;8(4):276.
DOI:10.5847/wjem.j.1920-8642.2017.04.005.
PMID: 29123605; PMCID: PMC5675968.
22. Devanathan S, Deshpande SG, Tote D, Shinde S. Efficacy in predicting negative appendectomy rates in operated acute appendicitis patients using the Raja Isteri Pengiran Anak Saleha Appendicitis (RIPASA) score versus modified Alvarado score. Cureus. 2023 Apr 20;15(4):e37873.
DOI: 10.7759/cureus.37873.
PMID: 37223156; PMCID: PMC10202674.
23. Singh A, Parihar US, Kumawat G, Samota R, Choudhary R. To determine validation of RIPASA score in diagnosis of suspected acute appendicitis and histopathological correlation with applicability to indian population: A single institute study. Indian Journal of Surgery. 2018 Apr 1;80(2):113–7.
DOI: 10.1007/s12262-018-1731-6. Epub 2018 Feb 2.
PMID: 29915475; PMCID: PMC5991012.
24. Sammalkorpi HE, Mentula P, Savolainen H, Leppäniemi A. The introduction of adult appendicitis score reduced negative appendectomy rate. Scandinavian Journal of Surgery. 2017 Sep 1;106(3):196–201.
DOI:10.1177/1457496916683099
25. Gupta V, Gupta P, Gill C, Gupta M. Appendicitis inflammatory response score in acute appendicitis: A study at a tertiary care center in North India. Int J Appl Basic Med Res. 2022;12(4):234.
DOI: 10.4103/ijabmr.ijabmr_287_22. Epub 2022 Dec 19
PMID: 36726654; PMCID: PMC9886148.
26. Andersson M, Kolodziej B, Andersson RE. Validation of the appendicitis inflammatory response (air) score. World J Surg. 2021 Jul 1;45(7):2081–91.
DOI: 10.1007/s00268-021-06042-2. Epub 2021 Apr 6
PMID: 33825049; PMCID: PMC8154764.
27. Sağ S, Basar D, Yurdadoğan F, Pehlivan Y, Elemen L. Comparison of appendicitis scoring systems in childhood appendicitis. Turkish Archives of Pediatrics. 2022 Sep 1;57(5):532–7.
DOI: 10.5152/TurkArchPediater.2022.22076
PMID: 36062441; PMCID: PMC9524470.
28. Yazar FM, Urfalioglu A, Bakacak M, Boran ÖF, Bülbüloğlu E. Efficacy of the evaluation of inflammatory markers for the reduction of negative appendectomy rates. Indian Journal of Surgery. 2018 Feb 1;80(1):61–7.
DOI: 10.1007/s12262-016-1558-y. Epub 2016 Oct 18
PMID: 29581687; PMCID: PMC5866802.
29. Alvarado A. Inflammatory markers in acute appendicitis: Are we still look-ing for the philosopher's stone? Journal of Surgery Review Article Alvarado A J Surg. 2018;1104.
DOI: 10.29011/2575-9760.001104
30. Sushruth S, Vijayakumar C, Srinivasan K, Raj Kumar N, Balasubramaniyan G, Verma SK et al. Role of C-reactive protein, white blood cell counts, bilirubin levels, and imaging in the diagnosis of acute appendicitis as a cause of right iliac fossa pain. Cureus. 2018 Jan 16; ;10(1):e2070.
DOI: 10.7759/cureus.2070.
PMID: 29552432; PMCID: PMC5854317.
31. Acharya A, Markar SR, Ni M, Hanna GB. Biomarkers of acute appendicitis: Systematic review and cost–benefit trade-off analysis. Surg Endosc. 2017 Mar 1;31(3):1022–31.
DOI: 10.1007/s00464-016-5109-1. Epub 2016 Aug 5
PMID: 27495334; PMCID: PMC5315733.
32. Liu L, Shao Z, Yu H, Zhang W, Wang H, Mei Z. Is the platelet to lymphocyte ratio a promising biomarker to distinguish acute appendicitis? Evidence from a systematic review with meta-analysis. PLoS ONE.

- Public Library of Science. 2020;15. DOI:<https://doi.org/10.1371/journal.pone.0233470>
33. Duyan M, Vural N. Assessment of the diagnostic value of novel biomarkers in adult patients with acute appendicitis: A cross-sectional study. *Cureus*. 2022 Dec 8;14(12):e32307. DOI: 10.7759/cureus.32307. PMID: 36632249; PMCID: PMC9828092.
34. Patel M, Thomas JJ, Sarwary H. We can reduce negative paediatric appendectomy rate: A cohort study. *Annals of Medicine and Surgery*. 2021 Nov 1;71. DOI:<https://doi.org/10.1016/j.amsu.2021.102901>
35. Lehmann B, Koeflerli U, Sauter TC, Exadaktylos A, Hautz WE. Diagnostic accuracy of a pragmatic, ultrasound-based approach to adult patients with suspected acute appendicitis in the ED. *Emergency Medicine Journal*. 2022 Mar 17;39(12):931–6. DOI-10.1136/emmermed-2019-208643
36. Fu J, Zhou X, Chen L, Lu S. Abdominal ultrasound and its diagnostic accuracy in diagnosing acute appendicitis: A meta-analysis. *Frontiers in Surgery*. *Frontiers Media S.A.* 2021;8. DOI [org/10.3389/fsurg.2021.707160](https://doi.org/10.3389/fsurg.2021.707160)
37. Löfvenberg F, Salö M. Ultrasound for appendicitis: Performance and integration with clinical parameters. *Biomed Res Int*. 2016. /5697692. Epub 2016 Dec 1. PMID: 28044133; PMCID: PMC5156797.
38. Crocker C, Akl M, Abdoell M, Kamali M, Costa AF. Ultrasound and CT in the diagnosis of appendicitis: Accuracy with consideration of indeterminate examinations according to STARD guidelines. *American Journal of Roentgenology*. 2020 Sep 1;215(3):639–44. DOI: 10.2214/AJR.19.22370. Epub 2020 May 14. PMID: 32406773.
39. Rao PM, Rhea JT, Novelline RA. Sensitivity and specificity of the individual CT signs of appendicitis: Experience with 200 helical appendiceal CT examinations. *J Comput Assist Tomogr*. 1997 Sep-Oct;21(5):686-92. DOI: 10.1097/00004728-199709000-00002. PMID: 9294553.
40. Kim K, Kim YH, Kim SY, Kim S, Lee YJ, Kim KP, Lee HS, Ahn S, Kim T, Hwang SS, Song KJ, Kang SB, Kim DW, Park SH, Lee KH. Low-dose abdominal CT for evaluating suspected appendicitis. *N Engl J Med*. 2012 Apr 26;366(17):1596-605. DOI: 10.1056/NEJMoa1110734. PMID: 22533576.
41. Mikwar Z, ALSaleh N, Hafez Mousa A, Alsuayri R. Role of preoperative computed tomography in the diagnosis of acute appendicitis and reduction of negative appendectomy rates: Retrospective cross-sectional study. *Ann Med Surg (Lond)*. 2022 Apr 15;77:103609. DOI: 10.1016/j.amsu.2022.103609. PMID: 35638008; PMCID: PMC9142381.
42. Webb EM, Nguyen A, Wang ZJ, Stengel JW, Westphalen AC, Coakley FV. The negative appendectomy rate: Who benefits from preoperative CT? *American Journal of Roentgenology*. 2011 Oct;197(4):861–6. DOI: 10.2214/AJR.10.5369. PMID: 21940573.
43. *Istan journal of radiology* PK, Jesrani A. Impact of CT-fact on negative appendectomy rates: Prospective evaluation from a tertiary care hospital [Internet]. Vol. 29. Available from:<http://www.openepi.com/>
44. Park JH. Diagnostic imaging utilization in cases of acute appendicitis: Multi-center experience. *J Korean Med Sci*. 2014;29(9):1308–16. DOI-10.3346/jkms.2014.29.9.1308

© 2023 Kumar and Soma; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/102421>