



Root Parameter Investigation in Standardized Media for Bougainvillea (*Bougainvillea comm. Ex Juss.*) Propagation

Shashi Kiran Minj^{a+++*}, P. Pooja Gupta^{b#}, Rajnish Kumar^{c++},
Madhavi Khilari^{a†} and Dayanand Sai Painkra^{d++}

^a Department of Floriculture and Landscape Architecture, IGKV, Raipur, Chhattisgarh, India.

^b Department of Floriculture and Landscape Architecture, College of Agriculture, IGKV, Raipur, Chhattisgarh, India.

^c Department of Fruit Science, IGKV, Raipur, Chhattisgarh, India.

^d Department of Forestry, College of Agriculture, IGKV, Raipur, Chhattisgarh, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i183448

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/103976>

Original Research Article

Received: 22/05/2023

Accepted: 25/07/2023

Published: 26/07/2023

ABSTRACT

The study was conducted in Completely Randomized Design (CRD) with 16 treatments and 5 replications comprising of soilless media and biofertilizer Azotobacter. The present experiment entitled "Standardization of media for propagation of Bougainvillea (*Bougainvillea comm. Ex Juss.*)" was carried out at Horticultural nursery, College of Agriculture, IGKV, Raipur, (C.G) during the year

⁺⁺ Ph.D.Scholar;

[#] Assistant Professor;

[†] M.sc Scholar;

*Corresponding author: E-mail: shashiminj1000@gmail.com;

2019-20. All of the media treatments compared to the control had better vegetative and rooting qualities. Soil + Sand + Cocopeat had the highest sprouting rate and survival percentage of all the media treatments tested. All other Root attributes viz. The minimum days to rooting (144.80), The longest root length (6.08), highest number of roots per cutting (63.54). The most cuttings that could successfully take root (24.00) were found to be idealistic in media including soil, sand, and cocopeat. The results showed that all increasing media treatments had a substantial impact on all variables related to Bougainvillea development, including shoot and root growth. The results showed that the development of Bougainvillea as well as all the growing media treatments had a significant impact on all the characters' associated root growth. The maximum overall production cost has been seen in T6 (Soil + Sand + Cocopeat) and the least production cost in T13 (Soil + Azotobacter). Combination of rooting media has been beneficial to significantly increase the vegetative growth parameters and economics as compared to control in Bougainvillea.

Keywords: *Azotobacter*; *cocopeat*; *Bougainvillea*.

1. INTRODUCTION

Bougainvillea is a genus of thorny ornamental vines, bushes, and trees belonging to the four o'clock family, Nyctaginaceae. This evergreen shrub has a variety of flower colours like Red, Orange, Pink, Purple, White etc. It is mainly grown in open field condition. They flower most heavily in winter and early spring, but some plants put forth scattered clusters all year. Depending on the variety, it can be grown as a bush, hedge, in pots, as a bonsai, as standard or semi standard shrub or can be trained in pergola or arches.

Likewise, other ornamental plants, the multiplication of Bougainvillea is done by using vegetative part *i.e.* Stem cutting, Layering, Budding. Stem cutting is regarded as the cheapest and easiest method of Bougainvillea; however the success rate of propagation is very limited. Rooting percentage or else we can say rooting responses varies from variety to variety. As we know, in most of ornamental plants, the success of rooting in cuttings is mainly depends upon variety, time of planting and use of growth regulators [1]. Rainy season is the best time for planting Bougainvillea.

Use of different types of rooting media having different characteristics is important for quality rooting in cuttings. Though there is no universal rooting mix, different medias are used according to cutting types, different seasons etc [2,3]. Highest rooting percentage (> 97%) was seen with basal stem cutting compared with terminal and middle cutting. Highest number of roots/cuttings (27.66) was seen with basal stem cuttings after 60 days of planting the cuttings. Highest plant height(67.66 cm) and number of

leaves/cutting (46.00) was seen with soil+sand (1:1) after 90 days of planting the cuttings [4].

We are aware of the fact that, success rate of bougainvillea is very limited. Hence, to improve rooting in cuttings, several rooting hormones are widely used during propagation. The variety 'Mahara' is easily rooted with the help of auxin treatment. Growth hormones which are used are indole -3-butyric acid (IBA), 1-naphthalene acetic acid (NAA), indole -3-acetic acid (IAA). These hormones with different concentrations enhance the growth and development of roots.

2. MATERIALS AND METHODS

The field experiment was performed during the year 2019-2020 at the Horticulture research cum Instructional Farm, College of Agriculture, IGKV, Raipur (C.G).

After cutting from the mother plants, the hardwood cuttings (10-15 cm) of pencil thickness were prepared. The leaves were carefully trimmed from the shoots to prevent damage to the axillary buds. The cuttings were prepared to differentiate both ends during planting by giving a slanting cut on the upper end and a straight cut on the lower end. The cuttings were planted immediately in media mixtures after quick dip treatment of IBA@ 4000 ppm for (<5 sec.) directly in the polythene bags.

1g of IBA was first dissolved in a few drops of 0.1N NaOH for 4000 ppm of IBA solution and then up to 250 ml of distilled water was make up in order to achieve the right concentration of IBA.

In the polybags with pre-punched bottom holes for drainage, various treatments containing

different media were filled in. Various intercultural operations were carried out when needed. Various shoot and root attributes were noted and Statistically evaluated using Anova for Completely Randomized Design (CRD) [5].

Table 1. Treatment details [5]

S.no	Treatments	Treatment details
1.	T ₁	Sand
2.	T ₂	Cocopeat
3.	T ₃	Rice husk
4.	T ₄	Sawdust
5.	T ₅	Soil + FYM + Sawdust
6.	T ₆	Soil + Sand + Cocopeat
7.	T ₇	Sand + FYM
8.	T ₈	Sand + Azotobacter
9.	T ₉	Soil + Cocopeat
10.	T ₁₀	Soil + Sand + Ricehusk
11.	T ₁₁	Soil + Sand
12.	T ₁₂	Cocopeat + Azotobacter
13.	T ₁₃	Soil + Azotobacter
14.	T ₁₄	Ricehusk + Azotobacter
15.	T ₁₅	Sawdust + Azotobacter
16.	T ₁₆	Soil (control)

3. RESULTS AND DISCUSSION

The minimum days to rooting (144.80) was recorded with T₆ (Soil + Sand + Cocopeat). Likewise, the maximum days to rooting (152.40) was recorded with control T₁₆ (Soil). this can be because of cocopeat, which has low nitrogen, calcium and magnesium but it is high in phosphorus and potassium [6] in ornamental pot plants. Benefits of phosphorus in plants is that, it is good for the root growth and development.

The longest root length (6.08) was observed in T₆ (Soil + Sand + Cocopeat). The least root length (3.60) was observed in T₁₅ (Sawdust + *Azotobacter*) at 180 DAP.

The increase in the length of the cutting roots treated with growth regulators may be due to the accumulation of metabolism at the site of auxin application, cell enlargement, increased carbohydrate hydrolysis, new protein synthesis and auxin-induced cell division [7]. The standard concentration of IBA combined with media increases the duration of the roots of *bougainvillea*, an effect recorded by [4].

Data recorded at 180 DAP, treatment T₆ (Soil + Sand + Cocopeat) showed highest number of roots per cutting (63.54). The lowest number of roots (39.78) was recorded with control T₁₆ (Soil). These findings are in agreement with the reports of Nazari *et al.* [8] in hyacinth.

The maximum number of cutting attained rooting (24.00) was recorded in T₆ (Soil + Sand + Cocopeat). The minimum number of cutting attained rooting (17.00) was recorded in Control T₁₆ (Soil). Highest survival percentage was also found in T₆. This may be attributed to metabolite aggregation at the site of auxin application [9], accelerated carbohydrate hydrolysis, new protein synthesis, auxin-induced cell elongation and cell division [10]. Increases in the rooting percentage may be attributable to proper IBA concentration, high carbohydrates in turn, and low nitrogen levels contribute to further root forming [11].




Fig. 1. Graphical representation of effect of growing media on *Bougainvillea* (Root parameters)

Table 2. Effect of growing media on *Bougainvillea*

Tr. No.	Treatments	Days to rooting	Length of root	Number of roots
			180 DAP	180 DAP
T1	Sand	145.80	4.14	42.24
T2	Cocopeat	149.80	4.18	51.04
T3	Ricehusk	145.40	4.42	57.73
T4	Sawdust	149.00	4.32	51.92
T5	Soil + FYM + Sawdust	147.60	5.24	63.36
T6	Soil + Sand + Cocopeat	144.80	6.08	63.54
T7	Sand + FYM	146.00	4.24	58.08
T8	Sand + <i>Azotobacter</i>	145.80	4.58	49.98
T9	Soil + Cocopeat	148.60	4.58	51.22
T10	Soil + Sand + Ricehusk	145.00	4.02	52.27
T11	Soil + Sand	146.00	4.26	52.10
T12	Cocopeat + <i>Azotobacter</i>	147.40	4.30	49.98
T13	Soil + <i>Azotobacter</i>	149.80	4.42	52.80
T14	Ricehusk + <i>Azotobacter</i>	146.00	2.40	55.62
T15	Sawdust + <i>Azotobacter</i>	147.00	3.60	46.29
T16	Soil (control)	152.40	4.54	39.78
	SEm ±	1.55	0.30	3.48
	C.D @ 5%	4.38	0.87	9.82

Table 3. Effect of growing media on *Bougainvillea*

Tr. No.	Treatments	Number of cutting attained rooting	Survival %
T1	Sand	19	64
T2	Cocopeat	20	68
T3	Ricehusk	19	64
T4	Sawdust	20	64
T5	Soil + FYM + Sawdust	22	80
T6	Soil + Sand + Cocopeat	24	88
T7	Sand + FYM	23	84
T8	Sand + <i>Azotobacter</i>	22	84
T9	Soil + Cocopeat	19	68
T10	Soil + Sand + Ricehusk	23	84
T11	Soil + Sand	19	68
T12	Cocopeat + <i>Azotobacter</i>	22	80
T13	Soil + <i>Azotobacter</i>	19	68
T14	Ricehusk + <i>Azotobacter</i>	23	80
T15	Sawdust + <i>Azotobacter</i>	22	80
T16	Soil (control)	17	56
	SEm ±	0.28	
	C.D @ 5%	0.78	

4. CONCLUSION

As cocopeat contains 20 percent to 30 percent fibre of varying length, it is considered as a good growing media. Cocopeat is high in phosphorus and potassium [6]. Benefits of phosphorus in plants is that, it is good for the root growth and development. Whereas, potassium helps in strong build up of stems. It holds upto 8-9 times its weight in water, hence it has high water holding capacity which may cause poor aeration within the media, thus affecting the oxygen diffusion to the roots. The solution to this problem is to incorporate coarser material into cocopeat

could improve the aeration status of the media [12].

FUTURE SCOPE

As there is scarcity of land, hence the supply of quality topsoil is declining rapidly. Thus, Promoting the use of soil-free media in ornamental crop production.

ACKNOWLEDGEMENT

The authors are thankful to Horticultural nursery, College of Agriculture, Indira Gandhi Agriculture

College, Raipur (C.G) for providing planting material to conduct the experiment.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Rowezak MMA. Response of some Ornamental plants to Treatment with Growth substances. Sc. Thesis, Fac. Agric., Cairo Univ; 2001.
2. Okunlola AI. The effects of cutting types and length on rooting of *Duranta repens* in the Nursery. Global Journal of Human Social Science, Geography, Geosciences, Environmental, and Disaster Management. 2013; 13(3): 1.
3. Gupta VN, Banerj BK, Datta SK. Effect of auxin on rooting and sprouting behavior of stem cuttings of Bougainvillea under mist. Haryana Journal of Horticulture Sciences. 2002; 31: 42.
4. Eed AM, Albana^a B, Almaqtari S. The effect of growing media and stem cutting type on rooting and growth of Bougainvillea spectabilis plants. Univ. Aden J. Nat. and Appl. Sc. 2015;19(1): 141-147.
5. Minj SK, Gupta P, Sahu RK, Khilari M. Standardization of media for propagation of Bougainvillea (*Bougainvillea comm. Ex Juss.*). The Pharma Innovation Journal. 2023; 12(5): 927-930.
6. Gohil P, Gohil M, Rajatiya J, Halepotara F, Solanki M, Malam VR, Barad R. Role of growing media for ornamental pot plants. Int. J. Pure App. Biosci. 2018;6(1):1219-1224.
7. Singh KK, Rawat JMS, Tomor YK. Influence of indole butyric acid (IBA) on rooting potential of Torch Glory Bougainvillea glabra during winter seasons. Journal of Horticulture Science and Ornamental Plants. 2011;3(2):162.
8. Nazari F, Farahmand H, Khos-Khui M, Salehi H. Effects of coir as a component of potting media on growth, flowering and physiological characteristics of hyacinth. Int. J. of Agric. and Food Sci., 2011;1(2): 34-38.
9. Minj SK, Gupta P, Sahu RK, Khilari M. Standardization of media for propagation of Bougainvillea (*Bougainvillea comm. Ex Juss.*). The Pharma Innovation Journal. 2023; 12(5): 927.
10. Fadwa MAA, Yahia HAE. Vegetative propagation of *Peltophorium petrocarpum* (DC) Backer ex k.Heyne: A multi-purpose tree. Net Journal of Agricultural Science. 2014; 2(4): 113.
11. Carlson MC. Microchemical studies of rooting and non-rooting rose cuttings. Bot. Gaz., 1929;87(1): 64.
12. Awang Y, Shaharom AS, Mohamad RB, Selamat A. Chemical and physical characteristics of cocopeat-based media mixtures and their effects on the growth and development of Celosia cristata. AM. J. Agri. & Biol. Sci. 2009;4(1): 63-71.

© 2023 Minj et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/103976>