



The Pattern of Hematological Abnormalities in NS1 Positive Dengue Fever

Habibur Rahman ^{a+++*}, Tahrina Akter ^{b#}, Sabrina Islam ^{c++},
A. B. M. Kamrul Hasan ^{d++}, Sabiya Afroz ^{a†},
Mahmood Rahman ^{e‡} and Shaleha Nazneen ^{f++}

^a Department of Transfusion Medicine, Shaheed Suhrawardy Medical College Hospital, Dhaka, Bangladesh.

^b Sheikh Hasina National Institute of Burn & Plastic Surgery, Dhaka, Bangladesh.

^c Department of Transfusion Medicine, Shaheed Tajuddin Ahmed Medical College Hospital, Gazipur, Bangladesh.

^d Department of Transfusion Medicine, M. Abur Rahim Medical College Hospital, Dinajpur, Bangladesh.

^e National Institute of ENT, Dhaka, Bangladesh.

^f Department of Hematology, Shaheed Suhrawardy Medical College Hospital, Dhaka, Bangladesh.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/108316>

Original Research Article

Received: 14/10/2023
Accepted: 01/11/2023
Published: 06/11/2023

⁺⁺Assistant Professor & Head;

[#]Medical Officer;

[†]Blood Transfusion Officer;

[‡]Junior Consultant (Anesthesia);

*Corresponding author: E-mail: habib.nidch@gmail.com;

ABSTRACT

Background: Dengue fever, caused by the dengue virus and transmitted by mosquitoes, is a global health threat, especially in tropical regions. The NS1 antigen is a key diagnostic marker for dengue. Hematological changes, such as platelet counts, white blood cells, and hematocrit levels, play a crucial role in diagnosing and managing NS1-positive dengue fever. Studying these abnormalities enhances our understanding of the disease and improves patient care.

Aim of the Study: The aim of the study was to assess the pattern of hematological abnormalities in NS1-positive dengue fever.

Methods: This prospective observational study was conducted at the Department of Transfusion Medicine, Suhrawardy Medical College Hospital, Dhaka, Bangladesh from February 2023 to September 2023. A total of 196 NS1 positive cases enrolled in this study using purposive sampling methods. Demographic and clinical data were meticulously recorded and analyzed with MS Office tools for dissemination.

Results: The majority of participants (51.53%) fell within the 16-28 age group, with 61% being male. Prominent symptoms included fever (82.1%), headache (62.8%), nausea (48.5%), and anorexia (42.9%). Dengue fever was the most frequent diagnosis (88.27%), followed by Dengue hemorrhagic fever (9.69%). Among those with leucopenia (n=68), lymphocytosis prevailed (41.18%), followed by neutrophilia (20.59%). In cases with platelet counts $\leq 20,000$, petechiae were observed in 37.9%, hematemesis in 34.4%, and melena in 27.5%.

Conclusion: In assessing the NS1 positive dengue fever patients, among cases of leucopenia, lymphocytosis is currently more prevalent than neutrophilia. Among individuals with platelet counts $\leq 20,000$, common bleeding manifestations are currently being observed. Among cases of leukocytosis, neutrophilia is predominantly found in half of such patients.

Keywords: NS1 positive; dengue hemorrhagic fever; haematological profile; shock syndrome.

1. INTRODUCTION

Dengue fever and dengue hemorrhagic fever have emerged as significant global public health concerns in recent decades. Dengue fever is typically caused by an arthropod-borne virus and manifests as an acute febrile illness characterized by sudden-onset fever, headache, body aches, generalized weakness, retro-orbital pain, and rash [1,2]. Diagnosis of dengue fever relies on a combination of clinical evaluation and laboratory data [3]. Clinicians consider clinical symptoms and signs, along with both non-specific parameters like blood count, platelet count, prothrombin time (PT), and liver function tests, as well as specific tests such as viral isolation and serology for antibody examination to make an accurate diagnosis [4]. Dengue infection encompasses a spectrum of manifestations, ranging from undifferentiated fever and dengue fever (DF) to more severe conditions like dengue hemorrhagic fever (DHF) with shock and expanded dengue syndrome [5]. Every year, approximately 50 million dengue infections occur, leading to approximately 500,000 hospitalizations, with a predominant number of these patients being children [6]. Dengue presents a wide clinical spectrum, ranging from asymptomatic fever to life-

threatening conditions like dengue hemorrhagic fever (DHF) and dengue shock syndrome (DSS). Since there is no licensed vaccine available, the primary approach to management involves fluid management and monitoring for complications [7]. One of the most notable hematological changes in dengue cases is thrombocytopenia and leukopenia, often accompanied by lymphocytosis characterized by reactive lymphocytes. Additionally, there is a progressive increase in hematocrit concentration during the progression of dengue fever, often reaching up to a 20% increase from the patient's baseline [8,9]. As part of its global strategy from 2012 to 2020, the World Health Organization (WHO) has set the first objective of reducing dengue mortality by 50% by the year 2020 [10]. In addition to dengue-specific parameters, platelet count serves as a prognostic laboratory parameter that can help identify the severity of dengue. While thrombocytopenia may not be an early indicator of severe dengue, it is useful in predicting the progression of the disease [11]. Raising awareness about dengue fever is crucial in its prevention. Unfortunately, the level of awareness among the population remains inadequate. Some studies have shown that only about one-third of the adult population possesses sufficient knowledge about dengue fever and its rapid

spread to others. Furthermore, only approximately 17% of them are aware of and knowledgeable about preventive measures [12]. The object of this study was to assess the pattern of hematological abnormalities in NS1-positive dengue fever.

2. METHODOLOGY

This prospective observational study was conducted at the Department of Transfusion Medicine, Suhrawardy Medical College Hospital, Dhaka, Bangladesh from February 2023 to September 2023. A total of 196 NS1 positive cases were included in this study as the study population. The sample selection process utilized a convenient purposive sampling technique. All cases that tested serologically positive for Dengue NS1 antigen, IgM antibody, or IgG antibody were eligible for inclusion in the study. A comprehensive patient history was documented, which encompassed age, gender, complaints, clinical presentation, and laboratory investigations for all cases. Before data collection, written consent was obtained from all participants following ethical guidelines. Exclusion criteria for this study involved cases

where non-cooperation was anticipated from the participants and individuals who tested positive for dengue but also exhibited co-existing infections such as malaria or typhoid. Demographic and clinical information of all participants was meticulously recorded, and data analysis and dissemination were performed using MS Office tools.

3. RESULTS

The majority of our participants (51.53%) were from the 16-28 years' age group. More than half of the participants (61%) were male and the rest of 39% were female. Among the clinical symptoms reported by patients, the most significant findings were fever (82.1%), headache (62.8%), nausea (48.5%), and anorexia (42.9%). These symptoms were the most prevalent and are likely to be critical indicators for diagnosis and management. Among our total participants, dengue fever was the most common diagnosis, accounting for 88.27% of cases (173 individuals). A smaller proportion of cases were diagnosed with Dengue hemorrhagic fever, representing 9.69% of cases (19 individuals). The least common diagnosis was Dengue shock

Table 1. Age distribution of participants (N=196)

Age (Years)	n	%
16-28 yrs.	101	51.53%
29-38 yrs.	34	17.35%
39-48 yrs.	29	14.80%
49-60 yrs.	18	9.18%
>61 yrs.	14	7.14%




Fig. 1. Column chart showed age group wise patients distribution. (N=196)




Fig. 2. Pie chart showed gender wise patients distribution (N=196)

Table 2. Clinical symptoms distribution (N=196)

Signs/Symptoms	n	%
Fever	161	82.1%
Headache	123	62.8%
Nausea	95	48.5%
Anorexia	84	42.9%
Vomiting	76	38.8%
Myalgia	69	35.2%
Abdominal pain	54	27.6%
Retro orbital pain	28	14.3%
Arthralgia	21	10.7%
Back pain	17	8.7%




Fig. 3. Bar chart showed clinical symptoms wise patients (N=196)

syndrome, observed in 2.04% of cases (4 individuals). Among those with leucopenia (n=68), lymphocytosis was the most common

pattern, observed in 41.18% of cases, followed by neutrophilia at 20.59%. In individuals with normal WBC counts (n=120), lymphocytosis

remained prevalent at 47.5%, while neutrophilia was observed at 17.5%. Among cases of leukocytosis (n=8), neutrophilia was predominant in 50% of individuals, with notable lymphocytosis in 25%. Among individuals with a platelet count of $\leq 20,000$, 37.9% (19 individuals) experienced petechiae, 34.4% (18 individuals) had

hematemesis, and 27.5% (15 individuals) had melena. No cases of these bleeding manifestations were observed in individuals with platelet counts higher than 50,000. The total number of cases considered in this categorization was 57.

Table 3. Clinical spectrum of dengue (N=196)

Diagnosis	n	%
Dengue fever	173	88.27%
Dengue hemorrhagic fever	19	9.69%
Dengue shock syndrome	4	2.04%




Fig. 4. Column chart showed Clinical spectrum of dengue patients (N=196)




Fig. 5. Line chart showed differential WBC count pattern of the patients (N=196)

Table 4. Differential WBC count pattern (N=196)

Pattern	Leucopenia		Normal WBC		Leukocytosis	
	(n=68)		(n=120)		(n=8)	
	n	%	n	%	n	%
Lymphocytosis	28	41.18%	57	47.5%	2	25.0%
Neutrophilia	14	20.58%	21	17.5%	4	50.0%
Within normal limit	26	38.24%	42	35.0%	2	25.0%
Total	68	34.7%	120	61.2%	8	4.1%

Table 5. Thrombocytopenia based on bleeding manifestations (n=57)

Manifestation	≤20,000	21-30,000	31-40,000	41-50,000	51-1 lac	>1 lac	Total
Petechiae	9	8	7	3	0	0	24
Hematemesis	6	6	6	2	0	0	18
Melena	4	4	2	0	0	0	10
Total	19(37.9)	18(34.4)	15(27.5)	5	0	0	57

4. DISCUSSION

This study aimed to assess the pattern of hematological abnormalities in NS1-positive dengue fever. The majority of our participants (51.53%) belonged to the 16-28 years' age group. In terms of gender distribution, more than half (61%) were male, while the remaining 39% were female. A similar finding was also observed in studies conducted by Shamsunder Khatroth [13] and Yashaswini LS et al. [14]. In this study, the clinical symptoms reported by patients revealed several significant findings. The most common symptoms included fever (82.1%), headache (62.8%), nausea (48.5%), and anorexia (42.9%). These prevalent symptoms are likely to be crucial indicators for diagnosis and management. Similar findings were also observed in studies conducted by other authors [15,16]. Among our total participants, dengue fever was the most common diagnosis, accounting for 88.27% of cases (173 individuals). Dengue hemorrhagic fever was diagnosed in a smaller proportion of cases, representing 9.69% (19 individuals). The least common diagnosis was Dengue shock syndrome, observed in only 2.04% of cases (4 individuals). Similar findings were noted in another study as well [17]. In our study, among the individuals with leucopenia (n=68), 41.18% exhibited lymphocytosis as the most common pattern, followed by 20.59% with neutrophilia. In those with normal white blood cell (WBC) counts (n=120), 47.5% had lymphocytosis, while 17.5% had neutrophilia. Among cases of leukocytosis (n=8), 50% showed neutrophilia as the predominant pattern and 25% had notable lymphocytosis. Ferede et al [16], Priya, and Bindu M [17] showed that maximum

cases in leucocyte count range within normal limits. Moreover, Joshi AA et al found that maximum patients in the leucopenia range. [18] In our study, among individuals with a platelet count of ≤20,000 (total cases considered: 57), 37.9% (19 individuals) exhibited petechiae, 34.4% (18 individuals) presented with hematemesis, and 27.5% (15 individuals) showed melena as bleeding manifestations. No instances of these bleeding manifestations were observed in individuals with platelet counts higher than 50,000. Here the findings of Tewari et al. [19] were comparable. All the findings of this current study may provide valuable insights for future research in similar studies.

5. CONCLUSION AND RECOMMENDATION

Drawing insights from this study, it becomes evident that in patients with confirmed NS1-positive dengue fever, several prominent symptoms tend to manifest frequently. These symptoms notably include fever, headache, nausea, and anorexia. Moreover, the most prevalent diagnoses encountered in these cases are dengue fever itself and its more severe form, dengue hemorrhagic fever. When delving into a more detailed examination of NS1-positive dengue fever patients, it becomes apparent that among those exhibiting leucopenia, lymphocytosis tends to prevail over neutrophilia. Furthermore, a noteworthy observation is the presence of common bleeding manifestations among individuals with platelet counts falling at or below 20,000. Among cases of leukocytosis, neutrophilia is found predominant among half of such patients.

6. LIMITATION OF THE STUDY

This single-center study featured a limited sample size and a short data collection period, potentially restricting its representativeness for the wider national population. While it offers valuable insights within its context, caution is needed when extrapolating to a broader population. Future research with larger, more diverse samples, and longer data collection can enhance our national-level understanding of the topic.

ETHICAL APPROVAL

As per international standard or university standards written ethical approval has been collected and preserved by the author(s).

CONSENT

As per international standard or university standards, patient(s) written consent has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Guha-Sapir D, Schimmer B. Dengue fever: new paradigms for a changing epidemiology. *Emerg Themes Epidemiol*. 2005;2(1):1-10.
2. Park K. Epidemiology of Communicable Diseases: Dengue syndrome. In: Park's Textbook of Preventive and Social Medicine. 20th ed. Jabalpur, India: M/s Bhanarsidas Bhanot. 2009;218-22.
3. De Paula SO, Fonseca BA. Dengue: A review of the laboratory tests a clinician must know to achieve a correct diagnosis. *Braz J Infect Dis*. 2004;8(6):390-8.
4. Srichaikul T, Nimmannita S. Haematology in dengue and dengue haemorrhagic. *Baillieres Best Pract Res Clin Haematol*. 2000;13(2):261-76.
5. Ministry of Health Sri Lanka. Guidelines on Management of Dengue Fever & Dengue Haemorrhagic Fever in Children and Adolescents. 2012;1-2.
6. Guzman MG, Halstead SB, Artsob H, Buchy P, Farrar J, Gubler DJ, et al. Dengue: A continuing global threat. *Nature Reviews. Microbiology*. 2010;2460:7-14
7. Paranavitane S, Gomes L, Kamaladasa A, Adikari T, Wickramasinghe N, Jeewandara C Et al. Dengue NS1 antigen as a marker of severe clinical disease. *BMC Infect Dis*. 2014;14 (1).
8. Kao CL, King CC, Chao DY, Wu HL, Chang GJ. Laboratory diagnosis of dengue virus infection: current and future perspectives in clinical diagnosis and public health. *J Microbiol Immunol Infect*. 2005;38(1):5-16.
9. Ageep AK, Malik AA, Elkarsani MS. Clinical presentations and laboratory findings in suspected cases of dengue virus. *Saudi Med J*. 2006;27(11):1711-3.
10. Nujum ZT, Vijayakumar K, Kavithai P. World Health Organization. Global strategy for dengue prevention and control 2012-2020. Geneva: WHO, 2012. *Dengue Bulletin*. 2013;37:11-30.
11. Brasier A, Scott T, Morrison A, Kochel T, Spratt H, Bazan I et al. A Three-Component Biomarker Panel for Prediction of Dengue Hemorrhagic Fever. *Am J Trop Med Hyg*. 2012;86(2):341-348.
12. Malhotra G, Yadav A, Dudeja P. Knowledge, awareness and practices of dengue in a city. *International Journal of Medical Science and Public Health*. 2014;3(3):295-299.
13. Shamsunder Khatroth. A Study on Clinical and Hematological Profile of Dengue Fever in a Tertiary Care Hospital. *IAIM*. 2017;4(8): 96-102.
14. Yashaswini LS, Priya. Vidyashri. A study of hematological parameters and requirement of platelet transfusion in dengue fever. *Int J Adv Med*. 2017;4:1668-71.
15. Tahlan A, Bhattacharya A, Singla N, Singh R. Haematological profile of dengue fever. *Int J Res Med Sci*. 2017;5:5367-71.
16. Ferede G, Tiruneh M, Abate E, et al. A study of clinical, hematological, and biochemical profiles of patients with dengue viral infections in Northwest Ethiopia: Implications for patient management. *BMC Infect Dis*. 2018;616:1-6.
17. Priya AV, Bindu M H. Clinico – Hematological profile of dengue in children in a tertiary care hospital. *IP Int J Med Paediatr Oncol*. 2020;6(4):147-151.
18. Joshi AA, Gayathri BR, Gowda YR. The total leucocyte count-its utility in dengue. *Int J Adv*. 2017;4:1621-6.

19. Tewari K, Tewari VV, Mehta R. Clinical and hematological profile of patients with dengue fever at a tertiary care hospital – an observational study. *Mediterr J Hematol Infect Dis.* 2018;10(1): 1-7.

© 2023 Rahman et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/108316>