



# Generation Mean Analysis for Estimating Gene Action, Heterosis and Inbreeding Depression in Yield and Foliar Disease Resistance in Three Faba Bean (*Vicia faba* L.) Crosses

Gehan G. Abo-Zaid <sup>a</sup>, Salwa M. Mostafa <sup>a\*</sup>,  
Shymaa F.A. Kalboush <sup>a</sup> and R.A. El-Refaey <sup>b</sup>

<sup>a</sup> Food Legumes Research Section, Field Crop Research Institute, Agricultural Research Center, Egypt.

<sup>b</sup> Agronomy Department, Faculty of Agriculture, Tanta University, Egypt.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ajrcs/2024/v9i2280>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/119770>

**Original Research Article**

**Received: 21/05/2024**

**Accepted: 23/07/2024**

**Published: 30/07/2024**

## ABSTRACT

This study was conducted at the Sakha Agricultural Research Station farm over three growing seasons (2019/20, 2020/21, and 2021/22) to introduce new faba bean genotypes possessing high yield potential and increased resistance to foliar diseases. The research involved six populations (P1, P2, F1, F2, BC1, and BC2) from three crosses (Cross-1: Santamora x Foal Sbai Labiade,

\*Corresponding author: E-mail: [salwamostfa82@gmail.com](mailto:salwamostfa82@gmail.com);

**Cite as:** Abo-Zaid, Gehan G., Salwa M. Mostafa, Shymaa F.A. Kalboush, and R.A. El-Refaey. 2024. "Generation Mean Analysis for Estimating Gene Action, Heterosis and Inbreeding Depression in Yield and Foliar Disease Resistance in Three Faba Bean (*Vicia faba* L.) Crosses". *Asian Journal of Research in Crop Science* 9 (2):192-203. <https://doi.org/10.9734/ajrcs/2024/v9i2280>.

Cross-2: Santamora x Misr 3, and Cross-3: Cairo 4 x Misr 3), analyzed using a randomized complete block design with three replications. Generation mean analysis revealed significant mid and better parent heterosis for most traits, indicating over-dominance. Inbreeding depression manifested negatively for chocolate spot and rust diseases but positively for seed count and seed yield per plant across all crosses. Non-allelic interactions were present for all traits, and dominance gene effects were more influential than additive gene effects for most traits. Mean performance analysis showed that F1 populations had lower disease reactions, but higher yields compared to the better parent, while F2 populations displayed greater genetic variability due to segregation. Heterosis was significant for yield and its components across all crosses but less beneficial for disease resistance. Inbreeding depression was significant and negative for disease reactions but positive for yield traits. The study found high broad-sense heritability for all traits, suggesting potential for phenotypic selection. Narrow-sense heritability values were moderate for qualitative traits and lower for quantitative traits. Genetic advance under selection indicated the highest gains were associated with moderate heritability values. Significant non-allelic interactions suggested the need for specific breeding strategies, with dominance gene effects more pronounced than additive effects. These findings support the feasibility of selecting high-yielding, disease-resistant faba bean genotypes through targeted breeding methods, aligning with previous research on the genetic inheritance of these traits.

**Keywords:** *Faba bean; foliar disease resistance; generation mean analysis; genetic resources.*

## 1. INTRODUCTION

Faba bean is a crucial legume crop in Egypt, valued for its high nutritional content as a protein source and its role in enhancing soil fertility through nitrogen fixation. It is commonly used in crop rotation systems. The crop is partially self-pollinated and exhibits both heterosis and inbreeding depression [1]. During the 2020/21 season, Egypt faba bean cultivation covered approximately 120,000 feddan, producing around 281,000 tons with an average yield of 10 ardab per feddan (1 ardab = 155 kg), meeting about 41% of the country's consumption needs [2].

Understanding gene action is vital for breeders to plan effective breeding programs. Using generation mean analysis on multiple populations can provide essential genetic insights. Foliar diseases like chocolate spot and rust significantly impact faba bean crops globally, with susceptible varieties experiencing losses exceeding 55% [3]. To address these challenges, breeders use various biometrical techniques to enhance breeding procedures, tackling both biotic and abiotic stresses. These techniques help assess the genetic effects on quantitative traits, thereby optimizing crop yield potential.

Hybrid vigor for seed yield is linked to heterotic effects on yield and its components. In faba bean, these effects are particularly notable in the

F1 generation, especially among widely divergent parents [4,5]. Inbreeding depression reduces autofertility and yield in the absence of pollinators and diminishes yield due to the loss of hybrid vigor. Poulsen (1979) reported that inbreeding depression can reduce yield by approximately 11%.

This study aimed to explore the genetic factors influencing yield and disease resistance in three faba bean crosses. The goal was to identify traits with higher heritability for targeted breeding efforts, focusing on gene action, heterosis, inbreeding depression, and the expected and predicted genetic advances for resistance to foliar diseases, yield, and its components.

## 2. MATERIALS AND METHODS

This study, conducted over three seasons (2019/20, 2020/21, and 2021/22) at the Sakha Agricultural Research Station, Agricultural Research Center, Egypt aimed to identify new faba bean genotypes with high yield and resistance to chocolate spot and rust diseases. The research focused on estimating heterosis, inbreeding depression, and the inheritance of yield and its components using generation mean analysis for three faba bean crosses. Table 1 provides details on the parental genotypes, including their name, origin, botanical group, and special remarks.

**Table 1. Name, origin, botanical group, and special remarks of the parental genotypes**

Genotypes	Original	Botanical group	Special remarks
Santamora	Spain	Major	Medium flowering and maturity, high-yielding ability and resistance to foliar diseases
Foul Sbai Labiade	Morocco	Major	Medium flowering and maturity, high-yielding ability and resistance to foliar diseases
Misr 3 and Cairo 4	Egypt	Equina	This varieties exhibits early flowering and maturity, along with tolerance to Orobanche and moderate resistance to foliar diseases.

Reactions to foliar diseases were recorded in mid-February for chocolate spot and mid-March for rust, using a grading system from 1 to 9, where 1-4 indicates resistance, 5-6 moderate resistance, and 7-9 high susceptibility, based on the disease scales suggested by Bernier et al [6]. In the 2019/20 season, four parental genotypes were crossed in wire cages to produce three crosses: Cross-1 (Santamora x Foul Sbai Labiade), Cross-2 (Santamora x Misr 3), and Cross-3 (Cairo 4 x Misr 3). In the 2020/21 season, hybrid seeds from these crosses were sown to produce F<sub>1</sub> plants, which were then selfed to generate the F<sub>2</sub> generation. Each F<sub>1</sub> was also backcrossed to its parents to produce BC<sub>1</sub> and BC<sub>2</sub> seeds, and additional F<sub>1</sub> seeds were obtained by crossing the parents again. In the 2021/22 season, six populations (P<sub>1</sub>, P<sub>2</sub>, F<sub>1</sub>, F<sub>2</sub>, BC<sub>1</sub>, and BC<sub>2</sub>) from the three crosses were planted in a randomized complete block design with three replications. Plot sizes varied: four ridges for P<sub>1</sub>, P<sub>2</sub>, and F<sub>1</sub>; nine ridges for BC<sub>1</sub> and BC<sub>2</sub>; and 20 ridges for F<sub>2</sub>. Each ridge was three meters long and 60 cm wide, with hills spaced 20 cm apart and one seed per hill. The studied traits included flowering date, chocolate spot disease reaction, rust disease reaction, plant height, number of branches per plant, number of pods per plant, number of seeds per plant, 100-seed weight (g), and seed yield per plant (g). Data were collected from 30 plants for P<sub>1</sub>, P<sub>2</sub>, and F<sub>1</sub>; 250 plants for F<sub>2</sub>; and 120 plants for each of BC<sub>1</sub> and BC<sub>2</sub> in each cross.

### 2.1 Statistical and Genetic Analysis

The mean ( $\bar{x}$ ), variances (S<sup>2</sup>), and variance of the mean (S<sup>2</sup>m) were calculated for each population (P<sub>1</sub>, P<sub>2</sub>, F<sub>1</sub>, BC<sub>1</sub>, BC<sub>2</sub>, and F<sub>2</sub>) to assess the presence of non-allelic interactions. A scaling test, as outlined by Mather [7] and Hayman [8], was employed. Generation mean analysis was conducted following Gamble [9]. Heritability estimates were computed in both broad (H) and narrow (h<sup>2</sup>) senses according to Allard [10] and Warner [11], respectively. The

expected genetic advance from selection (Ga) was calculated using the formula proposed by Johnson et al [12], and the predicted genetic advance from selection was expressed as a percentage of the F<sub>2</sub> mean (Ga%) according to Miller et al [13]. Potence ratio estimates were determined according to Smith [14].

**Heterosis was expressed as the deviation of F<sub>1</sub> from the mid-parent and better parent as follows:**

$$\text{Heterosis over mid-parent (MP) \%} = \frac{\bar{F}_1 - \bar{M} \bar{P}}{\bar{M} \bar{P}} \times 100$$

$$\text{Heterosis over better-parent (BP)\%} = \frac{\bar{F}_1 - \bar{B} \bar{P}}{\bar{B} \bar{P}} \times 100$$

**Inbreeding depression:**

$$I. D \% = \frac{\bar{F}_1 - \bar{F}_2}{\bar{F}_1} \times 100$$

To test the significance of inbreeding depression (ID), the variance deviation was calculated as:

$$\text{Variance of (ID) deviation} = \bar{V} \bar{F}_1 + \bar{V} \bar{F}_2$$

## 3. RESULTS AND DISCUSSION

**Mean performance:** The mean performance and variance of traits in six populations (P<sub>1</sub>, P<sub>2</sub>, F<sub>1</sub>, F<sub>2</sub>, BC<sub>1</sub>, and BC<sub>2</sub>) from the first cross are presented in Table 2. The F<sub>1</sub> population showed lower mean values for chocolate spot and rust disease reactions compared to the better parent but had higher plant height, number of pods per plant, number of seeds per plant, seed weight per plant, and 100-seed weight. The F<sub>1</sub> number of branches per plant was intermediate. The F<sub>2</sub> population exhibited lower mean values for all traits compared to F<sub>1</sub>, except for chocolate spot and rust disease reactions. Mean values of BC<sub>1</sub> and BC<sub>2</sub> populations varied relative to F<sub>1</sub> and F<sub>2</sub>. Similar trends were observed in Tables 3 and 4

for crosses 2 and 3, with higher disease reactions. In cross-2,  $F_1$  mean values were lower than the better parent for plant height, number of branches per plant, and 100-seed weight. In cross-3,  $F_1$  values were lower for the number of branches per plant, number of seeds per plant, and 100-seed weight. Genetic variability in the  $F_2$  population was significantly larger than in  $P_1$ ,  $P_2$ , and  $F_1$  for all traits, likely due to genetic segregation.

**Heterosis and inbreeding depression:** Table 5 presents percentages of heterosis, inbreeding depression, and potency ratios for traits in the three crosses. Cross-1 exhibited significant negative mid and better parent heterosis for chocolate spot and rust disease reactions, indicating over-dominance ( $P > +1$ ). Positive over-dominance heterosis was also noted for plant height, number of branches, pods, seeds, and seed weight per plant. Cross-2 showed significant positive mid and better parent heterosis for a number of pods, seeds per plant, and seed yield per plant due to over-dominance ( $P > +1$ ). In Cross-3, significant positive heterosis over mid and better parents was observed for plant height, number of pods, and seed per plant, indicating over-dominance. However, no heterosis benefit was seen for chocolate spot and rust diseases. Positive mid-parent heterosis for the number of branches, number of seeds per plant, and 100-seed weight indicated partial dominance ( $P < +1$ ). Negative better parent heterosis was unfavorable for breeding. Inbreeding depression, calculated as the percent deviation of  $F_2$  from  $F_1$ , showed significant negative values for chocolate spot and rust disease reactions in Cross-1, and positive values for several branches, pods, seeds, and seed weight per plant. Cross-2 had significant negative inbreeding depression for disease reactions and positive values for several seeds and seed weight per plant. Cross-3 showed significant negative inbreeding depression for chocolate spot disease reaction and positive values for several branches per plant and seed weight per plant. Overall, significant negative inbreeding depression was observed for chocolate spot and rust diseases across all crosses, while positive inbreeding depression was noted for yield traits.

**Scaling test:** Generation mean analysis, a quantitative biometrical method, is used to measure the phenotypic performance of quantitative traits across various breeding generations. It helps breeders evaluate the potential for heterosis exploitation or pedigree

selection [15,16]. The significance of at least one scale indicates non-allelic interactions, which can be estimated using a six-parameter model like the Gamble procedure. If all scales are insignificant, a simple additive-dominance model is suitable for estimating genetic components of variance. Table 6 data show that at least one scale (A, B, C, or D) was significant for all traits in the three crosses.

**Type of gene action:** Breeding program success relies on genetic variability in breeding materials. If the additive genetic variance is predominant, the selection is more effective; if the non-additive variance is more significant, an inbred-hybrid program may be better [17]. The analysis provided estimates for six model parameters: (m), (a), (d), (aa), (ad), and (dd) based on Gamble [9]. The estimated mean effects (m) were highly significant for all traits in all crosses (Table 6), indicating quantitative inheritance. Dominant gene effects were significantly higher than additive effects for most traits, except for number of branches per plant in cross-2 and the number of seeds per plant and seed weight in cross-3, where additive genes had a higher impact. Additive effects can be overshadowed by significant dominant estimates when there is a high dispersion of alleles between parents [18]. Dominant effects being larger than additive effects suggest that dominant gene effects play a major role in genetic variance, requiring intensive selection in later generations. Large dominance effects may also explain the significantly better parent heterosis values observed. These conclusions align with findings by many researchers [19,20,21,22,23,24]. Table 6 shows significant additive (a) and dominant (d) genetic variance in cross-1 for flowering date, number of branches, and seed weight per plant; in cross-2 for number of seeds, seed weight per plant, and 100-seed weight; and in cross-3 for number of pods, seed weight per plant, and 100-seed weight. This indicates both additive and dominance in different proportions were involved in trait inheritance. These results are consistent with those reported by El-Refaey and Abd-El-Razek [20], Ibrahim et al [24], El-Refaey et al [25].

Negative values for main effects (a) or (d) or non-allelic interactions (aa, ad, dd) might indicate that alleles responsible for low trait values were over-dominant over those controlling high values. Table 6 data indicate that the aa epistatic effect was more important and higher in magnitude than dd for traits like plant height, number of

branches, seeds, seed weight per plant, and 100-seed weight in cross-1; flowering date, number of pods, seeds, seed weight per plant, and 100-seed weight in cross-2; and chocolate spot and rust disease reactions, number of pods, seed weight per plant, and 100-seed weight in cross-3. Other traits in all crosses had significant and greater dd effects. Traits more affected by aa suggest selection in early segregating generations, while those more affected by dd benefit from delayed selection to later generations. These results are consistent with El-Refaey and Abd-El-Razek [20].

Additive x dominance epistatic gene action was significant and either positive or negative for traits in the three crosses, indicating dominance towards increasing or decreasing, respectively. Ramalingam and Sivasamy [26] noted that a high (ad) epistatic effect for traits like flowering date,

number of seeds, and seed weight per plant in cross-1 and flowering date in cross-3 might suggest delaying selection and inter-mating segregates followed by pedigree selection. Negative (ad) interaction in some crosses for some traits may suggest gene dispersion in parents. Gene action effects where dominance (d) and dominance x dominance (dd) were similar for traits like flowering date, chocolate spot and rust disease reactions, number of pods and seeds per plant in cross-1, and flowering date, number of branches per plant, and seed weight per plant in cross-3, suggest complementary epistatic genes. This indicates considerable heterosis potential for these traits, confirming Table 5 results. El-Refaey and Abd-El-Razek [20] obtained similar results. Similar d and aa gene effects for most traits in all crosses might indicate no complementary non-allelic interaction in genetic control.

**Table 2. Mean performance ( $\bar{X}$ ), variance ( $S^2$ ) and variance of mean ( $S^2\bar{X}$ ) of  $P_1$ ,  $P_2$ ,  $F_1$ ,  $F_2$ ,  $BC_1$  and  $BC_2$  populations of Cross-1 (Santamora x Foal Sbai labiade) for all studied traits**

Trait	Statistical Parameter	$P_1$	$P_2$	$F_1$	$F_2$	$BC_1$	$BC_2$
Flowering date (day)	$\bar{X}$	67.80	68.83	69.27	49.66	61.52	48.81
	$S^2$	2.44	3.45	1.03	41.93	32.56	34.29
	$S^2\bar{X}$	0.08	0.12	0.03	0.14	0.31	0.33
Chocolate spot disease reaction	$\bar{X}$	3.30	3.02	1.83	4.14	4.04	4.02
	$S^2$	0.04	0.03	0.04	1.29	0.95	0.88
	$S^2\bar{X}$	0.001	0.001	0.001	0.004	0.01	0.01
Rust disease reaction	$\bar{X}$	2.30	2.00	1.13	3.46	3.03	2.76
	$S^2$	0.03	0.03	0.02	1.36	0.98	0.81
	$S^2\bar{X}$	0.001	0.001	0.001	0.005	0.01	0.01
Plant height	$\bar{X}$	145.3	128.97	157.0	128.48	139.14	142.19
	$S^2$	4.56	4.65	5.38	101.62	85.35	86.14
	$S^2\bar{X}$	0.15	0.16	0.18	0.34	0.81	0.82
No of branches plant <sup>-1</sup>	$\bar{X}$	4.83	5.07	5.63	3.23	4.78	4.12
	$S^2$	0.13	0.15	0.09	3.05	2.44	2.65
	$S^2\bar{X}$	0.004	0.01	0.003	0.01	0.02	0.03
No of pods plant <sup>-1</sup>	$\bar{X}$	28.83	25.47	44.77	30.52	32.28	31.61
	$S^2$	6.42	5.36	5.56	113.51	105.68	95.45
	$S^2\bar{X}$	0.21	0.18	0.19	0.38	1.01	0.91
No of seeds plant <sup>-1</sup>	$\bar{X}$	98.53	116.57	145.33	96.25	110.96	105.86
	$S^2$	7.09	8.94	11.95	207.98	197.48	178.07
	$S^2\bar{X}$	0.24	0.30	0.40	0.69	1.88	1.70
Seed yield plant <sup>-1</sup> (g)	$\bar{X}$	114.12	132.00	165.80	92.12	119.56	120.91
	$S^2$	6.54	9.41	11.08	203.95	182.77	171.58
	$S^2\bar{X}$	0.22	0.31	0.37	0.68	1.74	1.63
100- seed weight(g)	$\bar{X}$	114.93	113.98	116.45	105.71	115.48	114.08
	$S^2$	9.47	7.61	8.54	156.66	133.43	136.52
	$S^2\bar{X}$	0.32	0.25	0.28	0.52	1.78	1.82

**Table 3. Mean performance ( $\bar{X}$ ), variance ( $S^2$ ) and variance of mean ( $S^2\bar{X}$ ) of P<sub>1</sub>, P<sub>2</sub>, F<sub>1</sub>, F<sub>2</sub>, BC<sub>1</sub> and BC<sub>2</sub> populations of Cross-2 (Santamora x Misr 3) for all studied traits**

Trait	Statistical Parameter	P <sub>1</sub>	P <sub>2</sub>	F <sub>1</sub>	F <sub>2</sub>	BC <sub>1</sub>	BC <sub>2</sub>
Flowering date (day)	$\bar{X}$	68.50	63.00	65.30	56.87	66.43	65.00
	$S^2$	1.43	2.21	1.57	45.67	38.24	34.39
	$S^2\bar{X}$	0.05	0.07	0.05	0.15	0.36	0.33
Chocolate spot disease reaction	$\bar{X}$	2.30	4.00	3.40	4.76	3.34	4.11
	$S^2$	0.02	0.01	0.01	0.56	0.43	0.46
	$S^2\bar{X}$	0.001	0.0003	0.0004	0.002	0.004	0.004
Rust disease reaction	$\bar{X}$	2.13	2.60	3.00	4.37	2.51	3.50
	$S^2$	0.02	0.01	0.01	0.64	0.53	0.42
	$S^2\bar{X}$	0.001	0.0004	0.0004	0.002	0.01	0.004
Plant height	$\bar{X}$	128.83	140.00	138.50	152.38	132.38	144.82
	$S^2$	8.32	6.90	8.19	172.16	133.52	148.12
	$S^2\bar{X}$	0.28	0.23	0.27	0.57	1.27	1.41
No of branches plant <sup>-1</sup>	$\bar{X}$	5.90	3.63	5.30	4.73	4.92	4.04
	$S^2$	0.11	0.12	0.11	3.65	3.48	2.29
	$S^2\bar{X}$	0.004	0.00	0.004	0.01	0.03	0.02
No of pods plant <sup>-1</sup>	$\bar{X}$	25.73	32.03	36.33	33.42	35.22	40.85
	$S^2$	4.96	5.76	5.40	98.25	81.95	84.52
	$S^2\bar{X}$	0.17	0.19	0.18	0.33	0.78	0.80
No of seeds plant <sup>-1</sup>	$\bar{X}$	116.57	94.13	127.15	107.62	123.93	115.39
	$S^2$	7.39	5.77	6.65	105.23	93.25	88.64
	$S^2\bar{X}$	0.25	0.19	0.22	0.35	0.89	0.84
Seed yield plant <sup>-1</sup> (g)	$\bar{X}$	136.34	63.01	148.36	105.22	127.88	116.74
	$S^2$	5.01	4.24	6.28	113.14	91.24	95.13
	$S^2\bar{X}$	0.17	0.14	0.21	0.38	0.87	0.91
100- seed weight(g)	$\bar{X}$	116.69	66.80	111.65	92.87	108.74	96.43
	$S^2$	3.37	5.02	4.75	149.82	129.41	123.74
	$S^2\bar{X}$	0.11	0.17	0.16	0.50	1.73	1.65

**Table 4. Mean performance ( $\bar{X}$ ), variance ( $S^2$ ) and variance of the mean ( $S^2\bar{X}$ ) of P<sub>1</sub>, P<sub>2</sub>, F<sub>1</sub>, F<sub>2</sub>, BC<sub>1</sub> and BC<sub>2</sub> populations of cross 3 (Cairo 4 x Misr 3) for all studied traits**

Trait	Statistical Parameter	P <sub>1</sub>	P <sub>2</sub>	F <sub>1</sub>	F <sub>2</sub>	BC <sub>1</sub>	BC <sub>2</sub>
Flowering date (day)	$\bar{X}$	68.50	61.67	73.00	69.07	70.68	63.81
	S <sup>2</sup>	2.43	1.75	2.21	61.90	42.70	48.04
	S <sup>2</sup> $\bar{X}$	0.08	0.06	0.07	0.21	0.41	0.46
Chocolate spot disease reaction	$\bar{X}$	2.00	4.83	3.80	3.93	3.96	4.28
	S <sup>2</sup>	0.01	0.02	0.02	0.70	0.54	0.51
	S <sup>2</sup> $\bar{X}$	0.0004	0.0007	0.0007	0.002	0.005	0.005
Rust disease reaction	$\bar{X}$	2.17	4.17	3.65	3.44	3.11	4.08
	S <sup>2</sup>	0.01	0.01	0.02	0.66	0.53	0.44
	S <sup>2</sup> $\bar{X}$	0.0004	0.0003	0.0007	0.002	0.01	0.004
Plant height (cm)	$\bar{X}$	128.33	133.67	140.07	147.83	132.76	134.26
	S <sup>2</sup>	9.89	7.13	5.69	157.33	136.22	128.74
	S <sup>2</sup> $\bar{X}$	0.33	0.24	0.19	0.52	1.30	1.23
No of branches plant <sup>-1</sup>	$\bar{X}$	6.10	4.93	5.93	4.75	5.10	4.82
	S <sup>2</sup>	0.12	0.14	0.11	3.44	2.92	3.04
	S <sup>2</sup> $\bar{X}$	0.004	0.00	0.004	0.01	0.03	0.03
No of pods plant <sup>-1</sup>	$\bar{X}$	27.27	25.00	33.37	28.33	30.37	27.60
	S <sup>2</sup>	5.26	4.28	5.41	92.32	79.31	73.58
	S <sup>2</sup> $\bar{X}$	0.18	0.14	0.18	0.31	0.76	0.70
No of seeds plant <sup>-1</sup>	$\bar{X}$	117.27	87.37	115.44	107.33	111.31	94.89
	S <sup>2</sup>	6.55	7.03	5.42	124.97	98.41	106.14
	S <sup>2</sup> $\bar{X}$	0.22	0.23	0.18	0.42	0.94	1.01
Seed yield plant <sup>-1</sup> (g)	$\bar{X}$	138.07	59.42	140.37	95.52	106.90	71.42
	S <sup>2</sup>	7.62	5.98	6.40	128.77	108.45	100.44
	S <sup>2</sup> $\bar{X}$	0.25	0.20	0.21	0.43	1.03	0.96
100- seed weight(g)	$\bar{X}$	117.77	68.21	105.33	96.31	111.12	89.62
	S <sup>2</sup>	4.89	5.43	3.98	129.17	108.89	113.22
	S <sup>2</sup> $\bar{X}$	0.16	0.18	0.13	0.43	1.45	1.51

Table 5. Heterosis over mid-parent (M.P) and better parent (B.P), potance ratio (P) and inbreeding depression (ID) for the studied traits in three faba bean crosses

Trait	M.P %	P	B.P %	ID %
<b>Cross-1 (Santamora x Foal Sbai labiade)</b>				
Flowering date (days)	1.39**	1.84	2.16**	28.30**
Chocolate spot disease reaction	-41.95**	-9.35	-39.23**	-125.64**
Rust disease reaction	-47.29**	-6.78	-43.33**	-205.59**
Plant height	14.49**	2.43	8.05**	18.16
No of branches plant <sup>-1</sup>	13.80**	5.86	11.18**	42.66**
No of pods plant <sup>-1</sup>	64.89**	10.47	55.26**	31.83**
No of seeds plant <sup>-1</sup>	35.13**	4.19	24.68**	33.78**
Seed yield plant <sup>-1</sup> (g)	34.74**	4.78	25.61**	44.44**
100-seed weight(g)	1.74**	4.20	1.32	9.22
<b>Cross-2 (Santamora x Misr 3)</b>				
Flowering date (days)	-0.68*	-0.16	3.65**	12.91
Chocolate spot disease reaction	7.94**	0.45	47.83**	- 40.00**
Rust disease reaction	27.11**	2.71	40.63**	- 45.78**
Plant height	3.04**	0.73	-1.07	-10.02
No of branches plant <sup>-1</sup>	11.19**	0.47	-10.17**	10.75**
No of pods plant <sup>-1</sup>	25.78**	2.36	13.41**	8.01
No of seeds plant <sup>-1</sup>	20.69**	1.94	9.08**	15.36
Seed yield plant <sup>-1</sup> (g)	48.84**	1.33	8.81**	29.08**
100-seed weight(g)	21.70**	0.80	-4.32**	16.82
<b>Cross-3 (Cairo 4 x Misr 3)</b>				
Flowering date (day)	12.16**	2.32	18.38**	5.38
Chocolate spot disease reaction	11.22**	0.27	90.00**	-3.42**
Rust disease reaction	15.26**	0.48	68.46**	5.75**
Plant height	6.92**	3.40	4.78**	-5.54
No of branches plant <sup>-1</sup>	7.55**	0.71	-2.73**	20.00**
No of pods plant <sup>-1</sup>	27.69**	6.39	22.38**	15.10
No of seeds plant <sup>-1</sup>	12.83**	0.88	-1.56*	7.03
Seed weight plant <sup>-1</sup> (g)	42.15**	1.06	1.67*	31.95**
100-seed weight(g)	13.28**	0.50	-10.56**	8.56

\* and \*\*refer to significance at 0.05 and 0.01 levels of probability, respectively



**Table 6. Scaling test and types of gene actions for the studied traits in the three faba bean crosses**

Trait	Scaling test				Types of gene action					
	A	B	C	D	m	a	d	aa	ad	dd
<b>Cross-1 (Santamora x Foal Sbai labiade)</b>										
Flowering date	-14.02**	-40.48**	-76.51**	-11.007**	49.66**	12.71**	22.96**	22.01**	13.23**	32.49**
Chocolate spot	2.94**	3.19**	6.56**	0.216	4.14**	0.02	-1.76**	-0.43	-0.12	-5.70**
Rust disease	2.62**	2.39**	7.29**	1.136**	3.46**	0.27*	-3.29**	-2.27**	0.12	-2.74**
plant height(cm)	-24.01**	-1.59	-74.33**	-24.367**	128.5**	-3.05*	68.60**	48.73**	-11.21**	-23.13**
No of branches plant <sup>-1</sup>	-0.90**	-2.45**	-8.25**	2.445**	3.23**	0.66**	5.57**	4.89**	0.77**	-1.53
No of pods plant <sup>-1</sup>	-9.05**	-7.01**	-21.77**	-2.852	30.52**	0.67	23.32**	5.70	-1.02	10.36
No of seeds plant <sup>-1</sup>	-21.94**	-50.19**	-120.8**	24.328**	96.25**	5.10**	86.44**	48.66**	14.12**	23.47**
Seed weight plant <sup>-1</sup> (g)	-40.80**	-55.98**	-209.3**	-56.239**	92.12**	-1.35	155.2**	112.5**	7.59**	-15.69
100-seed weight	-0.42	-2.27	-38.97**	-18.14**	105.7**	1.40	38.27**	36.28**	0.93	-33.59**
<b>Cross-2 (Santamora x Misr 3)</b>										
Flowering date	-0.94	1.70	-34.63**	17.697**	56.87**	1.43	34.94**	35.39**	-1.32	-36.15**
Chocolate	0.39**	0.83**	5.35**	2.070**	4.76**	-0.77**	-2.99**	-4.14**	-0.22*	2.92**
Rust disease	-0.10	1.39**	6.76**	2.737**	4.37**	-0.98**	-4.84**	-5.47**	-0.75**	4.19**
plant height(cm)	-2.57	11.14**	63.69**	27.559**	152.38**	-12.44**	-51.03**	-55.12**	-6.86**	46.55**
No of branches plant <sup>-1</sup>	-1.36**	-0.86**	-1.21**	0.502	4.73**	0.88**	-0.47	-1.00	-0.25	3.22**
No of pods plant <sup>-1</sup>	8.38**	13.33**	3.25	-9.228**	33.42**	-5.63**	25.90**	18.46**	-2.48	-40.16**
No of seeds plant <sup>-1</sup>	4.15*	9.50**	-34.52**	24.084**	107.62**	8.54**	69.97**	48.17**	-2.67*	-61.82**
Seed weight plant <sup>-1</sup> (g)	-28.94**	22.11**	-75.19**	-34.180**	105.22**	11.14**	117.04**	68.36**	-25.53**	-61.53**
100-seed weight	-10.86**	14.41**	-35.30**	19.43**	92.87**	12.31**	58.77**	38.86**	-12.63**	-42.42**
<b>Cross-3 (Cairo 4 x Misr 3)</b>										
Flowering date	-0.14	-7.05**	0.11	3.650**	69.07**	6.87**	0.62	-7.30**	3.45**	14.49**
Chocolate	2.12**	-0.08	1.29**	0.378**	3.93**	-0.31**	1.14**	0.76**	1.10**	-2.80**
Rust disease	0.41**	0.34*	0.13	-0.310*	3.44**	-0.96**	1.10**	0.62*	0.04	-1.37**
plant height(cm)	-2.88	-5.21*	49.20**	28.647**	147.83**	-1.50	-48.23**	-57.29**	1.17	65.39**
No of branches plant <sup>-1</sup>	-1.82**	-1.23**	-3.91**	0.430	4.75**	0.29	1.28*	0.86	-0.30	2.19*
No of pods plant <sup>-1</sup>	1.11	-4.17*	-5.69**	1.310	28.33**	0.77*	11.85**	6.62	-0.36	-11.56
No of seeds plant <sup>-1</sup>	-10.09**	-13.03**	-6.19*	8.460**	107.3**	16.42**	-3.80	-16.92**	1.47	40.03**
Seed weight plant <sup>-1</sup> (g)	-64.64**	-56.95**	-96.15**	12.716**	95.52**	35.48**	16.19**	-25.43**	-3.84**	147.01**
100-seed weight	-0.86	5.70**	-11.39**	8.120**	96.31**	21.50**	28.58**	16.24**	-3.28*	-21.09**

\* and \*\*refer to not significant and significant at 0.05 and 0.01 levels of probability, respectively

**Table 7. Heritability percentage in broad (H), narrow (h<sup>2</sup>) senses and expected (Ga) and predicted (Ga %) genetic advance for selection in the three crosses of faba bean for the studied traits**

Traits	H	h <sup>2</sup>	Ga	Ga%
Flowering date (days)	95.26	40.57	5.41	10.90
Chocolate spot disease reaction	97.09	58.14	1.36	32.88
Rust disease reaction	98.06	68.37	1.64	47.42
Plant height	95.09	31.25	6.49	5.05
No of branches plant <sup>-1</sup>	96.23	33.11	1.19	36.88
No of pods plant <sup>-1</sup>	94.96	22.81	5.01	16.40
No of seeds plant <sup>-1</sup>	95.20	19.43	5.77	6.00
Seed weight plant <sup>-1</sup> (g)	95.33	26.26	7.72	8.39
100-seed weight(g)	94.55	27.68	7.14	6.75
Cross-2 (Santamora x Misr 3)				
Flowering date (days)	96.29	40.96	5.70	10.03
Chocolate spot disease reaction	97.32	41.07	0.63	13.29
Rust disease reaction	97.77	51.56	0.85	19.43
Plant height	95.41	36.41	9.84	6.46
No of branches plant <sup>-1</sup>	96.88	42.09	1.66	35.02
No of pods plant <sup>-1</sup>	94.52	30.56	6.24	18.67
No of seeds plant <sup>-1</sup>	93.71	27.15	5.74	5.33
Seed weight plant <sup>-1</sup> (g)	95.18	35.27	7.73	7.35
100-seed weight(g)	97.01	31.03	7.82	8.43
Cross-3 (Cairo 4 x Misr 3)				
Flowering date (days)	96.53	53.41	8.66	12.53
Chocolate spot disease reaction	97.46	50.43	0.87	22.12
Rust disease reaction	97.62	54.17	0.91	26.39
Plant height	95.49	31.59	8.16	5.52
No of branches plant <sup>-1</sup>	96.51	26.74	1.02	21.53
No of pods plant <sup>-1</sup>	94.49	34.39	6.81	24.03
No of seeds plant <sup>-1</sup>	95.11	36.32	8.36	7.79
Seed weight plant <sup>-1</sup> (g)	94.87	37.78	8.83	9.25
100-seed weight(g)	96.46	28.05	6.57	6.82

**Heritability and genetic advance from selection:** Heritability estimates based on genetic variance components (a) and (d) are listed in Table 7. Broad-sense heritability for all traits ranged from 93.71% in Cross-2 for number of seeds per plant to 98.06% in Cross-1 for rust disease reaction. These high values suggest that superior genotypes for these traits can be identified phenotypically, emphasizing the importance of phenotypic selection [27]. The results also indicate that traits were minimally affected by environmental factors, with most variability due to genetics. Ibrahim et al [24] reported broad-sense heritability in faba bean ranging from 49.45% to 91.29%.

Narrow-sense heritability values ranged from 19.43% for the number of seeds per plant to 68.37% for rust disease reaction in Cross-1, from 27.15% for number of seeds per plant to 51.56% for rust disease reaction in Cross-2, and from 26.74% for the number of branches per plant to 54.17% for rust disease reaction in Cross-3. Moderate h<sup>2</sup> values (40.57% to 58.14%) were

observed for traits like flowering date and chocolate spot and rust disease, which are somewhat qualitative. In contrast, for quantitative traits like plant height, number of branches, pods, seeds, seed weight per plant, and 100-seed weight, h<sup>2</sup> values ranged from 19.43% to 42.09%, considered low. These results are expected as recent genetic material is controlled by non-additive gene effects. Ibrahim et al [24] reached similar conclusions.

Genetic advance under selection, as shown in Table 7, indicated expected genetic advance (Ga) ranged from 1.19 for number of branches per plant to 7.72 g for seed weight in Cross-1, from 0.63 for chocolate spot disease reaction to 9.84 cm for plant height in Cross-2, and from 0.87 for chocolate spot disease reaction to 8.83 g for seed weight in Cross-3. The highest predicted genetic advance (Ga%) values were coupled with moderate h<sup>2</sup> values across all traits in all crosses. Conversely, low expected (Ga) and predicted (Ga%) advance estimates were associated with low h<sup>2</sup> values for number of

seeds per plant, seed weight per plant, and 100-seed weight in all three crosses.  $Ga\%$  values ranged from 5.05% for plant height to 47.42% for rust disease reaction in Cross-1, from 5.33% for number of seeds per plant to 35.02% for number of branches per plant in Cross-2, and from 5.52% for plant height to 26.39% for rust disease reaction in Cross-3. These results align with Johnson et al [12], who reported that heritability estimates combined with genetic advance upon selection are more valuable for predicting selection effects. These findings support selection in late generations ( $F_4$  and  $F_5$ ) through the bulk method to obtain high-yielding faba bean genotypes with improved disease resistance. These results are consistent with those of El-Refaey and Abd-El-Razek [20] and Fouad [23].

#### 4. CONCLUSION

This study successfully introduced high-yielding, disease-resistant faba bean genotypes by analyzing six populations from three crosses over three growing seasons. Significant heterosis and high broad-sense heritability values indicated that phenotypic selection is effective for improving these traits. Dominant gene effects were more influential than additive effects, highlighting the need for intensive selection in later generations. The presence of non-allelic interactions necessitates specific breeding strategies. These findings support the feasibility of developing superior faba bean genotypes through targeted breeding, providing valuable insights for future breeding programs.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Authors have declared that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Lawes DS, Bond DA, Poulsen MH. Classification, origin, breeding methods and objectives in the faba bean (*Vicia faba* L.). P. D. Hebblethwaite (ed.), Butterworths, London. 1983;32-76.
2. Eliw M. Economic analysis of supply response of broad beans crop in Egypt. New Valley Journal of Agricultural Science. 2021;1(1):10-25.
3. Heiba HE, Mahgoub E, Mahmoud A, Ibrahim M, Mahdy EMB. Genetic improvement of faba bean (*Vicia faba* L.) genotypes selected for resistance to chocolate spot disease. Agronomic Colombian. 2022;40(2):186-197.
4. El-Hady MM, Sabah M, Attia AAM, Ashrei EAA, El-Emam, Shaaban M. Heterosis, combining ability and genetic variability and some related components in faba bean using six populations model. Minia Journal of Agric. Res. and Develop. 2009;29(3):417-430.
5. Abdallah MMF, Shafik MM, Sabah M Attia, Hend A Ghannam. Heterosis, GCA and SCA effects of diallel cross among six faba bean (*Vicia faba* L) genotypes. ARIA. 2017;4(4):1-10.
6. Bernier CC, Hanounik SB, Hussein MM, Mohamed HA. Field manual of common faba bean diseases in the Nile Valley. International Center for Agricultural Research in the Dry Areas (ICARDA) Information Bulletin No. 1993;3.
7. Mather K. Biometrical Genetics. Dover Publications, Inc., London; 1949.
8. Hayman BI. The theory and analysis of diallel crosses. Genetics. 1954;39:789-809.
9. Gamble EE. Gene effects in corn (*Zea mays* L.). I-Separation and relative importance of gene effects for yield. Canadian Journal of Plant Science. 1962; 42:330-335.
10. Allard RW. Principles of plant breeding. Jon. Wiley and Sons. Inc. NY, USA. 1962; 92.
11. Warner JN. A method for estimating heritability. Agron. J. 1952;44:427-430.
12. Johnson HW, Robinson HF, Comstock RE. Estimating of genetic and environmental variability in soybeans. Agron. J. 1955;47: 314-318.
13. Miller PA, Williams JC, Robinson HF, Comstock RE. Estimates of genotypic and environmental variances and covariance in upland cotton and their implications in selection. Agron. J. 1958;50:126-131
14. Smith HH. Fixing transgressive vigor in *Nicotiana rustica*. in gowen heterosis, Iowa State Collage Press. Anes, Iowa, U. S. A; 1952.

15. Kearsey MJ, Pooni HS. The Genetic Analysis of Quantitative Traits. Chapman and Hall, 1<sup>st</sup> edition, London. 1996;46.
16. Sharma SN, Sain RS. Genetic architecture of grain weight in durum wheat under normal and late sown environments. Wheat Information Service. 2003;96:28-32.
17. Cockerhan CC. Implication of genetic variance in hybrid breeding program. Crop Sci. 1961;1:52-74.
18. Snap JW. Conventional methods of genetic analysis in wheat. In Hupton F.G. H(ed) wheat breeding its scientific basis, Chapman and Hall, London, New York. 1987;109-128.
19. El-Refaey RA. Generation mean analysis of earliness, yield and its components in three faba bean crosses. Menoufia J. Agric. Res. 1999;24:409-424.
20. El-Refaey RA, Abd-El-Razek UA. Generation mean analysis for yield and its components in four crosses of Egyptian cotton (*Gossypium Barbadians* L.). Asian Journal of Crop Science. 2013;5(2):153-166.
21. Haridy MH, El-Said MAA. Estimation of genetic parameters using populations in faba bean (*Vicia faba* L.) J. Plant production, Mansoura Univ. 2016;7:1443-1447.
22. Abou-Zaid Gehan GA. Genetical analysis of yield, its components for three faba bean crosses using six parameters model. Menoufia Journal Plant Production. 2018;3:121 – 135.
23. Fouad HM. Six generation mean analysis using scaling and joint scaling tests in Faba bean (*Vicia faba* L.). Journal of Sustainable Sciences. 2020;46(1):1-11.
24. Ibrahim MA, Gehan G, Abo-Zaid, Salwa M Mostafa. Estimation of some genetic parameters using six populations mean analysis in three faba bean crosses exposed to natural infection of foliar diseases. Egyptian J. Agric. Res. 2023; 101(2):843-854.
25. El-Refaey RA, El-Moslhy MA, El-Gammaal AA, El-Naggar AA. Generation mean analysis for yield and its components in five crosses of barley (*Hordeum vulgare* L.) under water stress conditions. J. Agric. Res. Kafr El-Sheikh Univ. 2015;4: 670-683.
26. Ramalingam A, Sivasamy N. Genetics and order effects of seed cotton yield in upland cotton (*Gossypium hirsutum* L.) diallel analysis. Indian J. Genetics. 2002; 62(4):359-360.
27. Awaad HA. Diallel analysis of yield and its contributing characters in wheat (*Triticum sativum* L.) Zagazig J. Agric. Res. 1996; 23(6):999-1012.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/119770>