



# Pre- Harvest Foliar Spray of Zinc, Calcium, Chitosan, and Salicylic Acid on Quality Standards of Indian Ber (*Zizyphus mauritiana* Lamk.) cv. Gola

Mukesh Chand Bhatেশwar <sup>a\*</sup>, S. K. Khandelwal <sup>a</sup>,  
Deepak Gupta <sup>b</sup>, Pooja Sharma <sup>a</sup> and Kamal Mahala <sup>a</sup>

<sup>a</sup> Department of Horticulture, SKNAU, Jobner, Jaipur 303 329, Rajasthan, India.

<sup>b</sup> Department of Genetics and Plant Breeding, SKNAU, Jobner, Jaipur 303 329, Rajasthan, India.

## Authors' contributions

This work was carried out in collaboration among all authors. Authors MCB and SKK designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. Author DG, PS and KM managed the analyses and literature searches of the study. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/jsrr/2024/v30i102513>

**Open Peer Review History:**  
This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/125178>

Original Research Article

Received: 12/08/2024

Accepted: 16/10/2024

Published: 18/10/2024

## ABSTRACT

**Aim:** The present experiment was designed to find out the effect of a foliar spray of zinc, calcium, chitosan and salicylic acid on the fruit quality of ber.

**Place and Duration of Study:** This experiment was conducted at Horticulture farm, SKN College of Agriculture, Jobner in the newly established orchard of Indian jujube cv. Gola during the two-consecutive years 2022 and 2023.

\*Corresponding author: E-mail: [mukeshchandbhatেশwar@gmail.com](mailto:mukeshchandbhatেশwar@gmail.com);

**Cite as:** Bhatেশwar, Mukesh Chand, S. K. Khandelwal, Deepak Gupta, Pooja Sharma, and Kamal Mahala. 2024. "Pre- Harvest Foliar Spray of Zinc, Calcium, Chitosan, and Salicylic Acid on Quality Standards of Indian Ber (*Zizyphus Mauritiana* Lamk.) Cv. Gola". *Journal of Scientific Research and Reports* 30 (10):921-27. <https://doi.org/10.9734/jsrr/2024/v30i102513>.

**Study Design and Methodology:** The present experiment is laid out in factorial randomized block design with three replications. Factor A consists of 12 level of zinc, calcium, chitosan, and salicylic acid treatments along with the control, and factor B consists three foliar sprays i.e., single foliar spray at pea size stage, color turning stage of fruits and double foliar spray at pea size and color turning stage of fruits (green to yellow). Various quality parameters namely TSS, acidity, ascorbic acid, total, reducing and non-reducing sugars are observed as per standard procedures.

**Results:** Plants sprayed with  $ZnSO_4 @ 4g/L + CaCl_2 @ 2g/L + Chitosan @ 1.5g/L + Salicylic acid @ 1g/L$  recorded highest TSS (19.05°Brix), minimum acidity (0.415%) TSS: acid ratio (45.93%), ascorbic acid (89.08mg/100g), total sugars (10.25%), reducing sugars (6.68%) and non-reducing sugars (3.39%) were significantly more in the treated fruits in comparison to untreated fruits. These quality parameters were observed maximum with a double foliar spray of nutrients and agrochemicals at the pea size stage and color turning stage ( $S_3$ ).

**Conclusion:** It is concluded that  $ZnSO_4 @ 4g/L + CaCl_2 @ 2g/L + Chitosan @ 1.5g/L + Salicylic acid @ 1g/L$  is effective in improve the quality of ber fruits.

**Keywords:** Zinc; calcium; chitosan; salicylic acid; quality.

## 1. INTRODUCTION

Ber (*Zizyphus mauritiana* Lamk.) is one of the most ancient and common fruits of India and it is cultivated all over the country. Being a hardy fruit, it can be grown even on inferior and marginal lands. Ber is indigenous to an area stretching from India to China and Malaysia. Because of good returns from ber, its cultivation is becoming increasingly popular in Gujarat and other states of India, especially in Punjab, Haryana, and Rajasthan. Gola is the most common cultivar being farmed [1]. The Gola variety is a well-liked early type. The fruit has a medium size and an oblong to spherical shape. Its color changes from green to golden yellow as it ripens. Fruit typically weighs between 16 and 20 g on average [2]. A huge loss of horticultural produce takes place during storage. Inadequate means of losses during storage due to poor handling and poor post-harvest management practices. Ber can be cultivated even under constraints of irrigation and fertilization. The essential element of fruit production is bioregulators. These regulators can be applied to fruits pre-harvest to improve their physical and biochemical characteristics and preserve quality.

Zinc is an essential micro element for plants, being involved in many enzymatic reactions and is necessary for their good growth and development. It is also involved in regulating protein and carbohydrate metabolism. Its availability to plants is reduced in high- pH soils. Fruits generally have a considerable amount of Calcium in their tissues. The Ca application makes the middle lamella of the fruit cell wall thicker by increased deposition of calcium pectate and so, maintains the firmness of the fruit

(Sudha et al. 2007). Calcium has long been associated with regulation of the ripening and post-harvest storage life of fruits. Specially, maintenance of relatively high ca concentrates in fruit tissue result in a slower rate of ripening, as seen in lower respiration rate, reduced ethylene production and slower softening of fruit flesh. Combined with pectin, the “glue” that holds plant cells together, calcium forms a pectate salt that helps to keep the cell wall sturdy and rigid. Chitosan applied topically at various phases of fruit development had a major impact on the postharvest life of jujube fruits during storage. Treatments with chitosan function as barriers, preventing the transfer of water, shielding fruit skin from mechanical damage, and sealing minor wounds to postpone dehydration [3]. Salicylic acid is an additional phenolic endogenous growth regulator that helps control a number of physiological functions in plants, including transpiration, ion absorption, stomata closure, and the suppression of ethylene production [4]. Keeping in mind the importance of nutrients and agrochemicals current study was planned to improve the quality of ber fruit by pre-harvest exogenous spray of different concentrations of zinc, calcium, chitosan and salicylic acid at various growth stages under semi-arid condition of Rajasthan.

## 2. MATERIALS AND METHODS

### 2.1 Experimental Site and Treatments Detail

This experiment was conducted at Horticulture farm, SKN College of Agriculture, Jobner in the newly established orchard of Indian jujube (*Zizyphus mauritiana* Lamk.) cv. Gola during

2022 and 2023. Factor A consists of 12 level of zinc, calcium, chitosan, and salicylic acid treatments along with the control, and factor B consists of three foliar sprays i.e., single foliar spray at pea size stage, color turning stage of fruits and double foliar spray at pea size and colour turning stage of fruits (green to yellow). The treatment details of zinc, calcium, chitosan and salicylic acid along with the control i.e., T<sub>0</sub> (Control), T<sub>1</sub> (Zinc sulfate @ 4 g / L), T<sub>2</sub> (Calcium chloride @ 2 g / L), T<sub>3</sub> (Chitosan @ 1.5 g / L), T<sub>4</sub> (Salicylic acid @ 1 g / L), T<sub>5</sub> (Zinc sulphate @ 4 g / L + Calcium chloride @ 2 g / L), T<sub>6</sub> (Zinc sulfate @ 4 g / L + Chitosan @ 1.5 g / L), T<sub>7</sub> (Zinc sulfate @ 4 g / L + Salicylic acid @ 1 g / L), T<sub>8</sub> (Calcium chloride @ 2 g / L + Chitosan @ 1.5 g / L), T<sub>9</sub> (Calcium chloride @ 2 g / L + Salicylic acid @ 1 g / L), T<sub>10</sub> (Chitosan @ 1.5 g / L + Salicylic acid @ 1 g / L) and T<sub>11</sub> (Zinc sulfate @ 4 g / L + Calcium chloride @ 2 g / L + Chitosan @ 1.5 g / L + Salicylic acid @ 1 g / L) laid out in factorial randomized block design with three replications. The treatments were applied with single foliar spray at the pea size stage of fruits during first week of November and double foliar spray at pea size and color turning stage of fruits (green to yellow) applied last week of December, 2022-23 and 2023-24 after recording initial (base) quality and parameters of ber fruits.

## 2.2 Quality Attributes of Ber Fruit

For determination of total soluble solids was measured by placing a drop of juice on the prism of the hand refractometer. The reading against the light was noted and presented in degree brix. For estimation of ascorbic acid, well-homogenized 2 g of sample was taken in a beaker volume made up to 20 ml and 5 ml of HPO<sub>3</sub> is added and titrated with the standard dye solution to a pink end-point which should persist for 15 seconds. The same procedure was repeated in random samples of fruits in both the year of an experiment to draw an average value of ascorbic acid content. The ascorbic acid content of the sample was calculated by using the following formula [5].

Ascorbic acid (mg/100g) = (Titre value x dye factor x volume made up) / (Aliquot extract taken for estimation x weight of sample taken for Estimation) x 100

Total and reducing sugars were estimated by the lane and Eynon copper titration method and expressed in percentage [6]. The percentage of non-reducing sugar was determined by

subtracting the percent reducing sugar from the percent total sugar and multiplying the same with 0.95 as given below [7].

Non-reducing sugars (%) = (Per cent total sugar – per cent reducing sugar) x 0.95

## 2.3 Statistical Analysis

The data were statistically analyzed as per the analysis of variance technique as suggested by Panse et al. [8]. The significance of the treatments was tested through the F test at a 5 percent level of significance. The critical difference CD was calculated to assess the significance of differences among the different treatments.

## 3. RESULTS AND DISCUSSION

Pre- harvest foliar spray of zinc, calcium, chitosan, and salicylic acid on quality standards viz., TSS, titratable acidity, TSS: acid ratio and ascorbic acid of ber (*Zizyphus mauritiana* Lamk.) cv. Gola is presented in (Table 1).

The maximum TSS content in fruits (19.05%) was recorded with the application of treatment (ZnSO<sub>4</sub> @ 4g/L + CaCl<sub>2</sub> @ 2g/L + Chitosan @ 1.5g/L + Salicylic acid @ 1g/L) followed by T<sub>9</sub> (CaCl<sub>2</sub> @ 2g/L + Salicylic acid @ 1g/L 18.90%), T<sub>7</sub> (ZnSO<sub>4</sub> @ 4g/L + Salicylic acid @ 1g/L 18.83%), T<sub>5</sub> (ZnSO<sub>4</sub> @ 4g/L + CaCl<sub>2</sub> @ 2g/L 18.69%) and minimum TSS content in fruits (16.80) was noted under control. The maximum total soluble solid content (18.05 °Brix) was observed with double foliar spray of nutrients and agrochemicals at the pea size stage and color turning stage (S<sub>3</sub>) and it was found at par with foliar spray at the S<sub>1</sub> stage. The foliar spray of nutrients and agrochemicals at the color-turning stage (S<sub>2</sub>) registered the lowest total soluble solid (17.38 °Brix). A significant decrease in titratable acidity was observed in ZnSO<sub>4</sub> @ 4g/L + CaCl<sub>2</sub> @ 2g/L + Chitosan @ 1.5g/L + Salicylic acid @ 1g/L (0.16%) followed by the T<sub>9</sub>, T<sub>7</sub> and T<sub>5</sub> and highest titratable acidity was observed in fruits of controlled treatment (0.469%). The application of nutrients and agrochemicals at different fruit development stages had not shown any significant effect on the titratable acidity in ber fruits.

The maximum TSS: acid ratio (45.93) was recorded with combined application of ZnSO<sub>4</sub> @ 4 g/l + CaCl<sub>2</sub> @ 2 g/l + Chitosan @ 1.5 g/l + Salicylic acid @ 1 g/l (T<sub>11</sub>) followed by this treatment T<sub>9</sub>, T<sub>7</sub> and T<sub>5</sub>. However, a minimum

TSS: acid ratio of (36.32) was recorded under control. Application of a double foliar spray of nutrients and agrochemicals exhibited maximum TSS: acid ratio (41.69) in fruits. However, this treatment (S<sub>3</sub>) was found statistically at par with treatment S<sub>1</sub>. The foliar spray at S<sub>2</sub> recorded the lowest TSS: acid ratio (38.90) during the experimentation.

ZnSO<sub>4</sub> @ 4g/L + CaCl<sub>2</sub> @ 2g/L + Chitosan @ 1.5g/L +Salicylic acid @ 1g/L significantly increase the ascorbic acid content of ber fruits. Highest ascorbic acid content was observed in fruits treated with ZnSO<sub>4</sub> @ 4g/L + CaCl<sub>2</sub> @ 2g/L + Chitosan @ 1.5g/L +Salicylic acid @ 1g/L that's 89.08 mg/100g followed by T<sub>9</sub> (85.77 mg/100g), T<sub>7</sub> (85.63 mg/100g) and T<sub>5</sub> (85.31 mg/100g). The lowest contents were observed in control (74.63 mg/100g). However, the maximum ascorbic acid content (83.42 mg/100g) was observed with a double foliar spray of nutrients and agrochemicals at the pea size stage and color turning stage (S<sub>3</sub>) and it was found statistically at par with S<sub>1</sub> (82.75 mg/100g). The treatment combination S<sub>2</sub> recorded minimum ascorbic acid content (81.96 mg/100g) during the investigation.

It is evident from data presented in Table 2 related to the application of nutrients and agrochemicals showed a significant effect on

total, reducing and non-reducing sugars in fruits. Significantly higher total sugars was observed in ZnSO<sub>4</sub> @ 4g/L + CaCl<sub>2</sub> @ 2g/L + Chitosan @ 1.5g/L +Salicylic acid @ 1g/L treatment (10.25%). Controlled treatment shows minimum total sugars among other treatments (8.16%). However, the highest total sugars of (9.54 %) were observed with a double foliar spray of nutrients and agrochemicals at the pea size stage and colour turning stage (S<sub>3</sub>) and it was found statistically at par with treatment S<sub>1</sub> (9.32 %). The treatment combination S<sub>2</sub> recorded the lowest total sugars content (8.96 %) during the experimentation.

ZnSO<sub>4</sub> @ 4g/L + CaCl<sub>2</sub> @ 2g/L + Chitosan @ 1.5g/L +Salicylic acid @ 1g/L significantly increases reducing and non-reducing sugars (6.52%, 4.23% respectively) followed by the fruits treated with T<sub>9</sub>, T<sub>7</sub> and T<sub>5</sub> and minimum values were observed in the control treatment. However, the maximum reducing (6.46%) and non-reducing (3.01%) sugars were observed with a double foliar spray of nutrients and agrochemicals at the pea size stage and color turning stage (S<sub>3</sub>) and this treatment (S<sub>3</sub>) was found statistically at par with treatment S<sub>1</sub>. Whereas, minimum reducing (6.17%) and non-reducing (2.56%) sugars were recorded with a foliar spray of nutrients and agrochemicals done at the color-turning stage (S<sub>2</sub>).

**Table 1. Pre- harvest foliar spray of zinc, calcium, chitosan and salicylic acid on TSS, titratable acidity, TSS: acid ratio and ascorbic acid of ber fruit**

Treatments	TSS (°Brix)	Titratable acidity (%)	TSS: acid ratio	Ascorbic acid (mg/100g)
<b>Nutrients and agrochemicals</b>				
T <sub>0</sub> Control	16.80	0.469	36.32	74.63
T <sub>1</sub> ZnSO <sub>4</sub>	16.90	0.457	37.03	80.38
T <sub>2</sub> CaCl <sub>2</sub>	16.94	0.457	37.15	80.72
T <sub>3</sub> Chitosan	16.83	0.465	36.26	76.44
T <sub>4</sub> Salicylic acid	16.94	0.447	37.96	81.15
T <sub>5</sub> ZnSO <sub>4</sub> + CaCl <sub>2</sub>	18.69	0.421	44.47	88.07
T <sub>6</sub> ZnSO <sub>4</sub> + Chitosan	17.43	0.448	38.97	81.53
T <sub>7</sub> ZnSO <sub>4</sub> + Salicylic acid	18.83	0.418	45.05	88.32
T <sub>8</sub> CaCl <sub>2</sub> + Chitosan	17.77	0.450	39.57	81.65
T <sub>9</sub> CaCl <sub>2</sub> + Salicylic acid	18.90	0.417	45.36	88.80
T <sub>10</sub> Chitosan+ Salicylic acid	17.94	0.434	41.55	81.77
T <sub>11</sub> ZnSO <sub>4</sub> + CaCl <sub>2</sub> + Chitosan + SA	19.05	0.415	45.93	89.08
<b>SEm ±</b>	<b>0.21</b>	<b>0.00</b>	<b>0.61</b>	<b>0.48</b>
<b>CD (P = 0.05)</b>	<b>0.59</b>	<b>NS</b>	<b>1.70</b>	<b>1.35</b>
<b>Different Stage of Foliar Spray</b>				
S <sub>1</sub>	17.83	0.440	40.81	82.75
S <sub>2</sub>	17.38	0.449	38.90	81.96
S <sub>3</sub>	18.05	0.435	41.69	83.42
<b>SEm±</b>	<b>0.11</b>	<b>0.00</b>	<b>0.30</b>	<b>0.24</b>
<b>CD (P = 0.05)</b>	<b>0.29</b>	<b>NS</b>	<b>0.85</b>	<b>0.68</b>

**Table 2. Pre- Harvest foliar spray of zinc, calcium, chitosan and salicylic acid on total, reducing and non-reducing sugars of ber fruit**

Treatments	Total sugars (%)	Reducing sugars (%)	Non-reducing sugars (%)
<b>Nutrients and agrochemicals</b>			
T <sub>0</sub> Control	8.16	5.92	2.13
T <sub>1</sub> ZnSO <sub>4</sub>	8.84	6.15	2.56
T <sub>2</sub> CaCl <sub>2</sub>	8.92	6.23	2.55
T <sub>3</sub> Chitosan	8.64	6.07	2.44
T <sub>4</sub> Salicylic acid	9.01	6.19	2.68
T <sub>5</sub> ZnSO <sub>4</sub> + CaCl <sub>2</sub>	9.93	6.56	3.21
T <sub>6</sub> ZnSO <sub>4</sub> + Chitosan	9.01	6.24	2.64
T <sub>7</sub> ZnSO <sub>4</sub> + Salicylic acid	10.05	6.58	3.30
T <sub>8</sub> CaCl <sub>2</sub> + Chitosan	9.10	6.28	2.68
T <sub>9</sub> CaCl <sub>2</sub> + Salicylic acid	10.16	6.63	3.35
T <sub>10</sub> Chitosan+ Salicylic acid	9.17	6.31	2.71
T <sub>11</sub> ZnSO <sub>4</sub> + CaCl <sub>2</sub> + Chitosan + SA	10.25	6.68	3.39
<b>S<sub>Em</sub> ±</b>	<b>0.17</b>	<b>0.10</b>	<b>0.17</b>
<b>CD (P = 0.05)</b>	<b>0.47</b>	<b>0.27</b>	<b>0.48</b>
<b>Different Stage of Foliar Spray</b>			
S <sub>1</sub>	9.32	6.33	2.84
S <sub>2</sub>	8.96	6.17	2.56
S <sub>3</sub>	9.54	6.46	3.01
<b>S<sub>Em</sub>±</b>	<b>0.08</b>	<b>0.05</b>	<b>0.09</b>
<b>CD (P = 0.05)</b>	<b>0.23</b>	<b>0.13</b>	<b>0.24</b>

The application of zinc, CaCl<sub>2</sub>, chitosan and SA increased TSS content in the fruits, which might be due to the enhancement of auxin synthesis. Auxin, in turn, enhances the synthesis of metabolites and their rapid translocation from one part of the plant to developing fruits [9,10] (Badway et al., 2019). The improvement in fruit quality might also be due to zinc's role in enzymatic reactions, such as hexokinase, and its involvement in carbohydrate formation and protein synthesis [11,12]. The reason for decreased TA is due to the breakup of fermentation of acids into sugars during the respiration process. An increase in the TSS: acid ratio is likely due to an increase in TSS content and a decrease in acidity [13,14]. Zinc, CaCl<sub>2</sub>, chitosan and SA application also increased the ascorbic acid content, which appears to be due to increased growth and the availability of more metabolites for ascorbic acid synthesis. The possible reason for the increase in total sugar and reducing sugars in fruits from foliar sprays of minerals and agrochemicals might be the conversion of starch and acid into sugars, along with the continuous mobilization of sugars from leaves to fruits [15,16] (Martínez et al., 2022). Treatments that delayed the start of fruit senescence and the conversion of complex carbohydrates into simple sugars may have contributed to the greater non-reducing sugar levels in treated samples [17,18,19,20].

#### 4. CONCLUSION

The findings of this study demonstrate that as compared with individual applications of zinc, calcium, chitosan, or salicylic acid, a combination of these elicitors had a much superior impact on fruit quality aspects of ber fruits. The present study is noteworthy because it offers evidence that a combination of these elicitors may enhance fruit quality both during the postharvest storage period and at the time of harvest. Future research should concentrate on optimizing the concentrations and combinations that might be used commercially on ber fruit from different crops or cultivars to increase the fruit's marketability.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

#### ACKNOWLEDGEMENTS

Authors are highly thankful to Department of Horticulture, Biochemistry, SKN Agriculture University, Jobner (Rajasthan) for providing all the necessary facilities and kind support.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Surani, BR., Patel, AP. and Yadav, L. Effect of post-harvest application of chemicals on bio-chemicals parameters of different Ber (*Ziziphus mauritiana* Lamk.) varieties during storage. International Journal of Chemical Studies. 2018;6(4): 2804-280
2. Chattopadhyay TK. A Text Book on Pomology (Subtropical Fruits). Kalyani Publishers, New Delhi; 2012.
3. Ribeiro C, Vicente AA, Teixeira JA, Miranda C. Optimization of edible coating composition to retard strawberry fruit senescence. Post-Harvest Biology and Technology. 2007;44(1):63-70.
4. Shafiee M, Taghavi TS, Babalar M. Addition of salicylic acid to nutrient solution combined with post-harvest treatments (hot water, salicylic acid, and calcium dipping) improved post-harvest fruit quality of strawjuber. Scientia Horticulturae. 2010;124(1):40-45.
5. Ranganna K, Yatsu FM, Hayes BE, Milton SG, Jayakumar A. Butyrate inhibits proliferation-induced proliferating cell nuclear antigen expression (PCNA) in rat vascular smooth muscle cells. Molecular and Cellular Biochemistry. 2000;205:149-161.
6. Ranganna, S. Handbook of analysis and quality control for fruit and vegetable products. Tata McGraw-Hill Education. 1986;670.
7. Somogyi M. Notes on sugar determination. Journal of Biological Chemistry, 1952;195: 19-23.
8. Panse VG, Sukhatme PV. Statistical methods for agricultural workers, 4th edition. ICAR, New Delhi. 1995;58-92.
9. Alrawi W, Jassim NA, Al-Hadethi ME. Effect of iron and zinc spray on yield, growth and content of the seeds of apricot amygdalin. Egyptian Journal of Applied Sciences. 2012;27(4):173-187.
10. Talat K, Abdel-Salam MM, Radwan EMA, Haleem AY, Mahmoud AA. Effects of some pre-and post-harvest treatments on physical and chemical properties of flame seedless grapevines during cold storage. New Valley Journal of Agricultural Science. 2024;2805-2420.
11. Pamila S, Chatterjee SR, Deb DL. Seed yield, harvest index, protein content and amino acid composition of chickpea as affected by sulphur and micronutrients. Annals of Agriculture Research. 1992;3(1): 7-11.
12. Wahab M, Ullah Z, Usman M, Nayab S, Ullah M. Effect of three ca-sources applications on fruit quality attributes of three peach cultivars in Pakistan. Pure and Applied Biology. 2021;5(3):464-470.
13. Abdel Rhman IE, Attia MF, Elhady ES, Haggag LF. Effect of foliar spraying of some antioxidants and micronutrients on yield, fruit quality and leaf mineral content of manfaloty pomegranate trees (*Punica granatum* L.) grown in a calcareous soil. Middle East Journal of Applied Science. 2017;7(4):713-725.
14. Yadav A, Kumar N, Upadhyay A, Fawole OA, Mahawar MK, Jalgaonkar K, Chandran D, Rajalingam S, Zengin G, Kumar M, Mekhemar M. Recent advances in novel packaging technologies for shelf-life extension of guava fruits for retaining health benefits for longer duration. Plants. 2022;11(4):547.
15. Yadav VK, Jain MC, Sharma MK, Suman M. Effect of micronutrients spray on physical and chemical characteristics of pomegranate (*Punica granatum* L.) Cv. Sindhuri. International Journal of Current Microbiology and Applied Science. 2018;7:998-1005.
16. Sharma N, Kaushik L, Kumar R, Reddy DK. Pre-harvest application of calcium chloride and salicylic acid enhances morphological and biochemical characteristics of guava (*Psidium guajava*). Agricultural Mechanization in Asia. 2021;52(1):2545-2552.
17. Sajid M, Basit A, Ullah Z, Shah ST, Ullah I, Mohamed HI, Ullah I. Chitosan-based foliar application modulated the yield and biochemical attributes of peach (*Prunus persica* L.) cv. Early Grand. Bulletin of the National Research Centre. 2020;44(1):1-11.
18. Shanbehpour F, Rastegar S, Ghasemi M. Effect of pre-harvest application of calcium chloride, putrescine, and salicylic acid on antioxidant system and biochemical changes of two Indian jujube genotypes. Journal of Food Biochemistry. 2020; 44(11):13474.
19. Shashi Garhwal OP, Choudhary MR, Bairwa LN, Kumawat KL, Kumar P, Basile

- B, Corrado G, Roupael Y, Gora JS. 20. Singh P, Singh N, Singh SN. Studies on Effects of time of pruning and plant bio-regulators on the growth, yield, fruit quality, and post-harvest losses of ber (*Ziziphus mauritiana*). Horticulturae. 2022;8(9):809. improving storability and quality of harvested guava fruits using chemicals. Biological Forum – An International Journal. 2023;15(2):1124-1133.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/125178>