



# **Effect of Fertilizers and Humic Acids on Soil Micro Nutrients of Foxtail Millet Crop**

**Y. S. Satish Kumar <sup>a\*</sup>, Ch. Sujani Rao <sup>a</sup>, P. R. K. Prasad <sup>a</sup>,  
T.Venkata Sridhar <sup>a</sup>, S. Jaffar Basha <sup>a</sup> and K. Jayalalitha <sup>a</sup>**

<sup>a</sup> *Agricultural College Bapatla, ANGRAU, Andhra Pradesh, India.*

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: <https://doi.org/10.9734/arja/2024/v17i4605>

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/126914>

**Original Research Article**

**Received: 16/09/2024**  
**Accepted: 20/11/2024**  
**Published: 27/11/2024**

## **ABSTRACT**

Humic acids is an important soil component that can improve nutrient availability and impact on other important chemical, biological, and physical properties of soils. The ecological benefits of Humic acids are diverse and represent profitable and effective solutions for environmental problems and preservation of the environment. It can be integrated into the soils in the form of manure; it improves the physico chemical and biological properties of the soil. A field experiment was conducted at the College Farm, Agricultural College, Mahanandi, ANGRAU during kharif & rabi seasons of 2020-21 and 2021-22. The experimental soil was sandy loam in texture with a pH of 7.52, an Electrical Conductivity (EC) of 0.42 dS/m, 0.32% Organic Carbon (OC), low available nitrogen (175 kg ha<sup>-1</sup>), medium phosphorus (P) at 18.48 kg ha<sup>-1</sup>, high potassium (K) at 580 kg ha<sup>-1</sup>, and sufficient zinc (Zn) status at 0.85 ppm. The experiment was laid out in split plot design with

\*Corresponding author: E-mail: [yekkaladevikumar@gmail.com](mailto:yekkaladevikumar@gmail.com);

**Cite as:** Kumar, Y. S. Satish, Ch. Sujani Rao, P. R. K. Prasad, T.Venkata Sridhar, S. Jaffar Basha, and K. Jayalalitha. 2024. "Effect of Fertilizers and Humic Acids on Soil Micro Nutrients of Foxtail Millet Crop". *Asian Research Journal of Agriculture* 17 (4):940-48. <https://doi.org/10.9734/arja/2024/v17i4605>.

three replications with four main plots and six sub plots total twenty four treatments. After critical observation of the study, available iron, Zinc, Copper and Manganese contents in soil revealed that there was significant difference between the subplots that received different levels of Humic acids but main plots that received different levels of inorganic fertilizers and their interaction was found to be non significant at panicle initiation and harvest stages of foxtail millet crop in two *kharif* seasons of 2020 and 2021. Similar results were recorded in soil Copper and Manganese contents during two *kharif* seasons of Foxtail millet crop. However, the similar trend was followed in succeeding bengalgram crop.

**Keywords:** Foxtail millet; bengalgram cropping system; humic acids; in-organic fertilizers; soil micro nutrients content.

## 1. INTRODUCTION

“Sustenance of soil fertility is the key to crop productivity. Use of chemical fertilizers and organic manures has been found promising in arresting the decline trend in soil-health and productivity through the correction of marginal deficiencies of some secondary and micro-nutrients, micro-flora and fauna and their beneficial influence on physical and biological properties of soil. Integrated nutrient management system can bring equilibrium between degenerative and restorative activities in the soil eco-system” [1]. “Humic substances are generated through organic matter decomposition and employed as soil fertilizers in order to improve soil structure and soil microorganisms. Soil organic matter has been fractionated on the basis of solubility in dilute mineral acid and alkali in to three groups *viz.* fulvic acid, Humic acids and humin. Fulvic acids are soluble in both acid and alkali, Humic acids are soluble in alkali but insoluble in acids and humins are insoluble in both. Fulvic acids are relatively simple in composition and assimilable by plants, are labile in the soil. Humins are highly complex of the three forms and are unavailable to the plants. Humic acids occupy an intermediate position between these three groups and persist in the soil for a prolonged period so as to be useful to the crop plants” [2].

“Humic acids are organic compounds that play crucial roles in enhancing the qualities of soil, the growth of plants, and other agronomic factors. In recent years, products based on Humic acids have been incorporated into crop production to ensure the agricultural output's continued viability. According to the research that was conducted, HA has the potential to have a beneficial effect on the soil's physical, chemical, and biological properties. These properties include the aggregation and relative proportion of

soil particles, the capacity of soil to hold water, cation exchange capacity (CEC), pH, carbon content in the soil, enzymes activity, macronutrients cycling, and availability” [3]. “Humic acids (HA) producers claim that 1 kg of HA is as much beneficial as 1 tonne of cattle manure because manure needs a lot of time for humidification, the form that can be utilized and assimilated by plants” [4].

“Humic acids fraction contains about 60% organic carbon (C), which plays an important role in the growth of soil microorganisms. In addition to C, Humic acids also contain nitrogen (N), oxygen (O), hydrogen (H), and sulfur (S). Humic acids play several important roles such as increase soil physical and biochemical activities by improving structure, texture, water holding capacity (WHC), and microbial population increase soil nutrients availability, especially micronutrients by chelating and co-transporting micronutrients to plants reduce the transportation of toxic heavy metals by precipitating them, thus reducing toxic heavy metals intake by plants” (Nardi et al., 2017, 2021), [5].

## 2. MATERIALS AND METHODS

A field experiment was conducted at the college farm, Agricultural College, Mahanandi, Andhra Pradesh during *kharif* & *rabi* seasons of 2020-21 and 2021-22. The experimental site was geographically situated at 15.51° N latitude, 78.61° E longitude with an altitude of 233.48 meters above the mean sea level in Scarce Rainfall Zone of Andhra Pradesh. The experimental soil was sandy loam in texture with 7.52 pH, 0.42  $\text{dsm}^{-1}$  EC, 0.32 % OC, low available N (175 kg  $\text{ha}^{-1}$ ), medium in P (18.48 kg  $\text{ha}^{-1}$ ), high in K (580 kg  $\text{ha}^{-1}$ ) and sufficient in Zn status (0.85 ppm). The experiment was laid out in Split plot design with three replications with four main plots and six sub plots total twenty four

treatments viz., Control (M<sub>1</sub>), 50 % RDF (M<sub>2</sub>), 75 % RDF (M<sub>3</sub>) and 100% RDF (M<sub>4</sub>) as main plots and six Humic acids levels to foxtail millet crop comprising of No Humic acids application (S<sub>1</sub>), 10 kg ha<sup>-1</sup> Humic acids as soil application (S<sub>2</sub>), 20 kg Humic acids as soil application (S<sub>3</sub>), 0.2% of foliar application of Humic acids (S<sub>4</sub>), 10 kg ha<sup>-1</sup> Humic acids as soil application + 0.2% foliar application of Humic acids (S<sub>5</sub>) and 20 kg ha<sup>-1</sup> Humic acids as soil application + 0.2% foliar application of Humic acids (S<sub>6</sub>) as sub-plot treatments. These treatments were imposed to foxtail millet crop during kharif season and bengalgram crop during rabi season.

The 100% RDF for foxtail millet crop is 40:20:0 kg N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O ha<sup>-1</sup>. P fertiliser was applied as basal doses and half of the N was applied as basal and other half at 30 DAS. Humic acids was applied as basal as per treatments mentioned. Available nitrogen in soil was estimated by alkaline permanganate method using macro Kjeldahl distillation unit [6]. Available phosphorus in the soil was extracted with 0.5 M NaHCO<sub>3</sub> buffered at pH 8.5 and the phosphorus in the extract was estimated by ascorbic acid method using spectrophotometer at 660 nm [7]. Available Potassium was extracted with neutral normal ammonium acetate and estimated with the help of flame photometer [8]. Available zinc, iron, manganese and copper in the soils were determined in DTPA extract, using atomic absorption spectrophotometer [9].

### 3. RESULTS AND DISCUSSION

#### 3.1 Available Micro-Nutrients

##### 1. Iron

After critical observation of the study, available iron content in soil presented in Tables 1 & 2 revealed that there was significant difference between the subplots that received different levels of Humic acids but main plots that received different levels of inorganic fertilizers and their interaction was found to be non significant at panicle initiation and harvest stages of foxtail millet crop in two *kharif* seasons of 2020 and 2021. Though the main plots that receives different doses of fertilizers have not shown significant influence on soil iron content, the minimum soil iron content was recorded in control, where there is no fertilizers were applied (M<sub>1</sub>-6.24, 6.21mg kg<sup>-1</sup> in 2020 and 6.22, 6.19 mg kg<sup>-1</sup> in 2021) at panicle initiation and harvest stages, respectively). Whereas maximum

available iron content was recorded in the main plot that received 100% RDF (M<sub>4</sub>-6.40, 6.36mg kg<sup>-1</sup> in 2020 and 6.77, 6.72 mg kg<sup>-1</sup> in 2021). Increase in available Fe might be due to lowering of pH as a result of decomposition of organics which was known to increase the solubility of metallic elements [10].

Irrespective of the Humic acids treatments applications to foxtail millet crop, significantly the highest iron at panicle initiation and harvesting stages of crop was recorded in the treatment with 20 kg ha<sup>-1</sup> Humic acids as soil application + 0.2% foliar application (S<sub>6</sub>-6.51, 6.47mg kg<sup>-1</sup> in 2020 and 6.93, 6.88 mg kg<sup>-1</sup> in 2021). However it was on par with the treatment 20 kg ha<sup>-1</sup> Humic acids as soil application (S<sub>3</sub>). Significantly lower available iron content was recorded in control (S<sub>1</sub>-6.15, 6.12 in 2020 and 6.11, 6.08 mg kg<sup>-1</sup> in 2021) at panicle initiation and at harvest stages of *kharif* foxtail millet respectively. Humic acids has a good tendency to form chelates with metal ions and the formation of humate-metal ion complexes make them easily available to plants. The formation of humate-Fe complexes might have contributed to increased Fe availability in Humic acids applied plots. Similar results were also reported by Sharif et al. [11], Kumar & Singh [12], Manjeera [13] and Mankotia et al. [14].

##### 2. Zinc

Data pertaining to available zinc content presented in Tables 3 & 4 revealed that there was significant difference between the subplots that received different levels of Humic acids but main plots that received different levels of inorganic fertilizers and their interaction was found to be non significant at panicle initiation and harvest stages of foxtail millet crop in two *kharif* seasons of 2020 and 2021.

Among the different levels of RDF, the minimum available zinc content was recorded in control (M<sub>1</sub>-0.89, 0.88mg kg<sup>-1</sup> in 2020 and 0.86, 0.85 mg kg<sup>-1</sup> in 2021) at panicle initiation and harvest stages, respectively. Whereas maximum zinc content was recorded in the main plot that received 100% RDF (M<sub>4</sub>-1.02, 1.01mg kg<sup>-1</sup> in 2020 and 1.00, 1.00 mg kg<sup>-1</sup> in 2021) at panicle initiation and harvest stages, respectively. The available zinc decreased with advancement of crop stage during both the years of study period. The decrease might be attributed to uptake of zinc by the growing plants. Subhalakshmi and Pratapkumarreddy [15] also concluded that due to precipitation as in soluble sulfides and carbonates.

Table 1. Effect of inorganic fertilizers and Humic acids on available iron content (mg kg<sup>-1</sup>) in soil at panicle initiation stage of Foxtail millet

Sub Plots (Humic acids)	Kharif 2020				Mean	Kharif 2021				Mean
	Main Plots (In-Organics)					Main Plots (In-Organics)				
	M <sub>1</sub>	M <sub>2</sub>	M <sub>3</sub>	M <sub>4</sub>		M <sub>1</sub>	M <sub>2</sub>	M <sub>3</sub>	M <sub>4</sub>	
S <sub>1</sub>	6.12	6.15	6.16	6.18	<b>6.15</b>	6.08	6.11	6.12	6.14	<b>6.11</b>
S <sub>2</sub>	6.23	6.28	6.32	6.35	<b>6.30</b>	6.20	6.86	6.89	6.94	<b>6.72</b>
S <sub>3</sub>	6.31	6.42	6.55	6.61	<b>6.47</b>	6.32	6.98	7.13	7.19	<b>6.90</b>
S <sub>4</sub>	6.14	6.16	6.16	6.20	<b>6.17</b>	6.10	6.12	6.12	6.16	<b>6.12</b>
S <sub>5</sub>	6.25	6.31	6.37	6.39	<b>6.33</b>	6.26	6.89	6.96	6.98	<b>6.77</b>
S <sub>6</sub>	6.38	6.45	6.56	6.64	<b>6.51</b>	6.39	7.00	7.14	7.21	<b>6.93</b>
<b>Mean</b>	<b>6.24</b>	<b>6.30</b>	<b>6.35</b>	<b>6.40</b>		<b>6.22</b>	<b>6.66</b>	<b>6.73</b>	<b>6.77</b>	
	<b>SEm ±</b>	<b>CD (p=0.05)</b>		<b>CV (%)</b>		<b>SEm ±</b>	<b>CD (p=0.05)</b>		<b>CV (%)</b>	
<b>M</b>	0.01	NS		6.5		0.02	NS		7.5	
<b>S</b>	0.04	0.11		6.2		0.04	0.13		6.5	
<b>M X S</b>	0.02	NS				0.08	NS			
<b>S X M</b>	0.06	NS				0.04	NS			
<b>Main Plots</b>	<b>Sub Plots</b>									
M1-Control	S1-0kg/ha									
M2-50%RDF	S2-10kg/ha H.A									
M3-75%RDF	S3-20Kg/ha H.A									
M4-100%RDF	S4-0.2% H.A									
	S5-10kg/ha H.A+0.2%									
	S6-20kg/ha H.A+0.2%									

**Table 2. Effect of inorganic fertilizers and humic acid on available iron content (mg kg<sup>-1</sup>) in soil at harvesting stage of Foxtail millet**

Sub Plots (Humic acids)	Kharif 2020					Kharif 2021				
	Main Plots (In-Organics)				Mean	Main Plots (In-Organics)				Mean
	M <sub>1</sub>	M <sub>2</sub>	M <sub>3</sub>	M <sub>4</sub>		M <sub>1</sub>	M <sub>2</sub>	M <sub>3</sub>	M <sub>4</sub>	
<b>S<sub>1</sub></b>	6.08	6.12	6.13	6.14	<b>6.12</b>	6.05	6.08	6.09	6.11	<b>6.08</b>
<b>S<sub>2</sub></b>	6.21	6.25	6.28	6.32	<b>6.27</b>	6.17	6.83	6.86	6.90	<b>6.69</b>
<b>S<sub>3</sub></b>	6.28	6.39	6.52	6.58	<b>6.45</b>	6.28	6.92	7.06	7.13	<b>6.85</b>
<b>S<sub>4</sub></b>	6.10	6.13	6.13	6.16	<b>6.13</b>	6.07	6.09	6.09	6.13	<b>6.09</b>
<b>S<sub>5</sub></b>	6.23	6.28	6.34	6.36	<b>6.30</b>	6.22	6.86	6.92	6.94	<b>6.74</b>
<b>S<sub>6</sub></b>	6.35	6.41	6.53	6.60	<b>6.47</b>	6.35	6.94	7.07	7.14	<b>6.88</b>
<b>Mean</b>	<b>6.21</b>	<b>6.27</b>	<b>6.32</b>	<b>6.36</b>		<b>6.19</b>	<b>6.62</b>	<b>6.68</b>	<b>6.72</b>	
	<b>SEm ±</b>		<b>CD (p=0.05)</b>		<b>CV (%)</b>	<b>SEm ±</b>		<b>CD (p=0.05)</b>		<b>CV (%)</b>
<b>M</b>	0.02		NS		6.5	0.05		NS		6.0
<b>S</b>	0.05		0.15		6.8	0.04		0.13		6.8
<b>M X S</b>	0.06		NS			0.02		NS		
<b>S X M</b>	0.03		NS			0.01		NS		
<b>Main Plots</b>	<b>Sub Plots</b>									
M1-Control	S1-0kg/ha									
M2-50%RDF	S2-10kg/ha H.A									
M3-75%RDF	S3-20Kg/ha H.A									
M4-100%RDF	S4-0.2% H.A									
	S5-10kg/ha H.A+0.2%									
	S6-20kg/ha H.A+0.2%									

Table 3. Effect of inorganic fertilizers and Humic acids on available zinc content (mg kg<sup>-1</sup>) in soil at panicle initiation stage of Foxtail millet

Sub Plots (Humic acids)	Kharif 2020				Mean	Kharif 2021				Mean
	Main Plots (In-Organics)					Main Plots (In-Organics)				
	M <sub>1</sub>	M <sub>2</sub>	M <sub>3</sub>	M <sub>4</sub>		M <sub>1</sub>	M <sub>2</sub>	M <sub>3</sub>	M <sub>4</sub>	
S <sub>1</sub>	0.85	0.85	0.87	0.87	<b>0.86</b>	0.83	0.83	0.85	0.84	<b>0.84</b>
S <sub>2</sub>	0.86	0.89	0.94	1.05	<b>0.94</b>	0.85	0.88	0.93	1.04	<b>0.93</b>
S <sub>3</sub>	0.88	0.98	1.05	1.11	<b>1.01</b>	0.87	0.97	1.04	1.10	<b>1.00</b>
S <sub>4</sub>	0.87	0.87	0.88	0.88	<b>0.88</b>	0.83	0.83	0.84	0.84	<b>0.84</b>
S <sub>5</sub>	0.91	0.90	0.97	1.09	<b>0.97</b>	0.90	0.89	0.96	1.08	<b>0.96</b>
S <sub>6</sub>	0.94	0.98	1.08	1.12	<b>1.03</b>	0.90	0.97	1.07	1.11	<b>1.01</b>
<b>Mean</b>	<b>0.89</b>	<b>0.91</b>	<b>0.97</b>	<b>1.02</b>		<b>0.86</b>	<b>0.90</b>	<b>0.95</b>	<b>1.00</b>	
	<b>SEm ±</b>		<b>CD (p=0.05)</b>		<b>CV (%)</b>	<b>SEm ±</b>		<b>CD (p=0.05)</b>		<b>CV (%)</b>
<b>M</b>	0.01		NS		7.8	0.01		NS		6.5
<b>S</b>	0.01		0.03		6.4	0.01		0.04		6.2
<b>M X S</b>	0.02		NS			0.01		NS		
<b>S X M</b>	0.03		NS			0.02		NS		
<b>Main Plots</b>	<b>Sub Plots</b>									
M1-Control	S1-0kg/ha									
M2-50%RDF	S2-10kg/ha H.A									
M3-75%RDF	S3-20Kg/ha H.A									
M4-100%RDF	S4-0.2% H.A									
	S5-10kg/ha H.A+0.2%									
	S6-20kg/ha H.A+0.2%									

Table 4. Effect of inorganic fertilizers and Humic acids on available zinc content (mg kg<sup>-1</sup>) in soil at harvest of Foxtail millet crop.

Sub Plot (Humic acids)	Kharif 2020				Mean	Kharif 2021				Mean
	Main Plots (In-Organics)					Main Plots (In-Organics)				
	M <sub>1</sub>	M <sub>2</sub>	M <sub>3</sub>	M <sub>4</sub>		M <sub>1</sub>	M <sub>2</sub>	M <sub>3</sub>	M <sub>4</sub>	
<b>S<sub>1</sub></b>	0.84	0.85	0.86	0.87	<b>0.86</b>	0.82	0.83	0.84	0.83	<b>0.83</b>
<b>S<sub>2</sub></b>	0.85	0.88	0.93	1.04	<b>0.93</b>	0.85	0.88	0.93	1.04	<b>0.92</b>
<b>S<sub>3</sub></b>	0.87	0.97	1.04	1.10	<b>1.00</b>	0.87	0.97	1.04	1.10	<b>0.99</b>
<b>S<sub>4</sub></b>	0.86	0.87	0.88	0.88	<b>0.87</b>	0.83	0.83	0.84	0.84	<b>0.84</b>
<b>S<sub>5</sub></b>	0.90	0.89	0.96	1.08	<b>0.96</b>	0.90	0.89	0.96	1.08	<b>0.95</b>
<b>S<sub>6</sub></b>	0.93	0.97	1.07	1.11	<b>1.02</b>	0.89	0.97	1.07	1.11	<b>1.01</b>
<b>Mean</b>	<b>0.88</b>	<b>0.91</b>	<b>0.96</b>	<b>1.01</b>		<b>0.86</b>	<b>0.89</b>	<b>0.95</b>	<b>1.00</b>	
	<b>SEm ±</b>		<b>CD (p=0.05)</b>		<b>CV (%)</b>	<b>SEm ±</b>		<b>CD (p=0.05)</b>		<b>CV (%)</b>
<b>M</b>	0.03		NS		7.2	0.01		NS		6.8
<b>S</b>	0.02		0.05		6.1	0.02		0.05		6.4
<b>M X S</b>	0.01		NS			0.01		NS		
<b>S X M</b>	0.02		NS			0.02		NS		
<b>Main Plots</b>	<b>Sub Plots</b>									
M1-Control	S1-0kg/ha									
M2-50%RDF	S2-10kg/ha H.A									
M3-75%RDF	S3-20Kg/ha H.A									
M4-100%RDF	S4-0.2% H.A									
	S5-10kg/ha H.A+0.2%									
	S6-20kg/ha H.A+0.2%									

With increasing levels of Humic acids, significantly the highest zinc content was recorded with treatment S<sub>6</sub>(1.03, 1.01mg kg<sup>-1</sup> in 2020 and 1.01, 1.00mg kg<sup>-1</sup> in 2021) at panicle initiation and at harvest stages, respectively over control. This treatment was on par with the treatment which receives 20 kg ha<sup>-1</sup>Humic acids as soil application (S<sub>3</sub>). Lower zinc content was recorded in control (S<sub>1</sub>-0.86, 0.86 mg kg<sup>-1</sup> in 2020 and 0.84, 0.83 mg kg<sup>-1</sup> in 2021) at panicle initiation and at harvest stages, respectively. Humic acids has a good tendency to form chelates with metal ions and the formation of humate-metal ion complexes make them easily available to plants. Increased zinc availability might be attributed due to prevention of formation of immobile and insoluble hydroxides of zinc by Humic acids. Similar results were also reported by Singhal et al. [16], Kumar et al. [17] and Ameta et al., [18] & Kumar & Singh [12], De Ávila et al. [19], Keeling et al. [20], Mikkelsen [21] and Stevenson [22].

Similar results were recorded in soil Copper and Manganese contents during two kharif seasons of Foxtail millet crop.

#### 4. CONCLUSIONS

After critical observation of the study, available iron, Zinc, Copper and Manganese contents in soil revealed that there was significant difference between the subplots that received different levels of Humic acids but main plots that received different levels of inorganic fertilizers and their interaction was found to be non significant at panicle initiation and harvest stages of foxtail millet crop in two *kharif* seasons of 2020 and 2021. Similar results were recorded in soil Copper and Manganese contents during two kharif seasons of Foxtail millet crop.

#### FUTURE LINE OF WORK

Future research may be carried out to consider the potential effects of imposed treatments on Iron, Zinc fractions in soil, to test the treatments in other important cropping systems viz., rice-millet, cereal-pulse etc.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

#### ACKNOWLEDGEMENT

The authors are extremely grateful to Acharya N G Ranga Agricultural University, Guntur, Andhra Pradesh for generous assistance for the said project.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Upadhyay VB, Jain V, Vishwakarma SK, Kumhar AK. Production potential, soil health, water productivity and economics of rice (*Oryza sativa*)-based cropping systems under different nutrient sources. Indian Journal of Agronomy. 2011;56(4): 311-16.
2. Ravichandran M. Humic acids: A mystique substance in sustainable crop production. Journal of the Indian Society of Soil Science. 2011;59:49-57.
3. Pooja Bhatt, Singh VK. Effect of Humic acids on soil properties and crop production- A review. Indian Journal of Agricultural Sciences. 2022;92(12):1423-1430.
4. Satish Kumar YS, Sujani Rao Ch, Prasad PRK, Jayalalitha K, Jaffar Basha S, Sridhar T Venkata. Effect of inorganic fertilizers and humic acids on soil nutrients of foxtail millet crop under foxtail millet-bengal gram cropping system. The Andhra Agricultural Journal. 2021;68(4):481-488.
5. Among K, Thilakarathna MS and Gorim LY. Understanding the role of humic acids on crop performance and soil health. frontiers in agronomy. 2022;4:848621. DOI: 10.3389/fagro.2022.848621.
6. Subbiah BV, Asija CL. A rapid procedure for the estimation of available nitrogen in soils. Current Science. 1956;25: 259-260.
7. Watanabe FS, Olsen SR. Test of ascorbic acid method for determining phosphorous in water and sodium bicarbonate extracts of soil. Soil Science Society of American Journal. 1965;29:677-78.
8. Jackson ML. Soil Chemical Analysis. Prentice Hall of Inco. New York, USA. 1973;498.
9. Lindsay WL, Norvell WA. Developments of DTPA soil test for zinc, iron, manganese



- and copper. Soil Science Society of American Journal. 1978;42:421-428.
10. Prasad RK, Kumar V, Prasad B, Singh AP. Long-term effect of crop residues and zinc fertilizer on crop yield, nutrient uptake and fertility buildup under rice-wheat cropping system in Calciorthents. Journal of Indian Society of Soil Science. 2010b;58(2):205-211.
  11. Sharif M, Khattak RA, Sarir MS. Effect of different levels of lignitic coal derived Humic acids on growth of maize plants. Communications in Soil Science and Plant Analysis. 2002;33:3567– 3580.
  12. Kumar, Singh. Efficacy of Potassium Humate and Chemical Fertilizers on Yield and Nutrient Availability Patterns in Soil at Different Growth Stages of Rice, Communications in Soil Science and Plant Analysis. 2017;48(3):245- 261.
  13. Manjeera K 2020.Effect of Humic acids and inorganic nitrogen on soil properties and yield of direct sown rice. Msc.Thesis. Submitted to Acharya N.G. Ranga Agricultural University, Guntur, Andhra Pradesh.
  14. Mankotia S. Impact of Humic acids on various properties of soil and crop productivity- A review. Journal of Agriculture and Ecology. 2024;18:1-6. Available:<https://doi.org/10.58628/JAE-2418-101>
  15. Subhalakshmi C, Pratap Kumar Reddy A. Soil available nutrient status as influenced by organic sources and fertilizer levels in hybrid rice. International Journal of Science and Nature. 2017;8(1):40-43.
  16. Singhal VK, Shar K, Sharma, Teekam Singh. Integrated effect of enriched compost and fertilizer on yield and uptake of micronutrient by maize. Agriculture Science Digest. 2012;32(1):43–47.
  17. Kumar M, Yaduvanshi NPS, Singh YV. Effect of integrated nutrient management on soil fertility status in reclaimed sodic soils. Journal of the Indian Society of Soil Science. 2012;60(2):132-137.
  18. Ameta KD, Sharma SK, Dubey RB, Kaushik RA. Effect of Humic acids and Micro Nutrients on Growth and Yield of Poly House Grown Cucumber (*Cucumis sativus* L.). Chemical Science Review and Letters. 2017;6 (21):581-584.
  19. de Ávila MO, Moreira SG, Lima FR, Pimentel GV, Macedo JR, Nunes MR, Gomes LB, Morais EG. Effect of coating phosphorus with humic acids and micronutrients on yield of soybean and maize in succession. Journal of Agriculture and Food Research. 2024 Dec 1;18: 101318. Available:<https://doi.org/10.1016/j.jafr.2024.101318>.
  20. Keeling AA, McCallum KR, Beckwith CP. Mature green waste compost enhances growth and nitrogen uptake in wheat (*Triticum aestivum* L.) and oilseed rape (*Brassica napus* L.) through the action of water-extractable factors. Bioresource Technology. 2003 90(2): 127-137.
  21. Mikkelsen RL. Humic materials for agriculture, Davis, California, USA. Better Crops with Plant Food. 2005;89(3):6-7.
  22. Stevenson FJ. Humus chemistry: Genesis, composition, reaction. 2nd Ed. John Wiley and sons, New York. 1994;26-54.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/126914>