



International Journal of Environment and Climate Change

Volume 14, Issue 12, Page 82-90, 2024; Article no.IJECC.125786

ISSN: 2581-8627

(Past name: British Journal of Environment & Climate Change, Past ISSN: 2231-4784)

Effects of Climate Change and Greenhouse Gas Emission Relevance to Environmental Stress on Horticultural Crops

Hariom Katiyar ^{a++}, S.P. Singh ^{b##}, Rohit Maurya ^{ct},
Ajit Kumar Srivastava ^{d‡}, Rahul Kumar Verma ^{e†}
and Vijay Kumar Yadav ^{f^}

^a Department of Fruit Science, College of Horticulture, SVPUAT, Meerut-250110, India.

^b Department of Horticulture, KVK Belipar, Gorakhpur, India.

^c Department of Vegetable Science, KVK Turki, Muzaffarnagar, DRPCAU, Pusa Samstipur, Bihar, India.

^d Department of Horticulture, Mahayogi Gorakhnath Krishi Vigyan Kendra, Chauk Mafi, Pipiganj, Gorakhpur 273165, India.

^e Department of Horticulture, KVK, Madhepura -852113, BAU Sabour, India.

^f Department of Horticulture, N.N. PG College Nawabganj, Gonda, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijecc/2024/v14i124608>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/125786>

Review Article

Received: 24/09/2024

Accepted: 28/11/2024

Published: 03/12/2024

⁺⁺ Associate Professor;

[#] SMS/ Associate Professor;

[†] SMS Horticulture;

[‡] Subject Matter Specialist;

[^] Student;

*Corresponding author: E-mail: spskvkgkp73@gmail.com;

Cite as: Katiyar, Hariom, S.P. Singh, Rohit Maurya, Ajit Kumar Srivastava, Rahul Kumar Verma, and Vijay Kumar Yadav. 2024. "Effects of Climate Change and Greenhouse Gas Emission Relevance to Environmental Stress on Horticultural Crops". *International Journal of Environment and Climate Change* 14 (12):82-90. <https://doi.org/10.9734/ijecc/2024/v14i124608>.

ABSTRACT

Global climate change significantly impacts the yield and quality of sustainable horticulture crops. Rising global temperatures have forced the agricultural community to modify planting and harvesting timetables, frequently calling for earlier crop production. Notably, several concerning elements are brought about by climate change, including greenhouse gas emissions (GHGs), higher temperatures, higher concentrations of carbon dioxide (CO₂), nitrous oxide (N₂O) and methane (CH₄), ozone depletion (O₃), and deforestation. All of these factors exacerbate environmental stresses on crops. Climate change is, therefore, likely to have a negative impact on livestock production and crop yields. Thus, the main goal of the review paper is to provide a thorough summary of the various aspects that affect the production of fruits, vegetables, and plantation crops in horticulture, with a focus on greenhouse gas emissions and environmental stressors like high temperatures, droughts, salinity, and CO₂ emissions. This assessment will also look at how new greenhouse technologies and horticultural crop varieties can be used to lessen the adverse effects of climate change on crops.

Keywords: Carbon trade; carbon sink; climate change; greenhouse; horticultural crops.

1. INTRODUCTION

Climate change is defined as variations in the global temperature over a comparable length of time that are either directly or indirectly caused by human activities changing the composition of the atmosphere. Over the past century, the average global temperature of the earth's surface has risen by roughly 0.74°C. Regarding the worldwide and regional effects of anticipated climate change on agriculture, water resources, natural ecosystems, and food nutritional security, the Fourth Intergovernmental Panel on Climate Change (IPCC) report (IPCC 2007) was appropriately envisioned. Abiotic pressures like drought, hailstorms, heavy rain, floods, frost, cyclones, and other natural disasters affect different provinces and areas every year, and these events are attributed to the effects of climate change. A shifting climate due to altered weather patterns has put agricultural productivity at risk of high and low-temperature regimes and increased rainfall variability (Malhotra and Srivastava 2014; Eduardo et al. 2013).

“One of the main issues impacting the performance of agriculture, particularly horticultural crops, both annual and perennial, is climate change and its unpredictability. The short growing season, which will have a detrimental effect on growth and development, mainly because of terminal heat stress and lower water availability, is probably the cause of the reduction in fruit and vegetable yield. Rainfall variability and a decline in rainy days would significantly affect rainfed agriculture” (Venkateswarlu and Shanker 2012). “Horticultural production systems are now subject to additional constraints because

of the uncertainties and dangers associated with climate change and variability. Fruit and vegetable crop prices may rise as a result of climate change” (Hasan et al., 2024; Ramadhan et al., 2023). “The challenges ahead are sustainability and competitiveness, as well as achieving the targeted production to meet the increasing demands in the face of diminishing land and water resources and the threat of climate change. Climate-smart horticulture interventions are necessary to improve output in these challenging environments, and they require a high degree of location specificity and knowledge” (Malhotra and Srivastava 2014, Malhotra 2015).

“The greenhouse effect occurs in the troposphere, or lower atmosphere layer, where weather and life exist. The Earth's average surface temperature is predicted to be approximately -19°C in the absence of the greenhouse effect, as opposed to the current average of 14°C” (Le Treut et al., 2007). Greenhouse gasses (GHG) are the cause of the greenhouse effect. According to the IPCC (2014), greenhouse gases (GHGs) are the gaseous components of the atmosphere that both emit and absorb thermal infrared radiation. The troposphere contains traces of both naturally occurring and artificial greenhouse gases. In ascending priority order, the most prevalent greenhouse gases are water vapour, carbon dioxide (CO₂), methane (CH₄), nitrous oxides (N₂O) and ozone (O₃). GHG percentages vary daily, seasonally, and annually. The primary organic trace gas in the atmosphere is methane (CH₄). The main component of natural gas, a fuel used all over the world, is CH₄. Agriculture

and cattle raising produce significant amounts of greenhouse gas emissions, primarily due to the use of fossil fuels. Since the pre-industrial era, CH₄ concentrations have increased by a factor of two. As of 2014, the average concentration globally is 1.8 $\mu\text{mol}\cdot\text{mol}^{-1}$. Despite the fact that its concentration is just 0.5% of CO₂, worries are raised about an increase in the atmospheric release of CH₄. As a greenhouse gas, it is, in fact, thirty times more potent than CO₂ (IPCC, 2014). In addition to producing O₃, CH₄ and CO₂ help regulate the amount of OH in the troposphere (Wuebbles and Hayhoe, 2002).

1.1 Impact of Greenhouse Gases on Horticultural Crops

“Effects of increased CO₂ Twenty percent of the heat absorption is caused by carbon dioxide (Schmidt et al., 2010). Respiration, seawater release, and organic decomposition are examples of natural sources of CO₂. Anthropogenic sources of CO₂ emissions include cement production, deforestation, and burning fossil fuels, including coal, oil, and natural gas. Industry accounts for 21% of direct CO₂ emissions, and agriculture, forestry, and other land use accounts for 24%” (IPCC, 2014). “Over the last 200 years, atmospheric CO₂ concentrations have increased significantly, from approximately 270 $\mu\text{mol}\cdot\text{mol}^{-1}$ in 1750 to current levels of over 385 $\mu\text{mol}\cdot\text{mol}^{-1}$ ” (Mittler and Blum Wald, 2010; IPCC, 2014). Since the 1970s, around half of all anthropogenic CO₂ emissions between 1750 and 2010 have occurred (IPCC, 2014). According to calculations, the global mean surface temperature will rise by 3-5°C in 2100 due to rising CO₂ concentrations and positive feedback from water (IPCC, 2014).

“The impact of new cultivars and production system management on technological advancements is evident in higher productivity and production, which increased by more than eleven times to 283.2 million tonnes in 2015–16 (3rd estimates) from 25 million tonnes in 1950–51” (Malhotra 2016). The horticulture industry has undoubtedly advanced despite numerous obstacles and failings, and it is currently at a pivotal stage of growth that calls for initiatives for sustainable development. Vertical growth, through the use of new cultivars, efficient water and nutrient management, and effective plant health management coupled with strategies for reduced post-harvest losses, could be the approach, which would require appropriate innovation and investment, to achieve the

targeted production of 310 million tonnes of horticultural crops by the end of the XII Plan (2012–17). The challenges and effects of climate change, such as altered seasonal patterns, excessive rain, flooding, hailstorms, frost, high temperatures, and drought that cause extremes, must be addressed to achieve increased horticultural production. Reduced water availability, shorter growing seasons, and inadequate vernalization can all result in lower yields.

We require comprehensive data on the physiological reactions of the crops, as well as effects on growth and development, quality, and production, to effectively evaluate the effects of climate change on horticulture crops. To prepare for the impending difficulties of climate change, the horticulture industry must handle the diverse implications in a coordinated and organized manner. Increased respiration, altered photosynthesis, and the partitioning of photosynthates into economically significant portions are all caused by temperature increases. Moreover, it can change the phenology, accelerate fruit maturity, ripening, and senescence, and decrease the crop's growing season and number of blooming and fruiting days. A crop's sensitivity to temperature varies depending on its growing patterns and innate tolerance. Because they flower later than determinate crops, indeterminate crops are less susceptible to heat stress. Seasons and day and night variations in the rise in temperature may not be equally distributed (Srinivas Rao et al. 2010). In tropical areas, yield declines may be excessive, even with mild warming. A slight rise in temperature can lead to an improvement in agricultural yields in high-latitude regions. Since most emerging nations are found in lower latitudes, temperatures are already near or beyond thresholds, and more warming would decrease rather than boost production. The effects of climate change are likely to vary depending on the crop type and locality; these effects are discussed here for many horticultural crop subsectors.

1.2 Impact of NO₂

“Over the past few decades, there has been an increase in the emission of atmospheric nitrogen dioxide (NO₂) in various parts of the world (Munzi et al., 2009), particularly in some Asian nations” (Hu et al., 2015). “The NO₂ concentration is expected to continue rising and above the standard limit of NO₂ pollution due to

the expansion of industrialized production and the ongoing rise in vehicle exhaust emissions" (Hultengren et al., 2004). "Ozone and particle matter are two dangerous secondary air pollutants that are preceded by nitrogen dioxide (NO₂)" (Rahmat et al., 2013, Brmejo-Orduna et al., 2014). "It is crucial to employ ecological techniques, such as plant uptake and atmospheric NO₂ catabolism, to lower air concentrations. Scholars now disagree on many aspects of NO₂'s effects on plants. One theory states that NO₂ can generate organic nitrogenous molecules in plants by metabolization and incorporation into the nitrate absorption pathway. According to a different theory, most plants resist NO₂ in addition to low levels of NO₂-N incorporation into total plant nitrogen (N)" (Morikawa et al., 2001). "When plants are exposed to NO₂, complex physiological reactions can occur, such as modifications in the activity of antioxidant enzymes" (Liu et al., 2015). N metabolic enzyme activity and the makeup and arrangement of nitrogenous metabolic products in plant tissues (Vighi et al., 2017). Low amounts of NO₂ can dissolve in water to generate nitrate and nitrite, which are then used by plants in the course of their regular nitrate metabolism. In this way, NO₂ can function as an airborne fertilizer. On the other hand, distinct plant species react physiologically differently when exposed to NO₂. As a result, there is ongoing debate on the effects of NO₂ exposure on plants, and no consensus has been achieved. Furthermore, little is known about the various plants that naturally recover from high NO₂ concentrations. We investigated the physiological reactions of 41 garden plants from Jiangsu Province subjected to various NO₂ environments under controlled laboratory circumstances to find plants that show robust resistance and sound absorption. Few research has looked into whether these 41 plant species, frequently planted by roadsides in many nations' cities, have a high capacity to absorb NO₂ or are resistant to its effects. Numerous research works have examined how NO₂ concentration affects plant growth and have found that while low NO₂ concentrations (0.1 µl/L) did not significantly affect plant height, The study found that 0.5µl/L NO₂ considerably accelerated the leaf growth of *Populus Deltoides* and *Populus Nigra Italic a*. Still, a greater NO₂ concentration (1 µl/L) significantly inhibited the stem growth. The seedlings were 1-year-old *Buxus Sinica* seedlings. After being subjected to 0.85, 2, 4.25, and 9.4µl/L NO₂, *Arabidopsis thaliana* plants showed acute, evident leaf

damage before dying. Even though those plants suffered considerable damage when exposed to NO₂ concentrations ranging from 2 to 4.25 µl/L, there were no appreciable variations in leaf growth or chlorophyll (Chl) content. As a result, we determined that the NO₂ fumigation concentration should be six µl/L (Sheng et al., 2019).

1.3 Impact of SO₂

"Stress circumstances, especially air pollution, modify plant morphological, physiological, and biochemical processes by preventing net photosynthesis, slowing down development, and producing immediate observable damage" (Ashraf & Harris, 2013). "One of the first biological processes impacted by elevated SO₂ concentrations is photosynthesis; plants in these conditions show an initial reduction in photosynthetic rate and an increase in respiration rate" (Gheorghe and Ion, 2011). According to Verma and Singh (2006), the effects of SO₂ on physiological processes are linked to stomatal responses. This is probably because stressed plants are unable to seal their stomata appropriately. Moreover, SO₂ impacts chlorophyll levels and carbon allocation, both of which impact plant development and productivity and modify the pace of photosynthetic reaction and the distribution of photosynthates, which impacts plant growth.

Additionally, it modifies membrane permeability and alters the structure of the cell membrane. The carbon gain to water loss ratio is a plant's water usage efficiency or WUE. According to Swanepoel et al. (2007), prolonged exposure to SO₂, even at moderate concentrations, lowered WUE and chlorophyll content.

On the other hand, WUE was elevated by high concentrations and brief SO₂ exposure. Environmental stress is known to have an impact on chlorophyll concentration). There is consensus that a drop in chlorophyll concentration can signal SO₂ damage (Haworth et al., 2012). A plant's degree of harm from absorbing SO₂ is categorized as acute or chronic. While long-term absorption of SO₂ at threshold concentrations results in chronic harm, acute injury is produced by high concentrations of SO₂ absorbed within a brief period.

Nevertheless, it interferes with the electron transport chain and impairs thylakoid activity at high concentrations, which results in the collapse

of physiological systems. Some have proposed that the pace at which SO₂ is absorbed determines the extent of SO₂ harm. Through the roots in the soil or the leaves in the atmosphere, SO₂ can enter the plant tissues. Plant leaves absorb atmospheric SO₂ through stomata, which guard cells can open or close. Increased SO₂ concentrations typically result in stomatal alterations; prolonged high SO₂ concentrations tend to induce stomata to close. Guard cells absorb sulfur after exposure to SO₂ and become incapable of opening or closing their stomata. Consequently, SO₂ modifies the synthesis and distribution of photosynthates, which impacts plant growth. The reproductive process in plants can be hampered by the suppression of photosynthesis caused by high sulfur levels, perhaps resulting in fewer flowers or fruits (Hetherington and Woodward, 2003).

1.4 CH₄ Emission

Methanogenesis is the process by which bacteria create methane. The creation mechanisms for methane, which is found in rocks and soil, differ from those of other fossil fuels. Methane is derived from ancient biomass. Methane emissions originate from both natural and manmade sources. The oceans, termites, and wetlands are the primary natural sources. Three-quarters of methane emissions come from natural sources. Livestock farms and landfills are examples of human sources.

Nonetheless, fossil fuel production, consumption, and transportation constitute the primary source. Sixty-four percent of methane emissions originate from sources related to humans. Paddy fields account for 91% of methane emissions, with animal husbandry and the burning of agricultural wastes contributing less than 2% and 7%, respectively. When coal, natural gas, and oil are produced and transported, methane is released into the atmosphere. Methane emissions can also be attributed to the decomposition of organic waste in municipal solid waste dumps, livestock, and other agricultural processes. Higher frequency of physiological conditions such as blossom end rot and tip burn. Early maturity of citrus, grapes, melons, and onions is a result of rising temperatures. The citrus fruit set is low because of the warm weather. Many spring-flowering plants, like mangos, began to bud up to 4 days early for every one °C increase in the average spring temperature following two recent warm periods. The hilly mountain region of Himachal

Pradesh, where apples and other temperate crops are traditionally grown, was suitable for cultivation because it consistently experiences the ideal temperature for fruiting and flowering. However, as global warming continues, this temperature may eventually rise to a point where it is no longer suitable.

2. UTILIZING LOCATION-SPECIFIC, CLIMATE-SMART HORTICULTURE TO COMBAT CLIMATE CHANGE

Climate-smart horticulture is not a single, all-encompassing agricultural technique or technology. According to Malhotra (2014), this method necessitates site-specific assessments to find appropriate production technologies and techniques that may address the various issues that agriculture and food systems face concurrently and comprehensively. Although climate change's form, breadth, and scale vary among different countries and sites, it is a worldwide phenomenon. Therefore, local analysis, planning, and administration are needed to address the concerns of climate change and the difficulties that result from it. In order to effectively address the issues, it is necessary to assess and comprehend how climate change is affecting both annual and perennial horticultural crops at the regional level. This can be done by managing innovation and evaluating and improving technology (Malhotra and Srivastava 2014).

3. SIMULATION MODELS FOR IMPACT ASSESSMENT

"Using modelling methods for impact analysis for different horticulture crops will be helpful when developing adaptation and mitigation strategies. In India, there aren't many good simulation models available for horticulture crops (fruit and vegetables), possibly with the exception of coconut and potatoes. The Info Crop model has been modified for tomato and onion crops, and its validation for various agroecological zones is ongoing" (Naresh Kumar et al., 2008). Large fruit trees and bushes pose challenges in studies examining the direct effects of many factors on growth, development, and yield in controlled environments due to their perennial nature. Prioritizing the development of simulation models for significant horticulture crops such as mango, grape, apple, orange, citrus, litchi, guava, etc. calls for creative approaches. Developing crop simulation models for horticultural crops in India is now a priority area of research.

4. PRODUCTION SYSTEM MANAGEMENT

“The focus should be on using the suggested production technologies to increase water efficiency and adjust to the dry, hot weather. It is necessary to employ tactics like shifting the dates of sowing or planting in order to counteract the anticipated rise in temperature and periods of water stress during the crop-growing season. Utilizing soil amendments to increase soil fertility, improve nutrient uptake, and adjust fertilizer application to improve nutrient availability” (Srivastava et al. 2014, Malhotra and Srivastava 2015). “The two most crucial interventions are conserving soil moisture reserves and providing irrigation throughout crucial phases of crop growth” (Malhotra 2016). Soil moisture conservation is aided by crop management techniques such as mulching with crop waste and artificial mulches. Raised beds can be used to cultivate crops when heavy rains cause excessive soil moisture, which can occasionally become a severe issue. Vegetable production could be pursued with clear plastic rain shelters, which can lessen the direct effect on developing fruits and field water logging during the rainy season. Vegetables grown on raised beds during the rainy season will produce more because of better drainage and less anoxic stress on the root system. Vegetable scion cultivars would become resistant to soil-related environmental challenges such as drought, salinity, low soil temperature, and flooding if grafted onto tolerant rootstocks (Chieri et al. 2008).

5. CARBON SEQUESTRATION POTENTIAL

As a potential source of extra income for rural communities that might otherwise struggle financially and as a way to support climate change adaptation, mitigation measures in the agriculture and forestry sectors are attracting much attention. Carbon sequestration is one method of mitigation that helps lessen the adverse effects of climate change. Despite their considerable contribution, fruit trees' capacity to sequester carbon is little known. The mitigation potential of farm forestry fruit orchard block planting was assessed in a study conducted at IISC, Bangalore, using the PRO-COMAP model. *Mangifera indica*, *Tamarindus indica*, *Achras sapota*, *Artocarpus*, *Neem*, and *Guava* accounted for 75% of the projected area. For the 30 years (2005–2035), the carbon stock change under the baseline and mitigation scenario (apart from harvested wood products) and the carbon

increment per ha for different project activities came out to be 47.42 t C/ha. With a 5,381 ha area, agricultural forestry has an overall mitigation potential of 81,750 t C. The highest profitability is seen in fruit orchard farm forestry, with an IRR of 29.92% (Ravindranath et al., 2007; Laxman et al., 2010).

6. TECHNOLOGICAL CHANGE FOR MITIGATING EFFECT

Actually, grapes are a temperate fruit that is often grown in chilly climates, whether they are being grown for wine or table use. Nonetheless, technological advancements in plant architecture and production system management have made growing grapes in tropical climates with the highest global output possible. Similarly, high temperatures in the mid-hill agroclimatic conditions may produce pollen desiccation and fruit shrivelling, diminishing yield and increasing crop failure. Chilling alone will also not be sufficient to trigger flowering in apples. These are the anticipated effects that give rise to the worries. However, countless examples show how technologies have assisted in minimizing the effects of climate change. Alkalinity and salinity posed significant challenges to the practical cultivation of grapes, but the discovery of appropriate rootstocks has increased yield significantly. Potatoes, tomatoes, cauliflower, and cabbage are thermosensitive crops that were only productive during lengthy days in a temperate climate. However, it is now feasible to achieve extremely high productivity even in subtropical and mild subtropical regions with greater temperatures because of the introduction of heat-tolerant cultivars and modifications in production system management (Singh et al. 2008, Malhotra and Srivastava 2014). These historical examples eloquently illustrate that, with creative research, the threat posed by climate change could be turned into an opportunity. However, this will require depicting the change's expected course and effects and plans to lessen its adverse effects. Today's biotechnology techniques could contribute to a quicker delivery of study findings.

7. CONCLUSION

Numerous expert studies show how different horticultural crops may be affected by climate change. The measurement of the effects of differences in Numerous experts' studies shows how different horticultural crops may be affected by climate change. The first stage in preparing

the horticulture industry to develop adaptation strategies under climate change scenarios is quantifying the effects of temperature changes and surplus and limited moisture conditions. Coordinated efforts are required to investigate and evaluate the effects on specific crops under the main agro-ecological zones and growing seasons. Increased efforts should be made to create new cultivars that are suited for various agroecological zones with shifting climates. For annual crops, the adaptation tactics can be implemented quickly by utilizing a variety of cultivars and species, altering planting dates, or changing the season. However, when establishing and rearranging orchards, the longer-term effects of climate change must be considered. Therefore, a thorough analysis of how climate change is affecting perennial crops is required before pursuing any adaptation options.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Ashraf, M., & Harris, P. J. C. (2013). Photosynthesis under stressful environments: An overview. *Photosynthetic Research*, 51, 163-190.
- Bermejo-Orduna, R., McBride, J. R., Shiraishi, K., Elustondo, D., Lasheras, E., & Santamaría, J. M. (2014). Biomonitoring of traffic-related nitrogen pollution using *Letharia vulpina* (L.) Hue in the Sierra Nevada, California. *Science of the Total Environment*, 490, 205-212.
- Edenhofer, O. R., Pichs-Madruga, Y., Sokona, E., Farahani, S., Kadner, K., Seyboth, A., ... & IPCC. (Eds.). (2014). *Climate change 2014: Mitigation of climate change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press.
- Gheorghe, I. F., & Ion, B. (2011). The effects of air pollutants on vegetation and the role of vegetation in reducing atmospheric pollution. In *The Impact of Air Pollution on Health, Economy, Environment and Agricultural Sources* (pp. 256-259). Intech Publisher.
- Hasan, M. M., Nobil, F. B., Khan, N. A., Toma, N. I., Islam, M. W., Ali, M. A., & Akram, S. (2024). Greenhouse gas emissions and energy consumption in the garments, textile, and dyeing sectors in Savar, Dhaka (2019-2021). *Asian Journal of Environment & Ecology*, 23(10), 88-95.
- Haworth, M., Elliott-Kingston, C., Gallagher, A., Fitzgerald, A., & McElwain, J. C. (2012). Sulphur dioxide fumigation effects on stomatal density and index of non-resistant plants: Implications for the stomatal palaeo-[CO₂] proxy method. *Review of Palaeobotany and Palynology*, 182, 44-54.
- Hetherington, A. M., & Woodward, F. I. (2003). The role of stomata in sensing and driving environmental change. *Nature*, 424, 901-908.
- Hu, Y. B., Bellaloui, N., Tigabu, M., Wang, J. H., Diao, J., Wang, K., & et al. (2015). Gaseous NO₂ affects stomatal behavior, photosynthesis, and respiration of hybrid poplar leaves. *Acta Physiologiae Plantarum*, 37, 39.
- Hultengren, S., Gralen, H., & Pleijel, H. (2004). Recovery of the epiphytic lichen flora following air quality improvement in south-west Sweden. *Water, Air, & Soil Pollution*, 154, 203-211.
- Intergovernmental Panel on Climate Change (IPCC). (2007). *Climate change 2007: Fourth assessment report of the Intergovernmental Panel on Climate Change (IPCC)*. World Meteorological Organization (WMO), United Nations Environment Programme (UNEP).
- Kubota, C., McClure, M. A., Kokalis-Burelle, N., Bausher, M. G., & Roskopf, E. N. (2008). Vegetable grafting: History, use, and current technology status in North America. *Horticultural Science*, 43(6), 1664-1669.
- Kumar, N., Kasturi Bai, K. V., Rajagopal, V., & Aggarwal, P. K. (2008). Simulating coconut growth, development, and yield using the INFOCROP-Coconut model. *Tree Physiology*, 28, 1049-1058.
- Laxman, R. H., Shivasham, B. K. S., & Srinivasa Rao, N. K. (2010). An assessment of potential impacts of climate change on fruit crops. In H. P. Singh, J. P. Singh, & S. S. Lal (Eds.), *Challenges of climate change in*

- Indian horticulture* (pp. 23-30). Westville Publishing House.
- Le Treut, H., Somerville, R., Cubasch, U., Ding, Y., Mauritzen, C., Mokssit, A., ... & Solomon, S. (Eds.). (2007). Historical overview of climate change science. In *Climate change 2007: The physical science basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* (pp. 1-26). Cambridge University Press.
- Liu, X. F., Hou, F., Li, G. K., & Sang, N. (2015). Effects of nitrogen dioxide and its acid mist on reactive oxygen species production and antioxidant enzyme activity in *Arabidopsis* plants. *Journal of Environmental Sciences*, *34*, 93-99.
- Malhotra, S. K. (2014). Development strategies for climate-smart horticulture. In *Global conference held at Navsari on 28-30 May 2014*.
- Malhotra, S. K. (2015). Mission approach for the development of protected cultivation. *New Age Protected Cultivation*, *1*(1), 29-32.
- Malhotra, S. K. (2016). Water soluble fertilizers in horticultural crops – An appraisal. *Indian Journal of Agricultural Sciences*, *86*(10), 1245-1256.
- Malhotra, S. K. (2016). Water soluble fertilizers in horticultural crops – An appraisal. *Indian Journal of Agricultural Sciences*, *86*(10), 1245-1256.
- Malhotra, S. K., & Srivastva, A. K. (2014). Climate-smart horticulture addresses food, nutritional security, and climate challenges. In A. K. Srivastava et al. (Eds.), *Shodh Chintan- Scientific articles* (pp. 83-97). ASM Foundation.
- Mittler, R., & Blumwald, E. (2010). Genetic engineering for modern agriculture: Challenges and perspectives. *Annual Review of Plant Biology*, *61*, 443-462. <https://doi.org/10.1146/annurev-arplant-042809-112116>
- Morikawa, H., Higaki, A., Nohno, M., Takahashi, M., Kamada, M., Nakata, M., & et al. (2001). Effects of high nitrogen load on growth, photosynthesis, and nutrient status of *Cryptomeria japonica* and *Pinus densiflora* seedlings. *Trees*, *15*, 453-461.
- Munzi, S., Pisani, T., & Loppi, S. (2009). The integrity of lichen cell membrane as a suitable parameter for monitoring biological effects of acute nitrogen pollution. *Ecotoxicology and Environmental Safety*, *72*, 1085-1090.
- Rahmat, M., Maulina, W., Rustami, E., Azis, M., Budiarti, D. R., Seminar, K. B., & et al. (2013). Performance in real conditions of photonic crystal sensor-based NO₂ gas monitoring system. *Atmospheric Environment*, *79*, 480-485.
- Ramadhan, H., Khalam, H., Marselina, N., Nirmala, T., Aida, N., & Ratih, A. (2023). Analysis of economic growth on carbon dioxide gas emissions in G20 countries. *Asian Journal of Economics, Business and Accounting*, *23*(14), 1-7. <https://doi.org/10.9734/ajeba/2023/v23i141000>
- Rao, N. K., Laxman, R. H., & Bhatt, R. M. (2010). Impact of climate change on vegetable crops. In H. P. Singh, J. P. Singh, & S. S. Lal (Eds.), *Challenges of climate change in Indian horticulture* (pp. 113-123). Westville Publishing House.
- Ravindranath, N. H., Murthy, I. K., Sudha, P., Ramprasad, V., Nagendra, M. D. U., Sahana, C. A., Srivatha, K. G., & Khan, H. (2007). Methodological issues in forestry mitigation projects: A case study of Kolar districts. *Mitigation and Adaptation Strategies for Global Change*, *12*, 1077-1090.
- Schmidt, G. A., Ruedy, R. A., Miller, R. L., & Lacis, A. A. (2010). Attribution of the present-day total greenhouse effect. *Journal of Geophysical Research*, *115*, D20106. <https://doi.org/10.1029/2010JD014287>
- Sheng, Q., & Zhu, Z. (2019). Effects of nitrogen dioxide on biochemical responses in 41 garden plants. *Plants*, *8*, 45. <https://doi.org/10.3390/plants8020045>
- Singh, J. P., Govindakrishna, P. M., Lal, S. K., & Aggarwal, P. K. (2008). Increasing the efficiency of agronomy experiments in potatoes using the INFOCROP-POTATO model. *Potato Research*, *48*, 131-152.
- Somarrriba, E., Cerda, R., Orozco, L., Cifuentes, M., & Dávila, H. (2013). Carbon stocks and cocoa yields in agroforestry systems of Central America. *Agriculture, Ecosystems & Environment*, *173*, 46-57.
- Srivastava, A. K., Das, S. N., Malhotra, S. K., & Majumdar, K. (2014). SSNM-based rationale of fertilizer use in perennial crops: A review. *Indian Journal of Agricultural Sciences*, *84*(1), 3-17.
- Swanepoel, J. W., Krüger, G. H. J., & Van Heerden, P. D. R. (2007). Effects of sulphur dioxide on photosynthesis in the

- succulent *Augea capensis* Thunb. *Journal of Arid Environments*, 70, 208-221.
- Venkateswarlu, B., & Shanker, A. K. (2012). Dryland agriculture: Bringing resilience to crop production under changing climate. In *Crop Stress and its Management: Perspectives and Strategies* (pp. 19-44). Springer, Netherlands.
- Verma, A., & Singh, S. N. (2006). Biochemical and ultra-structural changes in plant foliage exposed to auto-pollution. *Environmental Monitoring and Assessment*, 120, 585-602.
- Vighi, I. L., Benitez, L. C., Amaral, M. N., Moraes, G. P., Auler, P. A., Rodrigues, G. S., & Braga, E. J. B. (2017). Functional characterization of the antioxidant enzymes in rice plants exposed to salinity stress. *Biologia Plantarum*, 61, 540-550.
- Wuebbles, D. J., & Hayhoe, K. (2002). Atmospheric methane and global change. *Earth-Science Reviews*, 57, 177-210. [https://doi.org/10.1016/S0012-8252\(01\)00062-9](https://doi.org/10.1016/S0012-8252(01)00062-9)

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/125786>