



# Evaluation of Indoor Radon and It's Health Risks Parameters within Azuabie, Trans-Amadi and Nkpogu Towns, in Port Harcourt, Rivers State, Nigeria

Briggs-Kamara, Margaret Apaemi, Briggs<sup>1</sup>, Sigalo, Friday Barikpe<sup>1</sup>, Iyeneomie, Tamunobereton-Ari<sup>1</sup> and Orlunta, Aloysius Ndubisi<sup>1\*</sup>

<sup>1</sup>Department of Physics, Faculty of Science, Rivers State University, Nkpolu Oroworukwo, Port Harcourt, Rivers State, Nigeria.

## Authors' contributions

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## Article Information

DOI: 10.9734/AJOPACS/2021/v9i230133

Editor(s):

(1) Dr. Thomas F. George, University of Missouri- St. Louis One University Boulevard St. Louis, USA.

Reviewers:

(1) Oumar Bobbo Modibo, Hirosaki University, Japan.

(2) Mohammed D Alenezy, Aljouf University, Saudi Arabia.

(3) Wiseman Bekelesi, Japan.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/67820>

Original Research Article

Received 18 March 2021

Accepted 22 May 2021

Published 02 June 2021

## ABSTRACT

Evaluation of indoor radon level and its health risk parameters has been carried out in three communities Azuabie, Trans-Amadi and Nkpogu towns in Port Harcourt, Rivers State, Nigeria. A pocket sized Corentium Arthings digital radon detector meter was used to record the indoor radon concentration levels. The geographical coordinates were recorded using a hand-held geographical positioning system (GPS) for the various sample points. A total of 30 sample points were evaluated, with 10 sample points for each town respectively. The results of the concentration levels showed that for Azuabie (AZ) town, the concentration level varied from 6.660 Bqm<sup>-3</sup> to 13.690 Bqm<sup>-3</sup> with an average of 10.65±0.95Bqm<sup>-3</sup>. Nkpogu (NK) town the results of the indoor concentration level ranged from 9.250 Bqm<sup>-3</sup> to 18.870 Bqm<sup>-3</sup> with an average of 13.32±1.02 Bqm<sup>-3</sup>, Nkpogu (NK) town, the indoor concentration level ranged from 7.030 Bqm<sup>-3</sup> to 20.350 Bqm<sup>-3</sup> with an average of

\*Corresponding author: E-mail: [aloyndubisi@gmail.com](mailto:aloyndubisi@gmail.com);

$12.25 \pm 1.34 \text{ Bq m}^{-3}$ . The annual absorbed dose for Azuabie, Trans-Amadi and Nkpogu varied as follows,  $1.680 \text{ mSvy}^{-1} - 3.921 \text{ mSvy}^{-1}$ ,  $2.334 \text{ mSvy}^{-1} - 47610 \text{ mSvy}^{-1}$  and  $1.774 \text{ mSvy}^{-2} - 5.134 \text{ mSvy}^{-1}$  respectively. The annual effective dose rate for the three towns ranged from  $0.403 \text{ mSvy}^{-1} - 0.941 \text{ mSvy}^{-1}$ ,  $0.560 \text{ mSvy}^{-1} - 1.143 \text{ mSvy}^{-1}$  and  $0.426 \text{ mSvy}^{-1} - 1.143 \text{ mSvy}^{-1}$ . The excess life time cancer risk varied from  $1.4117 - 3.294$ ,  $1.9607 - 3.999$  and  $1.4901 - 3.999$  respectively. The results of the indoor concentration levels annual and the absorbed dose and the annual effective dose rate are all below the ICRP safe limit. However, the results of the excess life time cancer risk are all higher than the ICRP safe standard limit of  $0.029 \times 10^{-3}$ .

**Keywords:** Background Ionization radiation; annual effective dose; excess life cancer risk.

## 1. INTRODUCTION

Radon is a radioactive gas which is colourless, odourless and tasteless. It is predominantly found in rock samples, bedrock formations, soil and ground water all over the world, [1]. Radon-222 is the immediate decay product of radium-226 during the decay series of Uranium – 238 which is the source of radon-222. It has a half-life of 3.81 days. Radon-222 decays to polonium – 298 with the release of an alpha particle and consequently to other progenies of radon or radon daughters [2,3]. Radon has been classified by the international Agency for Research on cancer (I A R C, 1988) as a carcinogenic gas.

The World Health Organization (WHO, 1999) also classified radon as the second highest cause of lung cancer after cigarette smoking. It was estimated that radon-222 causes between 3% to 14% of all lung cancers. Radon is the main cause of lung cancer among non-smokers. It is the cause of about fifteen per cent of all lung cancer cases throughout the world, (WHO, 2009). People are exposed to high radon level in small houses than in bigger apartments (UNSCEAR, 2000). The entry routes of indoor into the houses includes, the door, the windows, cracks on the walls, sinks, basements, floors etc [4-6].

The health effects as a result of higher exposure to radon-222 radiation induced. Inhalation is the main route through which humans are exposure to radon radiation [7-10]. The dose contribution to radon may be small, but the inhalation of the progenies of radon which have very short half-life can deposit non-homogenously in the human respiratory track and irradiate the bronchial epithelium which is usually very harmful [11-13].

Two of the radon progenies (daughters) Polonium-214 and Polonium-218 releases the highest amount of alpha radiation dose to the lungs (NCRP, 1994). When these radioactive

particles settle in a person's lung, they can cause damage to the mucosa linings of the lungs. A prolonged exposure to radon-222 radiation can also lead to series of damage to the pulmonary mucosa which could result to lung cancer [14-16].

An individual's average radiation come from the decay products of radon -222 such as , polonium – 218 and polonium – 214. These products of radon – 222 are in the solid form. They could be attached or unattached to the surface of aerosols, dusts and smoke particles. They can adhere deeply or stick to the lungs where they irradiate and penetrate the cells of the mucosa membranes, bronchi and pulmonary tissues. The ionizing radiation which emanate from the radioactive decay of radon-222 begins the process of carcinogenesis in the human lungs [17-19].

It is estimated that about 90% of the radon progenies can attach to airborne particles [20]. Also, the unattached fraction of the particles which constitutes about 10% has a higher rate of deposition and is more efficient in delivering doses to the sensitive cells of the lungs (UNSCEAR, 1993). The more the concentration of radon in homes the greater the risk of lung cancer due to radon exposure. The risk of lung cancer increases for every  $100 \text{ Bq/m}^3$  increase in radon concentration, according to the World Health Organization (WHO, 2016).

However, there is no thresholds value below which radon exposure carries no risk of lung cancer. Therefore it is important that every country should set up a national reference level "AS Low As Reasonably Achievable" (ALARA) [21].

The reason for carrying out radon measurements in these towns is because radon-222 concentration level has not been done within Port Harcourt local government. Therefore, we




Fig. 1. Map of Study Area

wanted to find out the radon-222 concentration level and compare the results with international set standards. Also, measurement of radon-222 concentration level in these areas will provide baseline studies and literature for other researchers. It is well known fact that these areas are densely populated with building styles that are poorly ventilated therefore, it was necessary to find out if the residents of these towns are exposed to higher dose of radon-222 from their environment. Additionally, there has been incident of lung cancer cases within these areas in recent times even though not officially reported, therefore it was important to confirm if it was as a result of high level of radon exposure.

## 2. MATERIALS AND METHODS

The radon detector used in this work is Airthings Corentium Digital Radon Detector designed in Oslo Norway in the year 2019. It is calibrated to measure radon-222 in picocurie per litre (pCi/L). The Airthings Corentium digital detector can be used to monitor radon concentration levels for a minimum period of 48 hours, for daily, weekly, and monthly and yearly. The radon meter works on the principle that the radon diffuses into a detection chamber. As the atom decay they emit

energetic alpha particles. The energetic alpha particles are detected by a silicon photodiode. The alpha particle generates a small signal current when it hits the photodiode. By the use of a low power amplifier stage, the signal current is converted into a large voltage signal. The maximum amplitude of the voltage signal is detected and sampled by an analog to digital converter (ADC). The amplitude of this signal is proportional to the energy of the alpha particle that hit the photodiode. The brain of the monitor is a micro-controller which registers the time and the energy of every detected particle. This information is used to calculate the mean Radon concentration for, daily, weekly, monthly and yearly periods.

Indoor radon concentration measurement were carried out in three communities in Port Harcourt using the Airthings digital radon detector while the geographical position system (GPS) was used for the measurement of the geographical coordinates of the sample points. A total of 30 sample points were measured for the three communities, 10 sample points for Azuabie town, 10 sample points for Nkpogwu town and 10 sample points for Woji town.

The Airthings digital radon detector was placed in the dwelling room at a height of about 50cm above the ground and about 150cm from both the window and door and 25cm from the walls, [22]. The detector was kept for a period of two days (48hrs.) in a given dwelling before relocating to another house or home.

The windows and doors were kept closed throughout the period of the measurement to ensure that the indoor air is not distorted to achieve accuracy within the period of 48 hours.

## 2.1 Radon Risk Parameters

### 2.1.1 Annual absorbed dose from radon concentration

The annual absorb dose from the radon concentration is the magnitude of energy delivered in a tissue from exposure to ionizing radiation within a specified period. The physics of the equation employed in the computation of the annual absorbed dose rate due to report of United Nation's Scientific Committee on the Effects of Atomic Radiation [20] (UNSCEAR, 2000) and is given as;

$$DRn(mSvy^{-1}) = CRn. D.H.T.F \quad (1)$$

Where:

CRn = the measured concentration of indoor radon-222 in Bqm<sup>-3</sup>,  
 F = Radon-222 equilibrium indoor factor of (0.4)  
 T = Is the indoor-222 occupancy time 7000hr, (0.8 x 24 hr x 365)  
 H = Indoor radon-222 occupancy factor (0.4)  
 D = Dose conversion factor (9.0x 10<sup>-6</sup> mSv/hr per Bqm<sup>-3</sup>)

### 2.1.2 Annual Effective Dose Rate (AEDR) from radon concentration

The annual effective dose rate was calculated by applying a tissue and radiation weighting factor (ICRA 1991). The inhalation dose equation is given by;

$$AEDR (mSvy^{-1}) = DRn. WR. WT \quad (2)$$

Where:

DRn = Indoor radon concentration

WR = Radiation weighting factor for alpha particles (20)  
 WT = Tissue weighting factor for the lung (0.12)

### 2.1.3 Excess life cancer risk from radon concentration

The excess life time cancer risks (ELCR) is the potential carcinogenic effects, from the calculation based on probability of cancer induced incidence in a population. It indicates the chances of contracting a cancer from the exposure from radiation or toxic chemical substances for a specific life time. According to [23], the excess life time risk was calculated from the equation.

$$ECLR = AEDR \times DL \times RF \quad (3)$$

Where:

DL = Average duration of life (70 years)  
 RF = Risk Factor 0.05 x 10<sup>-5</sup>

## 2.2 Statistical Analysis

The mean values and standard deviation for the radon concentrations was computed using equations (4) and (5). This can also be computed with the aid of a histogram and contour maps showing the spatial distribution of radon concentration and its health risk parameters which will also be plotted with the aid of statistical package for social sciences (SPSS) version (22) and surfer – 8 contour map software.

$$\text{Mean Value } \mu_x = \frac{\sum_i^n x}{n} \quad (4)$$

$$\text{Standard Deviation (S.D)} = \sqrt{\frac{\sum_i^n (x_i - \mu_x)^2}{n-1}} \quad (5)$$

## 3. RESULTS AND DISCUSSION

The result of the indoor radon concentration and their geographical coordinates for the three towns; Azuabie, Trans-Amadi, and Nkpogu town are presented in Tables, 1 - 3 respectively. The indoor radon concentration was measured in PicoCuri per litre (PCi/L). Tables 4 – 9 represent the results of the concentration in Becqueri per metre cube with their health risk parameters.

Also, the indoor radon concentration level for Nkpogu town varied from 6.660Bq/m<sup>3</sup> to 13.690 Bq/m<sup>3</sup> with an average indoor radon concentration of 10.65±0.95 Bq/m<sup>3</sup>; the result of

indoor radon concentration level in Trans-madi varied from 9.250 Bq/m<sup>3</sup> to 18.870B/m<sup>3</sup> with an average of 13.32±1.02Bq/m<sup>3</sup>; The indoor radon concentration level at Nkpogu town ranged from 7.030 to 20.350 Bq/m<sup>3</sup> with an average indoor concentration level of 12.25±1.34 Bq/m<sup>3</sup>. The annual absorbed dose for the indoor radon varied as follows; 1.68 mSv/yr-3.921 mSv/yr, 2.334 mSv/yr-4.761 mSv/yr, 1.774 mSv/yr-5.134 mSv/yr for Azuabie town, Trans-Amadi town and Nkpogu town respectively. The annual effective dose rate (AEDR) for the three towns which was calculated from the indoor radon concentration ranged from 0.403mSvm/yr- 0.941mSv/y; 0.560 mSv/yr-1.143mSv/yr; 0.426mSv/yr-1.143mSv/yr respectively. These values are all below the safe standard by International Commission on Radiological Protection (ICRP). The Excess Life Time Cancer Risk calculated from the indoor radon concentration were; 1.4117-3.294; 1.9607-3.999 and 1.4901-3.999 for Azuabie, Trans-Amadi and Nkpogu respectively. The results of the Excess Life Time Cancer Risk for the three towns were all higher than the world average.

The mean annual absorbed dose for the three towns, Azuabie, Trans-Amadi and Nkpogu are 2.69±0.24mSv/yr, 3.36±0.26mSv/yr and 3.08±0.34mSv/yr. Also, the average Annual Absorbed Dose Rate ( AEDR) for Azuabie town, Trans-Amadi town and Nkpogu town are 0.64±0.05mSv/yr, 0.81±0.06mSv/yr and 0.74±0.08mSv/yr respectively. The average Excess Life Time Cancer Risk calculated for Azuabie was (2.26±0.20)×10<sup>-3</sup>, Trans-Amadi was (2.82±0.22)×10<sup>-3</sup>, Nkpogu was (2.60±0.28)×10<sup>-3</sup>. These values are all higher than the world average. These results have been represented in their respective tables and barcharts. The excess life time cancer risk were

calculated for the age duration of 70 ,60, 50,40 and 30 years. The result show that the higher the years the higher the risk of cancer due to exposure to rado-222.

From the result obtained, there is no appreciable difference between the different study areas, this is due to the fact that there is similarity in the building design, materials used in the building, ventilation pattern of the living houses and the life style of the people. These results obtained are comparable to the indoor radon concentration level for different types of buildings in covenant university, Nigeria (Usilan *et al.*, 2020). The mean radon concentration for the three different building types Glass, Brick house and Basement house were 14.96Bqm<sup>-3</sup>, 10.74 Bqm<sup>-3</sup> and 144,6Bqm<sup>-3</sup> respectively.

The indoor radon concentration for the glass house ranged from 11.03 – 17.46Bqm<sup>-3</sup>, the concentration level of brick house varied from 6.62 – 20.85Bqm<sup>-3</sup>, while that for basement house ranged from 15.75 to 614.52Bqm<sup>-3</sup>. However, the concentration level for the basement house was above the international safe standard, but the glass house and brick house indoor concentration levels were of the same range for the Azuabie, Trans-Amadi and Nkpogu locations in Port Harcourt. Also, the indoor radon level concentration measured by(Olusegun *et al.*, 2015) in Obasfemi Awolowo University Ile-Ife are similar to the indoor radon level measured in Azuabie, Trans-Amadi and Nkpogu town. The radon concentration level obtained from the offices varied from 0.0 - 5.3Bqm<sup>-3</sup>. The average value obtained was 0.9 PCi/L.

**Table 1. Radon levels in azuabie town (AZB)**

S/N	Azuabie town	Gps coordinates		Radon levels IN pico curi PER litre (PC <sub>l</sub> /L)
		Latitudes	Longitudes	
1.0	AZB <sub>1</sub>	N4 <sup>0</sup> 48 <sup>1</sup> 35.832 <sup>11</sup>	E7 <sup>0</sup> 3 <sup>1</sup> 9.594 <sup>11</sup>	0.18
2.0	AZB <sub>2</sub>	N4 <sup>0</sup> 48 <sup>1</sup> 39.914 <sup>11</sup>	E7 <sup>0</sup> 3 <sup>1</sup> 92.242 <sup>11</sup>	0.22
3.0	AZB <sub>3</sub>	N4 <sup>0</sup> 48 <sup>1</sup> 34.32 <sup>11</sup>	E7 <sup>0</sup> 3 <sup>1</sup> 5.316 <sup>11</sup>	0.32
4.0	AZB <sub>4</sub>	N4 <sup>0</sup> 48 <sup>1</sup> 26.136 <sup>11</sup>	E7 <sup>0</sup> 3 <sup>1</sup> 13.626 <sup>11</sup>	0.26
5.0	AZB <sub>5</sub>	N4 <sup>0</sup> 48 <sup>1</sup> .29.28 <sup>11</sup>	E7 <sup>0</sup> 3 <sup>1</sup> 13.404 <sup>11</sup>	0.37
6.0	AZB <sub>6</sub>	N4 <sup>0</sup> 48 <sup>1</sup> 13.98 <sup>11</sup>	E7 <sup>0</sup> 3 <sup>1</sup> 4.902 <sup>11</sup>	0.33
7.0	AZB <sub>7</sub>	N4 <sup>0</sup> 48 <sup>1</sup> 20.304 <sup>11</sup>	E7 <sup>0</sup> 2 <sup>1</sup> 59.154 <sup>11</sup>	0.35
8.0	AZB <sub>8</sub>	N4 <sup>0</sup> 48 <sup>1</sup> 10.758 <sup>11</sup>	E7 <sup>0</sup> 3 <sup>1</sup> 1.374 <sup>11</sup>	0.19
9.0	AZB <sub>9</sub>	N4 <sup>0</sup> 48 <sup>1</sup> 13.632 <sup>11</sup>	E7 <sup>0</sup> 3 <sup>1</sup> 2.388 <sup>11</sup>	0.42
10.0	AZB <sub>10</sub>	N4 <sup>0</sup> 48 <sup>1</sup> 14.772 <sup>11</sup>	E7 <sup>0</sup> 3 <sup>1</sup> 1.824 <sup>11</sup>	0.24

**Table 2. Radon levels in trans-amadi/ogoniba town**

S/N	Location, trans amadi/ogonigba (TA)	GPS coordinates		Radon levels in pico curi per litre (PC <sub>i</sub> /L)
		Latitudes	Longitudes	
1.0	TA <sub>1</sub>	N4 <sup>0</sup> 49 <sup>1</sup> 42.936 <sup>11</sup>	E7 <sup>0</sup> 2 <sup>1</sup> 22.878 <sup>11</sup>	0.51
2.0	TA <sub>2</sub>	N4 <sup>0</sup> 49 <sup>1</sup> 40.818 <sup>11</sup>	E7 <sup>0</sup> 2 <sup>1</sup> 19.488 <sup>11</sup>	0.25
3.0	TA <sub>3</sub>	N4 <sup>0</sup> 49 <sup>1</sup> 26.798 <sup>11</sup>	E7 <sup>0</sup> 2 <sup>1</sup> 29.934 <sup>11</sup>	0.33
4.0	TA <sub>4</sub>	N4 <sup>0</sup> 49 <sup>1</sup> 21.252 <sup>11</sup>	E7 <sup>0</sup> 2 <sup>1</sup> 24.990 <sup>11</sup>	0.34
5.0	TA <sub>5</sub>	N4 <sup>0</sup> 48 <sup>1</sup> 3.744 <sup>11</sup>	E7 <sup>0</sup> 3 <sup>1</sup> 25.836 <sup>11</sup>	0.37
6.0	TA <sub>6</sub>	N4 <sup>0</sup> 49 <sup>1</sup> 33.492 <sup>11</sup>	E7 <sup>0</sup> 2 <sup>1</sup> 28.902 <sup>11</sup>	0.31
7.0	TA <sub>7</sub>	N4 <sup>0</sup> 49 <sup>1</sup> 37.056 <sup>11</sup>	E7 <sup>0</sup> 2 <sup>1</sup> 25.506 <sup>11</sup>	0.34
8.0	TA <sub>8</sub>	N4 <sup>0</sup> 49 <sup>1</sup> 34.374 <sup>11</sup>	E7 <sup>0</sup> 2 <sup>1</sup> 19.668 <sup>11</sup>	0.28
9.0	TA <sub>9</sub>	N4 <sup>0</sup> 49 <sup>1</sup> 33.702 <sup>11</sup>	E7 <sup>0</sup> 2 <sup>1</sup> 28.308 <sup>11</sup>	0.36
10.0	TA <sub>10</sub>	N4 <sup>0</sup> 49 <sup>1</sup> 39.006 <sup>11</sup>	E7 <sup>0</sup> 2 <sup>1</sup> 15.642 <sup>11</sup>	0.32

**Table 3. Radon levels in nkpogwu town (NK)**

S/N	Nkpogwu town	GPS coordinates		Radon levels in pico curi per litre (PC <sub>i</sub> /L)
		Latitudes	Longitudes	
1.0	NK <sub>1</sub>	N4 <sup>0</sup> 48 <sup>1</sup> 3.921080 <sup>11</sup>	E7 <sup>0</sup> 1 <sup>11</sup> 15.12055 <sup>11</sup>	0.51
2.0	NK <sub>2</sub>	N4 <sup>0</sup> 48 <sup>1</sup> 5.732280 <sup>11</sup>	E7 <sup>0</sup> 1 <sup>1</sup> 13.724620 <sup>11</sup>	0.33
3.0	NK <sub>3</sub>	N4 <sup>0</sup> 48 <sup>1</sup> 6.435790 <sup>11</sup>	E7 <sup>0</sup> 1 <sup>1</sup> 16.291020 <sup>11</sup>	0.28
4.0	NK <sub>4</sub>	N4 <sup>0</sup> 48 <sup>1</sup> 6.293740 <sup>11</sup>	E7 <sup>0</sup> 1 <sup>1</sup> 11.700480 <sup>11</sup>	0.26
5.0	NK <sub>5</sub>	N4 <sup>0</sup> 48 <sup>1</sup> 3.573170 <sup>11</sup>	E7 <sup>0</sup> 1 <sup>1</sup> 806956000 <sup>11</sup>	0.29
6.0	NK <sub>6</sub>	N4 <sup>0</sup> 48 <sup>1</sup> 5.524240 <sup>11</sup>	E7 <sup>0</sup> 1 <sup>1</sup> 28.163170 <sup>11</sup>	0.55
7.0	NK <sub>7</sub>	N4 <sup>0</sup> 48 <sup>1</sup> 3.389317 <sup>11</sup>	E7 <sup>0</sup> 1 <sup>1</sup> 59.070420 <sup>11</sup>	0.19
8.0	NK <sub>8</sub>	N4 <sup>0</sup> 48 <sup>1</sup> 3.292538 <sup>11</sup>	E7 <sup>0</sup> 059.9982000 <sup>11</sup>	0.32
9.0	NK <sub>9</sub>	N4 <sup>0</sup> 48 <sup>1</sup> 12.93592 <sup>11</sup>	E7 <sup>0</sup> 1 <sup>1</sup> 9.1599200 <sup>11</sup>	0.34
10.0	NK <sub>10</sub>	N4 <sup>0</sup> 48 <sup>1</sup> 4.817770 <sup>11</sup>	E7 <sup>0</sup> 1 <sup>1</sup> 18.051850 <sup>11</sup>	0.24

**Table 4. Computed values of annual effective dose and annual equivalent dose rate of Azuabie Town (AZB)**

S/N	S/Pts	CRN (PC <sub>i</sub> /L)	CRN (Bq/m <sup>3</sup> )	DRn (mSvy-1)	AEDR (mSvy-1)
1.0	AZB1	0.18	6.6600	1.6802	0.403
2.0	AZB2	0.22	8.1400	2.0536	0.493
3.0	AZB3	0.32	11.8400	2.9871	0.717
4.0	AZB4	0.26	9.6200	2.4270	0.582
5.0	AZB5	0.37	13.6900	3.4538	0.829
6.0	AZB6	0.33	12.2100	3.0804	0.739
7.0	AZB7	0.35	12.9500	3.2671	0.784
8.0	AZB8	0.19	7.0300	1.7736	0.426
9.0	AZB9	0.42	15.5400	3.9206	0.941
10.0	AZB10	0.24	8.8800	2.2403	0.538
AVERAGE		0.29 ± 0.02	10.65 ± 0.95	2.69 ± 0.24	0.64 ± 0.05

Similarly, the health risk parameters, the annual absorbed dose, annual equivalent dose rate and the excess life time cancer risk agreed with the calculated value in Azuabie, Trans-Amadi and Nkpogwu towns. The indoor radon level concentration measured at Azuabie, Trans-Amadi and Nkpogwu towns is also comparable to the work done earlier by (Sokari, S.A. 2018) in Okirika local government area in River's state, Nigeria. The average concentration level was 19.36±Bqm<sup>-3</sup> for mud house, while the overall

average indoor radon concentration was 11.37±3.28Bqm<sup>-3</sup> for Okirika local government.

These results are lower than the safe limit by ICRP. Also, the overall average of the annual absorbed dose was 3.36±0.26mSv/yr, the mean annual effective dose rate was 0.15±0.42 mSv/yr while the overall excess life time cancer risk calculate from the indoor radon concentration level was (0.52±0.15)×10<sup>-3</sup> and were all above the world standard of 0.029 x10<sup>-3</sup>.

**Table 5. Computed values of excess lifetime cancer risk for ages of 70(Std), 60, 50, 40 and 30 yrs of Azuabie Town (AZB)**

S/N	S/Pts	ELCR x10 <sup>-3</sup> 70 yrs(Std)	ELCR x10 <sup>-3</sup> 60 yrs	ELCR x10 <sup>-3</sup> 50 yrs	ELCR x10 <sup>-3</sup> 40 yrs	ELCR x10 <sup>-3</sup> 30 yrs
1.0	AZB1	1.412	1.210	1.008	0.807	0.605
2.0	AZB2	1.725	1.479	1.232	0.986	0.739
3.0	AZB3	2.510	2.151	1.792	1.434	1.075
4.0	AZB4	2.039	1.747	1.456	1.165	0.874
5.0	AZB5	2.902	2.487	2.072	1.658	1.243
6.0	AZB6	2.588	2.218	1.848	1.479	1.109
7.0	AZB7	2.745	2.352	1.960	1.568	1.176
8.0	AZB8	1.490	1.277	1.064	0.851	0.638
9.0	AZB9	3.294	2.823	2.352	1.882	1.411
10.0	AZB10	1.882	1.613	1.344	1.075	0.807
AVERAGE		2.25± 0.20	1.94 ± 0.17	1.61 ± 0.14	1.29 ± 0.11	0.97 ± 0.08

**Table 6. Computed values of annual effective dose and annual equivalent dose rate of Trans-Amadi/Ogoniba Town**

S/N	S/Pts	CRN (PCi/L)	CRN (Bq/m3)	DRn (mSvy-1)	AEDR (mSvy-1)
1.0	TA1	0.51	18.8700	4.7607	1.143
2.0	TA2	0.25	9.2500	2.3337	0.560
3.0	TA3	0.33	12.2100	3.0804	0.739
4.0	TA4	0.34	12.5800	3.1738	0.762
5.0	TA5	0.37	13.6900	3.4538	0.829
6.0	TA6	0.31	11.4700	2.8937	0.694
7.0	TA7	0.34	12.5800	3.1738	0.762
8.0	TA8	0.28	10.3600	2.6137	0.627
9.0	TA9	0.36	13.3200	3.3605	0.807
10.0	TA10	0.32	18.8700	4.7607	1.143
AVERAGE		0.34 ± 0.03	13.32 ± 1.02	3.36 ± 0.26	0.81 ± 0.06

**Table 7. Computed values of excess lifetime cancer risk for ages of; 70(Std), 60, 50, 40 and 30 yrs of Trans-Amadi/Ogoniba Town**


S/N	S/Pts	ELCR x10 <sup>-3</sup> 70 yrs(Std)	ELCR x10 <sup>-3</sup> 60 yrs	ELCR x10 <sup>-3</sup> 50 yrs	ELCR x10 <sup>-3</sup> 40 yrs	ELCR x10 <sup>-3</sup> 30 yrs
1.0	TA <sub>1</sub>	4.000	3.428	2.856	2.285	1.714
2.0	TA <sub>2</sub>	1.961	1.680	1.400	1.120	0.840
3.0	TA <sub>3</sub>	2.588	2.218	1.848	1.479	1.109
4.0	TA <sub>4</sub>	2.667	2.285	1.904	1.523	1.143
5.0	TA <sub>5</sub>	2.902	2.487	2.072	1.658	1.243
6.0	TA <sub>6</sub>	2.431	2.083	1.736	1.389	1.042
7.0	TA <sub>7</sub>	2.667	2.285	1.904	1.523	1.143
8.0	TA <sub>8</sub>	2.196	1.882	1.568	1.255	0.941
9.0	TA <sub>9</sub>	2.823	2.420	2.016	1.613	1.210
10.0	TA <sub>10</sub>	2.510	2.151	1.792	1.434	1.075
AVERAGE		2.67 ± 0.54	2.29 ± 0.47	1.90 ± 0.39	1.52 ± 0.10	1.14 ± 0.23

**Table 8. Computed values of annual effective dose and annual equivalent dose rate of Nkpgwu Town (NK)**

S/N	S/Pts	C <sub>RN</sub> (PCi/L)	C <sub>RN</sub> (Bq/m3)	D <sub>Rn</sub> (mSvy-1)	AEDR (mSvy-1)
1.0	NK <sub>1</sub>	0.51	18.8700	4.7607	1.143
2.0	NK <sub>2</sub>	0.33	12.2100	3.0804	0.739
3.0	NK <sub>3</sub>	0.28	10.3600	2.6137	0.627
4.0	NK <sub>4</sub>	0.26	9.6200	2.4270	0.582
5.0	NK <sub>5</sub>	0.29	10.7300	2.7071	0.650
6.0	NK <sub>6</sub>	0.55	20.3500	5.1341	1.232
7.0	NK <sub>7</sub>	0.19	7.0300	1.7736	0.426
8.0	NK <sub>8</sub>	0.32	11.8400	2.9871	0.717
9.0	NK <sub>9</sub>	0.34	12.5800	3.1738	0.762
10.0	NK <sub>10</sub>	0.24	8.8800	2.2403	0.538
AVERAGE		0.33 ± 0.04	12.25± 1.34	3.08 ± 0.34	0.74± 0.08

**Table 9. Computed values of excess lifetime cancer risk for ages of 70(Std),60, 50, 40 and 30 yrs of Nkpgwu Town (NK)**

S/N	S/Pts	ELCR x10 <sup>-3</sup> 70 yrs(Std)	ELCR x10 <sup>-3</sup> 60 yrs	ELCR x10 <sup>-3</sup> 50 yrs	ELCR x10 <sup>-3</sup> 40 yrs	ELCR x10 <sup>-3</sup> 30 yrs
1.0	NK <sub>1</sub>	4.000	3.428	2.856	2.285	1.714
2.0	NK <sub>2</sub>	2.588	2.218	1.848	1.479	1.109
3.0	NK <sub>3</sub>	2.196	1.882	1.568	1.255	0.941
4.0	NK <sub>4</sub>	2.039	1.747	1.456	1.165	0.874
5.0	NK <sub>5</sub>	2.274	1.949	1.624	1.299	0.975
6.0	NK <sub>6</sub>	4.314	3.697	3.080	2.464	1.848
7.0	NK <sub>7</sub>	1.490	1.277	1.064	0.851	0.638
8.0	NK <sub>8</sub>	2.510	2.151	1.792	1.434	1.075
9.0	NK <sub>9</sub>	2.667	2.285	1.904	1.523	1.143
10.0	NK <sub>10</sub>	1.882	1.613	1.344	1.075	0.807
AVERAGE		2.59± 0.28	2.23± 0.24	1.85± 0.20	1.48± 0.16	1.11 ± 0.12



**Fig. 2. Average Indoor Radon Level of sample locations and ICPR reference level**






Fig. 3. Annual absorbed dose of sample locations and ICRP Standard




Fig. 4. Annual effective dose rate (AEDR) and ICRP standard




Fig. 5. Excess life time cancer risk (ELCR) of sample locations with world standard

#### 4. CONCLUSION

The evaluation of indoor radon concentration level and its health risk parameters has been done in Azuabie town, Nkpogu town and Trans-Amadi in Port Harcourt, Rivers State, Nigeria using the Corentium Arthings radon digital detector. The concentration levels measured for the three towns were all below the action level of 200-600 Bq/m<sup>3</sup> given by the International Commission on Radiological Protection (ICRP). However, the excess life time cancer risk for all the three towns were all higher than the world average. It was also observed that there was no appreciable difference in the indoor radon concentration level between the three towns of Azuabie, Trans-Amadi, Nkpogu. This is because of the similarities in the building styles, materials used for the buildings, the life style of the people and the ventilation method.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Ojo TJ, Ajayi I. Outdoor production of radon from the uranium-238 decay serie; 2015. Available:www.Nuclearsafety.Gc.Ca On 17<sup>th</sup> April, 2019
- Angre CY, Miyittah M, Andam AB. Dose Assessment of indoor radon concentration level in the southern Dayi District of the volta region, Ghana; 2015. URL:http://197.255.68.203/handle/123456789/8217
- Anita PS, Davo S, Nikola S, Dragana D, Tonislav T. Short term measurement of indoor radiation concentration in Northern Croatia. *Journal of Applied Sciences*. 2020;10(7):104-115.
- Abu – Haija O, Salameh B, Abdul-Wali A, Abdelsalam M, Al-Ebarisat H. Measurement of radon province. *Jordan International Journal of the Physical Sciences*. 2010;5(6):696–699.
- Adegun IK, Anyaegbuna BE, Olayemi OA, Jolayemi TS, Ibiwoye MO. Radon concentration assessment in bank cellars in three Nigerian cities. *Nigerian Journal of Technology (NIJOTECH)*. 2019;38(4):957–964.
- Akabuogu EU, Nwaokoro E, Oni EA. Measurement and analysis of indoor radon level at a university in South-Eastern Nigeria. *International Research Journal of Pure and Applied Physics*. 2019;6(1):8–17.
- Arhin I. Assessment of indoor radon levels and emanation from soil in Abirem community in the Eastern region of Ghana. *Journal of International Nuclear Information system, INIS(IAEA)*. 2018;50(4):2910-2916.
- Chutima K, Yuki T, Masahiro H, Sinji T. Importance of discriminative map for dom Isotopes and its utilization in the environment and lessons learned from using RADUET monitor. *International Journal Res. Public Health*. 2020;17(11):4141-4146.
- Badoe LG, Andam A, Sosu EK. Measurement of radon gas level in ashanti and design of a radon vulnerability map for Ghana. *Journal of Environmental Radioactivity*. 2012;148:154-162.
- Environmental Protection Agency. A citizen's guide to Radon. Retrieved; 2016.
- Okoji MC, Agwu KK, Idigo FU, Anakwe AC. Measurement of indoor radon concentration levels in offices of University of Nigeria, Enugu Campus. *Journal of Association of Radiographers of Nigeria*. 2013;27(1):32–37.
- Production of Radon From the Uranium-238 Decay Series Available:www.Nuclearsafety.Gc.Com On 17<sup>th</sup> April, 2019
- Radon Movement and Pathways. Available:Www.Legacyinspect.com, On 17<sup>th</sup> April 2019
- National Bureau of Statistics. Annual Abstract of Statistics; 2011.
- Nigus MD. Indoor Radon concentration and its associated health effects in the dwellings of fiche Selale North Shewa, Ethiopia. *Journal of Natural Sciences Research*. 2017;7(1):105-115.
- Ojo TJ, Ajayi I. Outdoor radon concentration in the township of Ado-Ekiti Nigeria. *Journal Of Atmospheric Pollution*. 2015;3(1):8-21.
- Fayaz K, Nawab A, Ehsan UK, Nimat UK, Iftikhar AR, Muzahir AB, Muhammad UR. Measurement of indoor radon concentration and the associated health risk in the five districts of Hazara division, Pankistan. *Journal of Environ. Monit*. 2012;(14 ):3015–3023.
- Germana W Mlay, Ismael N Makundi. Assessment of indoor radon – 222 concentrations in the vicinity of manyoni

- Uranium deposit, Singida. Tanzania C; 2018.
19. ICRP. recommendations of the international commission on radiological protection ICRP Publication 60 Ann. Icrp. 1991;(21):1-3.
  20. ICRP. Protection against radon-222 at home and at work international commission on radiological protection ICRP Publication 65; Ann Icrp. 1993;(23):2.
  21. Yeung AWK. Oral and maxillofacial radiology, applied oral sciences, faculty of dentistry, the university of Hong Kong, Hong Kong, Pr China, Radioprotection. 2019;54(2):103-109.
  22. Kent W Scheller, William S Elliott Jr. Geochemical and ray characterization of pennsylvanian black shales: Implications for elevated home. Kent, W. S. and William S. E. (2015): Geochemical and Ray Characterization of Pennsylvanian Black Shales: Implications for Elevated Home Radon Levels In Vanderburgh County, Indiana. Journal Of Environmental Radioactivity. 2015;(148):154-162.
  23. ICRP. Lung cancer risk from radon and progeny and statement on radon. ICRP Publication 115. Annal ICRP. 2010;40(1).

---

© 2021 Briggs et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<http://www.sdiarticle4.com/review-history/67820>