



# A Comparative Study on the Haematological and Biochemical Alterations in Catfish (*Clarias gariepinus*) under Natural and Artificial Environments in Sokoto Metropolis, Nigeria

Y. S. Baraya <sup>a\*</sup> and A. Abdulrazak <sup>a</sup>

<sup>a</sup> Department of Veterinary Pathology, Usmanu Danfodiyo University, Sokoto, Nigeria.

## Authors' contributions

This work was carried out in collaboration between both authors. Author YSB conceived the idea and designed the study, performed the statistical analysis and wrote the original draft of the manuscript. Author AA performed the literature search and carried out the investigation. Both authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJBCRR/2022/v31i10784

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/93287>

Original Research Article

Received 01/09/2022  
Accepted 03/11/2022  
Published 14/11/2022

## ABSTRACT

**Aims:** The study was conducted to compare the alteration in haematological and biochemical parameters in African catfish (*Clarias gariepinus*) under natural (river) and artificial (ponds) habitats in Sokoto metropolis, Nigeria.

**Study Design:** Whole blood and serum samples harvested from each group of the catfish were used for assessment of the haematological and biochemical parameters, respectively. Data were

\*Corresponding author: E-mail: ysbgobir123@gmail.com;

analyzed to compare changes between the two groups of African catfishes using *T*-test and  $P < 0.05$  was considered statistically significant.

**Place and Duration of Study:** Catfish samples were collected over a period of four weeks from Kwalkwalawa river and commercial fish farms in Sokoto metropolis. Samples for haematological parameters were determined at the Clinical Pathology Laboratory, Veterinary Teaching Hospital, Usmanu Danfodiyo University Sokoto. The serum analyses were performed at the Chemical Pathology Laboratory, Usmanu Danfodiyo University Teaching Hospital Sokoto.

**Methodology:** A total of fifty (50) whole blood samples were obtained from live African catfishes representing twenty-five (25) samples each from natural and artificial environments.

**Results:** The results showed significant changes in some haematological parameters, packed cell volume ( $P = 0.039$ ), haemoglobin concentration ( $P = 0.001$ ), white blood cell count ( $P = 0.012$ ), and differential neutrophils ( $P = 0.001$ ) and lymphocytes ( $P = 0.001$ ) counts between the two groups of African catfishes studied. Significant biochemical alterations in the total protein ( $P = 0.001$ ), total bilirubin ( $P = 0.028$ ) and direct bilirubin ( $P = 0.019$ ) were observed in catfishes under natural and artificial environments.

**Conclusion:** This study shows that aquaculture environment has great influence on the haematological and biochemical parameters in African catfish. These changes were more evident in catfish reared under the natural environment. Hence, the alterations in haemato-biochemical parameters were suggestive of unhealthy state or chronic stressful condition in catfish perhaps due to natural and human caused stresses.

*Keywords:* African catfish; biochemistry; haematology; ponds; River; Sokoto; Nigeria.

## 1. INTRODUCTION

African mud catfish is called *Clarias gariepinus*, is one of the most commonly cultured fish species in tropical Africa including Nigeria [1]. It belongs to the family Claridae which exists in the wild and artificially cultured in the ponds, cages, and pens for commercial purposes. Catfish is an omnivorous creature with a preference for a planktonic diet and other types of foods such as insects, insect larvae, pupae, fish, fish remnants and commercially available pelleted feeds [2,3]. People's attraction to catfish farming is linked to some imperative features of high-quality meat, enduring in nature, high tolerance to adverse conditions and production performance [1]. Economically, catfish has significant market values in Sokoto state and Nigeria at large.

Fish leaves favorably in aquatic habitat, and their blood profile is an important tool in the development of aquaculture system [4]. Fish blood can inform measurable physiological changes more quickly than any other physiological assessment parameters [5]. Changes in the value of a component of a blood parameter in comparison with the reference values are used in the interpretation of health status of animals [6]. Fish haematology has been used as an index to detect different stressful conditions such as exposure to pollutants, diseases, heavy metals, hypoxia and anaemia [7,8]. Intensive system of fish management has been characterized by high tendency of disease

outbreaks. Diseases are part of the limiting factors that reduces fish production [9]. Therefore, haematological evaluation is recommended in fish culture for early detection of diseases and environmental research [8,5]. Similarly, the examination of serum constituents has produced valuable information for detection and diagnosis of metabolic disturbances and diseases in fish [7]. In particular, a number of liver injury biomarkers have often been used to detect unhealthy state in fish [10].

Despite the nutritional and economic benefits of African catfish, yet there is paucity of information on their blood profile under different aquaculture systems. Thus, this study was aimed at investigating the changes in haematological and biochemical parameters of African catfish under natural and artificial environments in Sokoto metropolis Nigeria.

## 2. MATERIALS AND METHODS

### 2.1 Study Area

African catfishes were purchased at the Kwalkalawa river Sokoto and some commercial fish farms in Sokoto metropolis Nigeria, representing natural and artificial fish habitats. Sokoto, the capital city of Sokoto state is located at the northwestern part of Nigeria, longitudes 11° 30' to 13° 50' East and latitudes 4° to 6° 40' North. The state shares boundary with Niger

Republic to the north, Kebbi state to the southwest and Zamfara state to the southeast [11,12].

## 2.2 Fish Samples

A total number of fifty (50) *C. gariepinus* were collected alive at intervals for 4 weeks using cast net, it was immediately transferred into plastic container for transportation to the laboratory for samples collection. Twenty-five (25) each from the wild (Kwarkwalawa River) and cultured (commercial farms) aquatic environments in Sokoto metropolis. Adult and juvenile catfishes of both sexes were used for the study. Catfishes were weighed before collection of blood samples and then individually restrained manually.

## 2.3 Blood Collection

Blood samples were collected from live *C. gariepinus* (80 - 330g body weight) by puncturing the posterior caudal vein [13] using a heparinized plastic syringe fitted with a 21-gauge hypodermic needle. The samples were immediately transferred into ethylene diamine tetra-acetic acid containing tubes to avoid coagulation of the blood and used for haematological analyses. Serum samples were harvested in plain containers after the whole blood was clotted, centrifuged using bench centrifuge at 10,000rpm for 5 minutes and kept at freezing point until required for biochemical analyses.

## 2.4 Haematological and Biochemical Analyses

Haematological parameters, red blood cell (RBC), hemoglobin (Hb) concentration, packed cell volume (PCV), white blood cell (WBC), and differential leukocyte counts, namely lymphocytes, neutrophils, monocytes, eosinophils, basophils and band cells were determined at the Clinical Pathology Laboratory, Veterinary Teaching Hospital, Usmanu Danfodiyo University Sokoto. Serum total protein (TP), total bilirubin (TBIL), direct bilirubin (DBIL), aspartate aminotransferase (AST), alanine aminotransferase (ALT) and alkaline phosphatase (ALP) concentrations were determined according to the standard laboratory procedures at the Chemical Pathology Laboratory, Usmanu Danfodiyo University Teaching Hospital Sokoto.

## 2.5 Data Analysis

Data were analyzed using statistical package for social sciences version 22. *T*-test was used to

compare the haematological and biochemical changes between the African catfish reared under natural and artificial environments in Sokoto metropolis.  $P < 0.05$  was considered statistically significant.

## 3. RESULTS AND DISCUSSION

Table 1 shows the mean haematological parameters of *C. gariepinus* reared under natural and artificial environments. The mean PCV, Hb, WBC, neutrophils and lymphocytes values were significant ( $p < 0.05$ ) between wild and cultured catfishes. However, the mean RBC, monocytes, eosinophils, basophils and band cells were not significant ( $p > 0.05$ ) between the two groups. Based on the result, it can be seen that the mean PCV values of catfish from the two research locations were within the normal range of 30.8 - 45.5% [14]. But, Hb concentrations were below the reference range of 10 -14g/dL. Normally, Hb binds to oxygen in order to produce energy. As such, low protein diet could cause the Hb content to be low as well, and inconsequence will predispose fish to an infection [14].

Red blood cell count was not remarkable among the two groups studied. The results were within normal values [15]. Low erythrocyte counts indicate anaemia, while high levels indicate stress induced polycythemia in fish [16]. Anaemia has an impact on growth performance of fish due to the low RBC counts, and by extension reduced food supply to cells, tissues and organs [14]. The WBC counts in both groups of catfishes, their values were below total range of leukocytes in health, suggesting leukopenia perhaps due to unhealthy state or chronic stressful conditions [17,18]. WBC is the main component of the body's defence system [16]. Mean differential leukocyte counts indicate neutropenia and lymphocytosis [19,20] in both natural and artificially managed catfishes, thus suggesting extravasation of neutrophils and immunogenic stimulation, respectively [21].

Table 2 shows some biochemical parameters of *C. gariepinus* reared under natural and artificial environments. The mean TP, TBIL and DBIL were significant among wild and cultured catfishes. However, the mean AST, ALT, ALP and ALB were not significant between the two groups. More importantly, serum proteins concentrations were showing significant hypoproteinaemia especially in the wild catfish. Under situations of stress, many organisms mobilize proteins as an energy source via

**Table 1. Haematological parameters (Mean ± SD) of *Clarias gariepinus* reared under natural and artificial environments**

Parameters	Wild catfish	Cultured catfish
PCV (%)	37.08 ± 11.26	31.96 ± 3.77*
RBC (x10 <sup>6</sup> cells/mm <sup>3</sup> )	1.68 ± 0.96	2.07 ± 0.49
Hemoglobin (g/dL)	3.52 ± 1.82	7.06 ± 1.98*
WBC (x10 <sup>3</sup> cells/mm <sup>3</sup> )	8.34 ± 3.59	11.64 ± 5.13*
Neutrophils (%)	11.32 ± 4.71	18.58 ± 7.42*
Lymphocyte (%)	84.64 ± 6.04	75.33 ± 8.06*
Monocyte (%)	3.88 ± 3.23	5.45 ± 3.22
Eosinophil (%)	0.00 ± 0.00	0.33 ± 1.09
Basophils (%)	0.00 ± 0.00	0.00 ± 0.00
Band cells (%)	0.16 ± 0.62	0.29 ± 0.75

Key; SD = standard deviation. Results were determined using student's T-test.  $P < 0.05$  was statistically significant. \* =  $P < 0.05$

**Table 2. Biochemical parameters (Mean ± SD) of *Clarias gariepinus* reared under natural and artificial environments**

Parameters	Wild catfish	Cultured catfish
AST (IU/L)	7.48 ± 0.59	7.6 ± 0.71
ALT (IU/L)	5.00 ± 0.82	4.84 ± 0.85
ALP (IU/L)	74.24 ± 5.11	77.49 ± 17.36
TP (g/dL)	4.16 ± 0.37	4.96 ± 0.36*
ALB (g/dL)	2.84 ± 0.07	3.09 ± 0.14
TBIL (mg/dL)	0.71 ± 0.21	1.07 ± 0.75*
DBIL (mg/dL)	0.26 ± 0.06	0.52 ± 0.51*

Results were determined using student's T-test.  $P < 0.05$  was statistically significant. \* =  $P < 0.05$

oxidation of amino acids. Hence, the observed hypoproteinaemia could be linked to stress [22] because of exposure to environmental pollutants, diseases or attack by pathogens including haemoparasites.

Furthermore, the decrease in the TP content is related to the decrease in protein synthesis and absorption or protein loss via haemodilution as well the increase in energy demand by the fish to cope with environmental condition [23]. Although, liver transaminases and ALP activities were not remarkable, as such the results suggest that there was no significant hepatic tissue damage, enzymes synthesis and metabolism in the catfish studied [24]. Overall, variations in the haematological and serum biochemistry in fish may occur depending on the time of the day and season, physiological status, and presence of disease and/or pollutant exposure [25]. Similarly, dietary supplementation with pelleted feed could interfere with the haematological profile, intestinal microbiota and immune system [26].

#### 4. CONCLUSION

The study established that aquaculture environment has great influence on the haematological and biochemical parameters in African catfish. These changes were more apparent in catfish reared under the natural habitat. In conclusion, the haemato-biochemical alterations observed were suggestive of unhealthy state or chronic stressful condition in catfish examined. Therefore, further research should be conducted to identify the specific stress factors due to natural and human causes in the study area. Future studies could also explore the situation in other fish species in order to have a wide view of aquatic conditions and proper appropriate measures to address the problems identified.

#### ACKNOWLEDGEMENTS

We would like to thank the staff members at the Clinical Pathology Laboratory, Department of Veterinary Pathology, Faculty of Veterinary

Medicine, Usmanu Danfodiyo University, Sokoto Nigeria for their technical assistance.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Ibrahim O, Adetola J, Kolawole AE, Yusuf AA. Natural occurrence of *Diplostomum spp.* in farm raised African catfish (*Clarias gariepinus*) from Oyo state, Nigeria. Int J Vet Sci Med. 2016;4(2):41-45.
2. Tekle-Giorgis Y, Wagaw S, Dadebo E. The food and feeding habits of the African catfish, *Clarias gariepinus* (Burchell, 1822) (Pisces: Clariidae) in lake Hawassa and Shallo swamp, Ethiopia. Ethiop J Biol Sci. 2016;15(1).
3. Adewumi AA, Olaleye VF. Catfish culture In Nigeria. Progress, prospects and problems. Afr. J Agric Res. 2011;6(6): 1281-1285.
4. O'Neal CC, Weirich CR. Effects of low-level salinity on production and haematological parameters of channel catfish, *Ictalurus punctatus* reared in multicrop ponds. In: Book of abstract. Aquaculture 2001. International Triennial Conference of World Aquaculture Society. Disney Colorado Springs Resort Lake Buena Vista, Florida. 2001;484.
5. Ezeri GNO, Gabriel UU, Opabunmi OO. Hematological response of cultured and wild *Clarias gariepinus* to acclimation. Environ Ecol. 2001;3:628-632.
6. Etim NN, Williams ME, Akpabio U, Offiong EEA. Haematological parameters and factors affecting their values. Agric Sci. 2014;2(1):37-47.
7. Jamalzadeh HR, Keyvan A, Ghomi MR, Gherardi F. Comparison of blood indices in healthy and fungal infected Caspian salmon (*Salmo trutta caspius*). Afr J Biotechnol. 2009;8(2):319-322.
8. Yanuhar U, Raharjo DKWP, Caesar, NR, Junirahma NS. Haematology response of catfish (*Clarias sp.*) as an indicator of fish health in Tuban Regency IOP Conference Series: Earth Environmental. Science. 2021;718:012059.
9. Food and Agriculture Organization (FAO). Towards safe and effective use of chemicals in coastal aquaculture. GESAMP Reports and Studies No. 1997;65. Rome.
10. Soulivongsa L, Tengjaroenkul B, Patawang I, Neeratanaphan L. Cytogenetic, serum liver enzymes and liver cell pathology of the *Hampala* Barb fish (*Hampala macrolepidota*) affected by toxic elements in the contaminated Nam Kok River near the Sepon Gold Copper Mine, Lao PDR. IJERPH. 2021;18:5854.
11. Sokoto BA. The study of local geography for Nigerian secondary schools (1<sup>st</sup> edition). Ministry of Education, Sokoto. 2001;104-110.
12. National population commission (NPC) Nigeria Census Data; 2006.
13. Kori-Siakpere O, Ake JEG, Idoge, E. Haematological characteristics of the African snakehead, *Parachanna obscura*. Afr J Biotechnol. 2022;4:527-530.
14. Alamanda IE, Handajani NS, Budiharjo A. The use of haematology method and blood endoparasite observation for determining catfish (*Clarias gariepinus*) health in fishery Mangkubumen, Boyolali. Biodiversitas. 2007;834-8.
15. Lukistiyowati I, Berk SI. Identification of lactic acid bacteria (bal) from biofloc from technology with tapioca carbon sources in red tilapia in cultivation media (*Oreochromis sp.*). Perikan Terubuk. 2019;45:23-43.
16. Ikun F, Salosso Y, dan Toruan LN. Prosiding Seminar Nasional Kelautan dan Perikanan I (Universitas Nusa Cendana). 2013;112-23.
17. Tort L. Stress and immune modulation in fish. DCI. 2011;35(12):1366-1375.
18. Adeyomo BT, Obande RA, Solomon SG. Haematological reference ranges of cultured *Clarias gariepinus* in the Lower Benue River Basin, Nigeria. Comp Clin Path. 2012;21(5).
19. Awolabi OD. Haematological and serum biochemical profile of the upside-down catfish, *Synodontis membranacea* Geoffroy Saint Hilaire from Jebba Lake, Nigeria. Comp Clin Path. 2011;20:163-172.
20. Adedeji OB, Adegbile AF. Comparative haematological parameters of the bagrid catfish (*Chrysichthys nigrodigitatus*) and the African catfish (*Clarias gariepinus*) from Asejire dam in Southwestern Nigeria. JASR. 2011;7(7):1042-1046.
21. Clauss TM, Dove ADM, Arnold JE. Hematologic disorders of fish. Vet

- Clin Exot Anim Pract. 2008;11:445–462.
22. Aita NAA. Genotoxic, hematological and biochemical changes induced by phenol exposure in African Catfish (*Clarias gariepinus*). Glob Vet. 2014;13(3):316-324.
23. Singh N, Samartha M. Alteration in hematological and biochemical parameters in fresh water catfish *Clarias batrachus* as compared to *Cyprinus carpio*. IJFAS. 2022;12(1):1-14.
24. Evelyn EN, Bebe Y, Chukwuemeka N. Changes in liver and plasma enzymes of *Clarias gariepinus* exposed to sublethal concentration of diesel. Afr J Biotechnol. 2013;12(4):414-418.
25. De Pedro N, Guijarro AE, Lopez-Patino MA, Marinez-Alvarez R, Delgado M. Daily and seasonal variation in hematological and blood biochemical parameters in touch, *Tinca yince*. Aquaculture. 2005; 36:85-96.
26. Neepa NN, Islam MR, Mely SS, Mrong CE, Kole K, Sultana N, Mostakim GM. Improvement of growth and hematological profile of juvenile silver barb (*Barbonymus gonionotus*) through dietary probiotic (*Bacillus sp.*) supplementation. J Surv Fish Sci. 2022;8(2):11-25.

© 2022 Baraya and Abdulrazak; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/93287>