



Phytochemical Profile, Antioxidant and Antibacterial Activities of *Ephedra alata* Extract against Pathogenic Bacteria

Djahra Ali Boutlelis ^{a*}, Benkaddour Mounia ^a,
Benkherara Salah ^b, Shaib Inass ^a, Laib Ibtissam ^a,
Benine Chaima ^a and Bdida Soumaya ^a

^a Laboratory of Biology, Environment and Health, Faculty of Natural and Life Sciences,
University of El Oued, Algeria.

^b Department of Biology, Faculty of Natural and Life Sciences and Earth Sciences,
University of Ghardaia, Algeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors participated in the literature searches, materials and methods, discussion, read and approved the final manuscript.

Article Information

DOI: 10.9734/JOCAMR/2022/v19i3392

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/93258>

Original Research Article

Received: 07/09/2022

Accepted: 10/11/2022

Published: 14/11/2022

ABSTRACT

Plants have been a source of important chemical compounds that can be used for treatment of different disease. Natural products in plants also are a source of active compounds and can be considered an alternative to synthetic molecules with high antioxidant potential. Objective of this study was to determine the antioxidant and antibacterial potential of *Ephedra alata* extract. Methods: Phytochemical values namely, total phenolic, DPPH radical scavenging effect were studied by established methods. Antimicrobial properties of *Ephedra alata* was tested using cup-plate diffusion method and disc diffusion.

*Corresponding author: E-mail: djahra_ab@yahoo.fr;

Results showed numerous secondary metabolic products including in plant *Ephedra alata* extract such as Flavonoids, Saponins, Tannins were detected. And other metabolic products such as Steroids and Cardinolids are absent. The yield for the aqueous extract was 9.89%. However, the polyphenols extract content was equal to 0.004243 (mg GAE/ml extract. Effectiveness of antioxidant plant using DPPH free root was estimated at $IC_{50} = 0.901$ mg/ml for aqueous extract. In antibacterial screening, the extract showed antibacterial power with zone of inhibitions compared to positive controls Penicillin against multi-resistant germs such as: *Pseudomonase aeruginosa* and *Serratia fonticola* responsible of many infectious. Growth inhibition varies with bacterial species and concentration of *Ephedra alta* extract.

Keywords: Antioxidant; antibacterial activity; *Ephedra alata*; penicillin; multi-resistant germs.

1. INTRODUCTION

Due to their powerful pharmacological activity, user-friendliness, economic viability, and low toxicity, numerous plants have been the subject of increasing research in the past few decades [1]. The discovery of natural antioxidants derived from plant materials has increased recently in order to replace synthetic antioxidants because the former are recognized as green medicine and are safe for the management of health [2]. Natural antioxidants from higher plants, especially common chemicals like vitamins, carotenoids, and phenolics, have antioxidant activity and lower chronic health issues linked to disease [3].

Reactive oxygen species (EROs) are implicated in physiological processes at low levels. However, the primary cellular components may become poisonous if EROs synthesis is too high; lipids, proteins and nucleic acids, and cause oxidative stress which will be implicated in various pathologies such as cancers, diabetes, Inflammatory diseases... [4]. In recent years, multiple drug resistance in human pathogenic microorganisms has developed due to indiscriminate use of commercial antimicrobial drugs commonly used in the treatment of infectious diseases [5]. In addition, the fight against bacterial infections becomes complicated because many bacteria have developed resistance to most antibiotics, which has been a major world health challenge [6].

However, It is concerning that synthetic compounds have negative consequences when used to treat bacterial infections and oxidative stress. Also finding a replacement for the usage of chemical antioxidants and traditional antibiotics seems to be crucial. Herbal therapies are a viable alternative to conventional primary care systems, making them a promising way to the development of conventionally enhanced

medications [7]. The abundance of natural substances found in medicinal plants, such as polyphenols, flavonoids, tannins, etc., that have antioxidant and antibacterial properties, has recently piqued the interest of many researchers.

Scientific research's top priorities now include finding novel bioactive compounds from natural sources that have few or no adverse effects and using them as therapeutic substitutes for manufactured molecules [8]. For this study, we investigated phytochemical compounds, antioxidant and antibacterial activity of *Ephedra alata*, an herb widely used in south Algerian traditional medicine.

2. MATERIALS AND METHODS

2.1 Materials

Plant material: Aerial parts of *Ephedra alata* were collected from Alenda village, Wilaya of El-Oued (South-east Algeria) and were identified by the botanist Prof. Atef Chouikh (Faculty of Natural and Life Sciences, University of El Oued, Algeria), were properly cleaned by being washed under running water to remove dust and other foreign objects, and then shade dried for 15 days.

Bacterial strains: The microbiological material consists of five bacterial strains. These are Gram-positive and Gram-negative bacteria and pathogenic to humans, often multi-resistant to antibiotics and causing serious infections. They come from the El Majed laboratory (Private laboratory) located in the wilaya of El-Oued (Algeria). The bacterial strains are: *Escherichia coli*, *Pseudomounase aeruginosa*, *Staphylococcus aureus*, *Proteus mirabilis* and *Serratia fonticola*.

Antiibotic: To compare the antibacterial effects of natural products isolated in the aqueous

extract with those antibiotics. Penicillin was used in (10 mg) [discs with a diameter of 6 mm contain 10 mg of the antibiotic].

2.2 Methods Followed

Preparation of plant extract: 400 g of shade-dried powdered aerial parts was extracted by using 4 L of water. The resultant extract was filtered using Whatman Filter paper (N^o:1) and evaporated at low pressure, thus dried. 13% w/w was the extract yield percentage of the initial raw material and stored in refrigerator at 4°C for further use.

Phytochemical qualitative analysis: Phytochemical analysis of the plant extract was evaluated using standard procedures [9-12]. Qualitative analysis were done to find the presence of the biochemical constituents such as Alkaloids, Flavonoids, Anthocyanins, Saponins, Terpenoids and steroids, Leucoanthocyanins, Cardinols, Gallic tannins and Catechic tannins.

Determination of total phenolic content: Total phenolic content were determined according according to method of Singleton & Rossi [13]. Briefly, 200 µl of Folin-Ciocalteu reagent added to 100 µl of the diluted sample. A saturated sodium carbonate solution (Approximately 20%) in 600 µl was added after 3 minutes. The absorbance of the sample was measured at 765 nm after 2 hours of incubation at room temperature. The standard curve was produced by applying the same process repeatedly to all standard Gallic acid solutions (0 - 100 µg/ml).

DPPH free radical scavenging assay: Of DPPH, 2.4 mg was dissolved in 100 ml of methanol to prepare the 1,1-diphenyl-2-picrylhydrazyl solution. Of extract, 50 µl (or

ascorbic acid as a control) was added to 1.950 ml of the DPPH solution previously produced. The reaction mixture is quickly agitated, then maintained at room temperature for 30 minutes in the dark to complete the reaction. The reaction medium's absorbance is measured at 515 nm [14].

Antibacterial activity: The technique of diffusion on solid medium has been used [15]. It is a method similar to that of antibiogram which consists in determining the sensitivity of a bacterial strain to one or more products. A sterile disc of Whatman paper (n^o1), the test items are submerged in a 6 mm diameter. Pure extract, Extract diluted to 1/2 and Extract diluted to 1/4. The three soaked discs of the three concentrations are placed on agar which is previously inoculated with strains and an inoculum concentration between 10⁶ and 10⁸ CFU/ml. Petri dishes are incubated at 37°C for 18 to 24 hours. After incubation, if the product is toxic to the bacterial species, it forms a halo or zone around the disc. The diameter of the inhibition zones is measured in millimeter's (mm). Control discs (sterile distilled water) and comparison discs (Penicillin 10 mg) are included in the tests. Each test of the antibacterial activity of the extract is performed in triplicate.

3. RESULTS

3.1 Phytochemical Qualitative Analysis

To determine if chemical groups were present in the studied plant extract, phytochemical screening was performed (Table 1), for Understand the active components responsible for biological actions.

Table 1. Phytochemical qualitative analysis of *Ephedra alata* aerial parts extract

Chemical constituents	Aerial parts of <i>Ephedra alata</i>
Alkaloids	(+)
Flavonoids	(+)
Anthocyanins	(+)
Saponins	(+)
Terpenoids	(+)
Leucoanthocyanins	(+)
Tannins	(+)
Steroids	(-)
Cardinols	(-)

(-) Absence of phytochemicals compounds.

(+) Presence of phytochemicals compounds

3.2 Determination of Total Phenolic Content

The phenolic content obtained from aqueous extracts was estimated using a calibration curve and a reference extract (Gallic acid) at different concentrations. Results are expressed in mg Gallic acid equivalent per mg extract (mg GAE/mg extract). The calibration curve is established with a correlation $y = 0.0009x + 0.001$ and coefficient $R^2 = 0.9926$. The results were represented in Table 2.

3.3 DPPH Free Radical Scavenging Assay

Ephedra alata aerial parts extract was tested for antioxidant activity by using DPPH free radical scavenging assay. The results of antioxidant activity were summarized in Fig. 1, and DPPH free radical scavenging of plant extract was compared to that of the standard (Ascorbic acid).

3.4 Antibacterial Activity

The results of the different tests performed with five strains of bacteria used for the products tested are grouped in Table 3. The results show that there is a great heterogeneity in the results obtained, the antibacterial action is more or less important in accordance with the nature of the strain. Better resistant registries for both bacterial species: *Staphylococcus aureus* and *Proteus mirabilis*.

The natural products tested of *Ephedra alata* still manage to limit their development on ranges between 5.83 and 5.67 mm for *Pseudomonas aeruginosa*. This effect is often greater than that recorded in the presence of the antibiotic (Penicillin). *Staphylococcus aureus*, although considered as a highly resistant species to most antibiotics, it has been shown to be sensitive to the effect of Penicillin.

Table 2. Quantitative analysis of total phenolic of *Ephedra alata* aerial parts extract

Aerial parts of <i>Ephedra alata</i>	
Total phenolic (mg GAE/mg extract)	13.72 ± 0.02




Fig. 1. Antioxidant activity by using DPPH of *Ephedra alata* aerial parts extract

Table 3. Inhibition zones (mm) of antibacterial activity for *Ephedra alata* aerial parts extract

Strains	Tested products				
	Pure Ext	Ext diluted to 1/2	Ext diluted to 1/4	Water	Penicillin 10 mg
<i>Escherichia coli</i>	1.12±0.32	0.89±0.98	0.56±0.37	00±00	06.15 ±1.22
<i>Pseudomounase aeruginosa</i>	5.83±1.53	5.67±0.74	4.33±0.53	00±00	00 ±00
<i>Staphylococcus aureus</i>	00±00	00±00	00±00	00±00	10 ±0.83
<i>Proteus mirabilis</i>	00±00	00±00	00±00	00±00	05.03 ±1.22
<i>Serratia fonticola</i>	6.33±1.86	4.67±2.24	4.33±1.56	00±00	10 ±2.19

4. DISCUSSION

The phytochemical content varies depending on the product concerned. These levels may be associated with several factors such as growth and maturity conditions, genotype, storage conditions and extraction methods [15,16]. In this case, these factors resulted in a high level of plant extract (13%) for aerial parts of *Ephedra alata*. Moreover, previous studies show that methanol and water are the most widely used solvents for high recovery of phenolic compounds [17-19].

The results of the phytochemical analysis carried out on the aqueous extracts of the aerial part of *Ephedra alata* showed the presence of certain active compounds Tannins, Flavonoids, Anthocyanins, Saponins, Alkaloids, Terpenoids and Leucoanthocyanins when sterols and Cardiolids are absent, these results are compatible with the literature [20-22]

Phenolic compounds are considered important contributors to the biological activities of plants [23]. These compounds possess various biological activities such as Antibacterial [24], Antifungal [25], Antidiabetic [26], Hepatoprotective [27], Haemato-protective [28]. For this reason, total phenolic content of *Ephedra alata* aerial parts were determined in this study. The results show that total phenolic content are abundant in the aerial part of *Ephedra alata* (13.72 ± 0.02 mg GAE/mg extract). This may be related to the harsh climatic conditions of the places where they grow (high temperature, high sun exposure, drought and salinity) that stimulate the biosynthesis of secondary metabolites such as polyphenols [29].

Antioxidant potential of polyphenols is great interest due to its chemo-protective against degenerative diseases such as neurological and cardiovascular diseases and its inhibitory effect of lipid peroxidation of foodstuffs [30]. Oxidative processes are multiple and the nature of antioxidant activity can be multifaceted and attributed to different mechanisms like the trapping of free radicals, chelation of transition metal ions, initiation prevention of EROs chain producing reactions and decomposition of peroxides. [31] Thus, the combination of several complementary antioxidant tests is useful in assessing the antioxidant potential of extracts [32]. The DPPH radical method used in this study. It's a common process for the antioxidant activity of the sample under investigation is

estimated by the degree of discoloration of the DPPH solution [33].

Our results indicate that the aqueous extract has a high activity against scavenging assay of free radical DPPH with an $IC_{50}=0.901$ mg/ml. In a similar study on ethanol extracts from *Ephedra procera* (same genus of our plant species) the authors reported an IC_{50} level of 0.056 mg/ml [34], a very low level, indicating an important antioxidant activity compared to ours, because the reductive power of a compound can serve as a significant indicator of its potential antioxidant activity, even that there is a direct correlation between antioxidant activities and the reduction power of the components of some plants [35,36]. In addition, aqueous extracts are assumed to contain polar compounds such as flavonoids and glycosides that may be responsible for their antioxidant activity [37].

Results found of antibacterial activity confirm once again the efficacy of the extracts of medicinal plants and their antiseptic power which rivals that of antibiotics [38,39]. Numerous studies have demonstrated the antibacterial effect of natural active substances. Indeed, an article in the journal of life sciences [40]. reports that natural products from *Ephedra foliate* extracts has inhibitory and lethal effects on a variety of strains. For strains tested, only *Staphylococcus aureus* and *Proteus mirabilis* showed a high performance in resistance to the natural products tested and all concentrations used. The crude extract of many plants have also been shown in several investigations to have some antibacterial action. While these plants extracts and essential oils severely restrict the development of strains, [41]. According to Candan et al. (2003) [42], hydrosoluble substances have a lower effect compared to no hydrosoluble substances. This probably refers to the ability of fat-soluble molecules for inserting into the bacterial membranes cells and damages them.

The constituents of the phenolic structure are very active against microorganisms and act as denaturants of proteins. These phenolic compounds are capable of binding to certain proteins and enzymes, thereby modifying the enzymatic equilibrium [43]. Finally, the activity of an extract is probably due to the presence of synergy between a number of components, which, when separated, become inactive individually [44].

5. CONCLUSION

Through the study of *Ephedra alata* phytochemical profile, antioxidant and antibacterial activity face pathogenic and multiresistant bacterial strains. It seems that these phytochemical compounds of *Ephedra alata* have an important antioxidant and antibacterial power on the tested germs responsible for infectious diseases.

CONSENT AND ETHICAL APPROVAL

It is not applicable.

ACKNOWLEDGEMENTS

This research was made by the D01N01UN390120230004. Research project, which was funded by the Algerian Ministry of Higher Education and the Directorate General for Scientific Research and Technological Development.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Chew AL, Jessica JJA, Sasidharan S. Antioxidant and antibacterial activity of different parts of *Leucas aspera*. *Asian Pac. J. Trop. Biomed.* 2012;2:176–180.
2. Chanwitheesuk A, Teerawutgulrag A, Rakariyatham N. Screening of antioxidant activity and antioxidant compounds of some edible plants of Thailand. *Food Chem.* 2005;92:491–497.
3. Duarte MCT, Figueira GM, Sartoratto A, Rehder VLG, Delarmelina C. Anti-candida activity of Brazilian medicinal plants. *J. Ethnopharmacol.* 2005;97:305–311.
4. Favier AE, Nève J, Faure P. Trace elements and free radicals in oxidative diseases, American Oil Chemists' Society Press, USA. 1994;1-3.
5. Janovská D, Kubiřková K, Kokosřka L. Screening for antimicrobial activity of some medicinal plants species of traditional Chinese medicine. *Czech J. Food Sci.* 2003;21:107–110.
6. Benbrinis S. Evaluation des activités antioxydante et antibactérienne des extraits de *Santolina chamaecyparissus*. These. Universty of Ferhat Abbas-SETIF. Algeria. 2012;143.
7. Djahra AB. Etude phytochimique et activité antimicrobienne, antioxydante, antihépatotoxique du Marrube blanc ou *Marrubium vulgare* L. These. University of El Oued. Algeria. 2014;114.
8. Pelt JM. Les nouveaux actifs naturels. Marabout. Paris. 2001;219-124.
9. Iyengar MA. Study of crude drugs. 8th Ed., Manipal Power Press, Manipal, India. 1995;2-5.
10. Iyengar MA. Study of crude drugs. 8th Ed. –Manipal Power Press, Manipal, India;1995.
11. Bouquet A. Plantes médicinales du Congo-Brazzaville: *Uvariopsis*, *Pauridiantha*, *Diospyros*. Travaux et Documents de l'ORSTOM, TOM 13. Paris. 1972;112.
12. Solfo RR. Etude d'une Plante Médicinale Malgache *Buxus madagascariensis* Bail et ses variétés. Travaux et Documents de l'ORSTOM.TOM 25. 1973;98.
13. Dahou N, Yamki K, Tahrouch S, Idrissi-Hassini LM, Gmira N. Screening phytochimique d'une endémique Ibéro Marocaine *Thymelaea lythroides*. Ed., Bull.Soc.Pharm. 2003;142:61–78.
14. Singleton VL, Rossi JA. Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *American Journal of Enology and Viticulture.* 1965;16(3):144–158.
15. Benhammou N, Atik Bekkara F, Kadifkova P. Antiradical capacity of the phenolic compounds of *Pistacia lentiscus* L. and *Pistacia atlantica* Desf, *Advances in Food Sciences.* 2007;29(3):155-161.
16. Djahra AB, Bordjiba O, Benkherara S. Extraction, séparation et activité antibactérienne des tanins de marrube blanc (*Marrubium vulgare* L.). *Phytothérapie.* 2013;11:348–352.
17. Zang D, Hamauru Y. Phenolic compounds, ascorbic acid, carotenoids and antioxidant properties of grebe. Red and yellow bell peppers. *Food Agric Environ.* 2003;2:22–7.
18. Sahreen S, Khan MR, Khan RA. Evaluation of antioxidant activities of various solvent extracts of *Carissa opaca* fruits. *Food Chemistry.* 2010;122:1205-1211.
19. Xia EQ, Deng GF, Guo YJ, Li HB. Biological activities of polyphenols from grapes. *International Journal of Molecular Sciences.* 2010;11:622-646.
20. Bouzid W, Yahia M, Abdeddaim M, Aberkane MC et Ayachi A. Evaluation de l'activité antioxydante et antimicrobienne

- des extraits de L'Aubepine Monogyne. Lebanese Science Journal. 2011;12(1): 59-69.
21. Amakura Y, Yoshimura M, Yamakami S, Yoshida T, Wakana D, Hyuga M, Hyuga S, Hanawa, T, Goda Y. Characterization of phenolic constituents from *Ephedra Herb* Extract, *Molecules*, 2013;18:5326-5334.3.
 22. Hussein SAM, Barakat HH, Nawar MAM, Willuhn G. Flavonoids from *Ephedra aphylla*. *Phytochemistry*. 1997;45(8):1529–1532.
 23. Nawwar M, El-Sissi HI, Barakat HH. Flavonoid constituents of *Ephedra alata*. *Phytochemistry*, 1984;23(12):2937-2939.
 24. Le K, Chiu F, Ng K. Identification and quantification of antioxidants in *Fructus lycii*. *Food Chemistry*. 2007;105:353-363.
 25. Djahra AB, Benkaddour M, Benkherara S. Evaluation of antimicrobial activity of medicinal plant *Cotula cinerea* against pathogenic strains. *PONTE International Scientific Researches Journal*. 2020;76(4):316-322.
 26. Djahra AB, Benkaddour M, Zeghib K, Benkherara Salah, Shaib I, Ghania A, Bbida S. Biofungicide activity of *Datura stramonium* leaf extract against phytopathogenic fungi. *International Journal of Biological and Agricultural Research*. 2019;2(1):1-5.
 27. Benkherara S, Bordjiba O, Harrat S, Djahra AB. Antidiabetic potential and chemical constituents of *Haloxylon scoparium* aerial part, an endemic plant from Southeastern Algeria. *International Journal of Secondary Metabolite*. 2021;8(4):398–413.
 28. Djahra AB, Benkaddour M, Benkherara S. Antioxidant and hepatoprotective potential of *Coriandrum sativum* L. against hepatic injury by lambda-cyhalothrin insecticide. *Journal of Drug Delivery and Therapeutics*. 2020;10(3-s):182-188.
 29. Zeghib K, Boutlelis DA, Menai S, Debouba M. Protective effect of *Atriplex halimus* extract against benzene-induced haematotoxicity in rats. *Ukr. Biochem. J*. 2021;93(4):66-76.
 30. Falleh H, Ksouri R, Chaieb K, Karray-Bourouai N, Trabelsi N, Boulaaba M, Abdelly C. Phenolic composition of *Cynara cardunculus* L. organs and their biological activities. *Comptes Rendus Biologies*. 2008;331:372-379.
 31. Bubonja-Sonje M, Giacometti J, Abram M. Antioxidant and antilisterial activity of olive oil, cocoa and rosemary extract polyphenols. *Food Chemistry*. 2011;127: 1821-1827.
 32. Ozen T. Investigation of antioxidant properties of *Nasturtium officinale* (Watercress) leaf extracts. *Acta Poloniae Pharmaceutica-Drug Research*. 2009;66(2):187-193.
 33. Ksouri R, Falleh H, Megdiche W, Trabelsi N, Mhamdi B, Chaieb K, Bakrouf A, Magné C, Abdelly C. Antioxidant and antimicrobial activities of the edible medicinal halophyte *Tamarix gallica* L. and related polyphenolic constituents. *Food and Chemical Toxicology*. 2009;47:2083-2091.
 34. Gulcin I, Huyut Z, Elmastas M, Aboul-Enein HY. Radical scavenging and antioxidant activity of tannic acid. *Arabian Journal of Chemistry*. 2010;3:43-53.
 35. Dehkordi NV, Kachouie MA, Pirbalouti AG, Maekpoor F, Rebeff M. Total phenolic content, antioxidant and antibacterial activities of the extract of *Ephedra procera* fisch. Et mey. *Acta Poloniae. Drug Research*. 2005;72(2):341-345.
 36. Jeong SM, Kim SY, Kim DR, Jo SC, Nam KC, Ahn DU, Lee SC. Effects of heat treatment on the antioxidant activity of extracts from citrus peels. *Journal of Agriculture and Food Chemistry*. 2004; 52:3389–3393.
 37. Yildirim A, Mavi A, Kara AA. Determination of antioxidant and antimicrobial activities of *Rumex crispus* L. extracts. *Journal of Agricultural and Food Chemistry*. 2001; 49:4083-4089.
 38. Giner Pons RM, Rios Canavate JL. *Santolina chamaecyparissus*: Especie mediterranea con potenciales aplicaciones terapeuticas en procesos inflamatorios y transtornos digestivos. *Revista de Fitoterapia*. 2000;1:27-34.
 39. Benkherara S, Bordjiba O et Djahra AB. Évaluation *In vitro* de l'activité antibactérienne de l'huile essentielle de *Salvia officinalis*. *Phytothérapie*. 2015;13: 14-18.
 40. Djahra AB, Bordjiba O, Benkherara S, Benkaddour M. Evaluation of Algerian spontaneous species: Phytochemical and biological study of aromatic and medicinal plant *Marrubium vulgare*. *PhytoChem & BioSub Journal*. 2015;9(1):2-9.
 41. Sharad B. Evaluation of antibacterial potential of *Ephedra foliate* boiss. ex. c. a. mey. *Journal of life sciences*. Jai Narain

- Vyas University, Jodhpur- 342 001 India. 2015;1172-2015.
42. Candan F, Unlu M, Tepe B, Daferera D, Polissiou M, Sökmen A, Akpulat HA. Antioxidant and antimicrobial activity of the essential oil and methanol extracts of *Achillea millefolium* subsp. *millefolium* Afan. (Asteraceae). Journal of Ethnopharmacology. 2003;87:215-220.
43. Ozawa T, Lilley TH, Haslam E. Polyphenol interactions: Astringency and the loss of astringency in ripening fruit. Photochemistry. 1987;26: 2937–42.
44. Sarker SD, Latif Z, Gray AI. Natural product isolation. In: Sarker SD, Latif Z, Gray AI. Natural products isolation. Humana Press (Totowa). 2005;1-23.

© 2022 Djahra et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/93258>