



Effect of Land use Change on Total Phosphorus and Its Fractions in North-Western Himalayas

Owais Bashir ^{a*}, Shabir Ahmad Bangroo ^a, Nasir Bashir Naikoo ^a,
Sandeep Kumar ^a, Rehana Rasool ^a, Aamir Hassan Mir ^a,
Sheikh Amjid ^a, Omer Reshi ^b, Roohi Jan ^c, Irfan gani ^d and Larieb Mir ^e

^a Division of Soil Science and Agricultural Chemistry, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, 190025, Kashmir, India.

^b Centre for Environment and Marine Studies, King Fahd University of Petroleum and Minerals, Saudi Arabia.

^c Division of Plant Pathology, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, 190025, Kashmir India.

^d Division of Floriculture, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, 190025, Kashmir, India.

^e Division of Soil Science, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, 190025, Kashmir, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i232535

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/93616>

Original Research Article

Received 06 September 2022

Accepted 12 November 2022

Published 16 November 2022

ABSTRACT

Aims: Conversion of land from forest to cropping has a serious effect on soil phosphorus and its fractions

Results: Land use is now widely understood to be a primary factor in environmental change across all time and space scales. The purpose of this research was to ascertain how different land uses affected the concentration of phosphorus in soil. Soil phosphorus (P) reserves are depleted when land is converted from natural vegetation to permanent agricultural cropping. The transformation of North-Western Himalayas from a forest-dominated to a grassland-dominated ecosystem is just one example of the diversity of land significantly less soil aggregation occurred when agricultural land was cleared of its native vegetation. Total organic carbon in soils was reduced when grassland was

*Corresponding author: E-mail: owaisbashir10@gmail.com;

converted to cropland. Reduced total organic carbon (TOC) concentrations by 62-79% and organic phosphorus (Po) concentrations by 47-53%. Even though, the total silt+clay fraction's contribution was negligible, it contained a significant amount of C and Po reserves and the C/Po ratio has been holding fairly steady, they have proven to be more robust. This impact of cropping on soil P reserves has been demonstrated in research, but changing land use practices can alleviate these problems significantly.

Keywords: Landuse; phosphorus fractions; pasture.

1. INTRODUCTION

Phosphorus (P) is a macronutrient that plants need to grow and flourish. However, it is believed that about 5.7 billion hectares (ha) of land around the world has insufficient amounts of plant-available P [1]. Phosphate rock is used in crop production at the current rate of about 150 million tons per year [2]. In addition, global reserves of P are the smallest of the 14 essential plant nutrients [3], and the vast majority of them are located in just one country (Morocco). Understanding P behavior in soils, including its effect on long-term agricultural production, is thus crucial. Despite total P concentrations between 200 and 800 mg kg⁻¹ soil [4], only 0.1% of this P is available for plant uptake in the soil solution [5]. Since plants uptake P from the soil solution, which has low P concentrations, that fraction must be replenished to meet plant needs. Labile P replenishes. The first labile pool (Pi) is P in mineral forms that dissolve, releasing P, or P adsorbing to soil surfaces. Plant and enzyme activity mineralizes the second labile pool (Po) [6].

Phosphorus in soil is dispersed among numerous geochemical phases, including soil solution and exchangeable phase, OM phase, Ca-bound phase, and Fe and Al-bound phases. The degree of P association with distinct geochemical forms greatly depends on soil physico-chemical characteristics owing to soil type, climate, and management approaches [7]. Under certain situations, these P fractions can be altered due to their different mobility, bioavailability, and chemical behaviors in soil. In soils, fractionation defines them qualitatively and quantitatively. Bioavailability, P leachability, and chemical form transitions in agricultural and contaminated soils can be predicted using such knowledge. Redistribution among fractions affects P mobility, unlike total P analysis. A high amount of P in labile pools indicates increased plant availability but also greater leaching or runoff [8].

Long-term cultivation and cropping reduce soil P concentrations in low-input agricultural systems [9, 10]. It was reported that both inorganic P and

Po concentrations reduced significantly. However, much remains unknown about cropping, along with how land management techniques affect P loss and how soil aggregates sustain and cycle P pools, particularly Po [7,11]. Only few studies have examined its effects on P forms. Fonte et al. [12]. Concluded that despite a 40% increased Po content in productive pasture soils, there wasn't any difference in total P content.

2. MATERIALS AND METHODS

2.1 Site Description

Study area: The study area is located in the temperate Himalaya of the North–Western part of India between the latitude 34°12' to 34°20' North latitude and 74°20' to 74°34' East longitude with an elevation of 1584 meters and an area of 3353 km². The region has the high, mid and low altitudes consisting of mountains, hills and valleys. The region has an annual rainfall of 1270 mm and an average temperature of 24° C. The highland is separated by a wide valley which is stretched by the river Jhelum in the eastern and western direction. In the north western part is the Jhelum and in the southern part is Pakistan. The topography of the area steep slopy to moderately slopy with some plain area as well.

2.2 Soil Sampling and Analysis


All the soil samples were air-dried, polished and pulverized using mortar and pestle and then sieved through a 2mm mesh sieve. The hydrometer method [13] was used to determine particle size distribution. Blake and Hartge's [14] method was used to estimate soil bulk density in which each core of soil sample was oven-dried at 105°C for at least two to three days. The soil pH was determined potentiometrically 1:2.5 (w/v) soil-water supernatant suspension [15]. SOC was estimated using [16]. Citrate-dithionite-extractable Fe and Al were determined using method of Kuo [17]. Concentrations of P within the soil samples were examined using Colwell-P

as well as sequential fractionation. For Colwell-P, concentrations were determined following extraction with NaHCO_3 [18], with P measured using the molybdate blue colorimetric method [19]. Sequential extraction of P was conducted following a modified Hedley fractionation method [20].

3. RESULTS

Overall, the pH ranged from 5.5 to 6.8 in the soils of different land uses (Table 1). Soil pH levels were typically lower in areas with forest compared to those in areas where the land had been converted to agriculture (Table 1). The EC

of the soils tested ranged from 0.01 to 0.21 dSm^{-1} , indicating no salinity hazard. A high percentage of clay was present in all soils, with values ranging from 25 % to 43 % (Fig. 1). Average concentration of SOC ranged from 3.2 to 10.3 g Kg^{-1} across all land uses with highest in forest soils and lowest in paddy soils. Total nitrogen varied between 0.4 to 7.7 g Kg^{-1} with highest in forest soils and lowest in paddy soils. Fe_d varied between the land uses from 1.2-15.5 g Kg^{-1} (Fig. 2) with lowest in paddy soils and highest in forest soils, similar trend was found in case of Al_d among all the land uses (Table 1).



Map 1. Gis map of the study area




Fig. 1. Mechanical analysis of different land uses

Table 1. Physico-chemical properties under different land uses

| Land use | pH | EC (dSm ⁻¹) | Bulk Density (Mgm ⁻³) | Sand (%) | Silt (%) | Clay (%) | SOC (g Kg ⁻¹) | Total Nitrogen (g Kg ⁻¹) | Fe _d (g Kg ⁻¹) | Al _d (g Kg ⁻¹) |
|----------|-----|-------------------------|-----------------------------------|----------|----------|----------|---------------------------|--------------------------------------|---------------------------------------|---------------------------------------|
| Forest | 5.5 | 0.07 | 1.15 | 20 | 15 | 25 | 10.3 | 7.7 | 13.5 | 8 |
| Pasture | 5.8 | 0.01 | 1.21 | 15 | 20 | 41 | 6.7 | 5.6 | 9.4 | 3.2 |
| Paddy | 6.8 | 0.15 | 1.42 | 24 | 25 | 30 | 3.2 | 0.4 | 1.2 | 2.1 |
| Orchard | 6.7 | 0.21 | 1.51 | 30 | 30 | 43 | 4.1 | 1.1 | 1.6 | 2.8 |

Table 2. P concentrations influenced by land use, including for total P, inorganic P (Pi), organic P (Po), and the Po as a proportion of total P

| Land use | Total P (mg Kg ⁻¹) | Total Pi (mg Kg ⁻¹) | Total Po (mg Kg ⁻¹) | Colwell-P (mg Kg ⁻¹) |
|----------|--------------------------------|---------------------------------|---------------------------------|----------------------------------|
| Forest | 975.3 | 536.2 | 439.1 | 65 |
| Pasture | 1193 | 665.5 | 527.5 | 53 |
| Paddy | 528.1 | 402.5 | 125.6 | 36 |
| Orchard | 760.8 | 485.5 | 275.3 | 42 |




Fig. 2. Chemical properties of different land uses




Fig. 3. Phosphorus fractions under different land uses

Total P ranged from 526 to 901 mg kg⁻¹ in studied soils. Pasture soils had the highest total P concentration of 901 mg kg⁻¹ (Table 2). 65-70 % of P was Pi and 25-50 % Po in all the land uses. Converting forest soils to agricultural production (cropping) affected these P fractions. First, transition of soil to cropping reduced total P concentrations. However, conversion of forest soils to other land uses decreased Po's contribution to total P. In all soils, the C/Po ratio (for Po) also decreased significantly for all land uses for all three soils. Finally, transition from forest to cropping reduced Colwell-P concentrations. These P fractions within the soil were influenced by the transition from grazing to cropping, though the extent of this influence varied across land uses. At first, it was discovered that when land was used for farming (cropping), total P concentrations fell. No matter the land use, it was obvious that conversion significantly reduced the contribution of Po to total P. Similar declines in C/Po ratio (for Po) were observed across all land uses. For any given shift in land use, bulk soil Pi/Po ratios rose, with cropping leading the way and plantation showing the least change. Furthermore, Colwell-P concentrations declined dramatically across all land types.

4. DISCUSSION

Changes in both total P and Pi were found to be highly site-specific. However, cropping consistently reduced Po compared to soil with forest soils Stutter et al. [21], who analyzed five Australian cropping soils, found that Po

decreased significantly for four of them, on average by 28%. Without fertilizer addition, Pi and Po did not change in the surface layer but showed depletion deeper within the soil profile. Total organic carbon (TOC) decreased by 45 percent after land was converted from forest to other land uses, but neither total P nor P fractionation were affected [22]. However, a shift from cropland to pasture led to a 94% increase in both total P and Po. Although Pi decreased by 31%, the total P and Po were barely impacted by the conversion of this pasture to other land uses (Fig. 2). Previous research has shown that after afforestation, total P and Po often decrease while plant-available Pi can increase [23].

Before being used for grazing, the soil in the pastures was cropped. Firstly, it was seen that the Po fraction was reduced by 63% after cropping, but then increased to almost the same levels as in the soil with native vegetation after being converted to pasture or plantation. Switching from grazing to cropping resulted in a significant drop in total organic carbon (85% in the bulk soils due to net mineralization). Total organic carbon (TOC) concentrations were higher in soil that had been converted from cropland to pasture, but not to pre-farm levels. Finally, related to the preceding points, we noticed that converting cropland to pasture also partially restored values for C/Po and Pi/o to levels closer to those for the soil with forest land use which might be attributed to the higher organic carbon, clay content compared to other land uses [24,25].

5. CONCLUSION

We analyzed how different types of land use affected the amount, fractionation, and distribution of phosphorus throughout the North-Western Himalayas. In addition to a 45% drop in soil C, we discovered that shifting land uses significantly altered P distribution. Changing the land use from forested to agricultural resulted in significant reductions in total organic carbon, total nitrogen, and phosphorus. More research into the impact of management practice on increasing P availability in these soils is needed.

ACKNOWLEDGEMENTS

A brief acknowledgement section may be given after the conclusion section just before the references. The acknowledgments of people who provided assistance in manuscript preparation, funding for research, etc. should be listed in this section. All sources of funding should be declared as an acknowledgement. Authors should declare the role of funding agency, if any, in the study design, collection, analysis and interpretation of data; in the writing of the manuscript. If the study sponsors had no such involvement, the authors should so state.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Rayne N, Aula L. Livestock manure and the impacts on soil health: A review. *Soil Systems*. 2020;4(4):64.
2. Hellal F, et al. Importance of phosphate pock application for sustaining agricultural production in Egypt. *Bulletin of the National Research Centre*. 2019;43(1):1-11.
3. Cordell D, Drangert JO, White S. The story of phosphorus: global food security and food for thought. *Global Environmental Change*. 2009;19(2):292-305.
4. Johnston A, Poulton P. Phosphorus in agriculture: A review of results from 175 years of research at Rothamsted, UK. *Journal of Environmental Quality*. 2019;48(5):1133-1144.
5. Naikoo NB, et al. Aggregation-dependent phosphorus adsorption under different land uses of district Kupwara of Kashmir Valley. *Journal of Plant Nutrition and Soil Science*; 2022.
6. Borie F, et al. Revisiting the nature of phosphorus pools in Chilean volcanic soils as a basis for arbuscular mycorrhizal management in plant P acquisition. *Journal of Soil Science and Plant Nutrition*. 2019;19(2)::390-401.
7. Vink JP, et al. Soils in lakes: the impact of inundation and storage on surface water quality. *Environmental Monitoring and Assessment*. 2020;192(6):1-18.
8. Zhang W, et al. Phosphorus fractionation related to environmental risks resulting from intensive vegetable cropping and fertilization in a subtropical region. *Environmental Pollution*. 2021;269: 116098.
9. Mäder P, et al., Arbuscular mycorrhizae in a long-term field trial comparing low-input (organic, biological) and high-input (conventional) farming systems in a crop rotation. *Biology and fertility of Soils*, 2000;31(2):150-156.
10. Stinner BR, Blair JM. Ecological and agronomic characteristics of innovative cropping systems, in *Sustainable agricultural systems*. 2020;CRC Press: 123-140.
11. O'Neil JM, et al. The rise of harmful cyanobacteria blooms: the potential roles of eutrophication and climate change. *Harmful Algae*. 2012;14:313-334.
12. Fonte SJ, et al. Pasture degradation impacts soil phosphorus storage via changes to aggregate-associated soil organic matter in highly weathered tropical soils. *Soil Biology and Biochemistry*. 2014;68:150-157.
13. Bouyoucos GJ. Hydrometer method improved for making particle size analyses of soils 1. *Agronomy Journal*. 1962;54(5):464-465.
14. Blake G, Hartge K, Peterson RG, Calvin LD. Bulk density Chapter 13 pp. 363-376. Klute, A. *Methods of soil analysis*. Part, 1986;1.
15. Jackson ML. *Soil chemical analysis-advanced course*. *Soil Chemical Analysis-Advanced Course*; 1969(Edition 2).
16. Walkley A, Black IA. An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. *Soil Science*. 1934;37(1):29-38.
17. Kuo S. Phosphorus. *Methods of soil analysis*. 1996;869-919.

18. Colwell J. The estimation of the phosphorus fertilizer requirements of wheat in southern New South Wales by soil analysis. *Australian Journal of Experimental Agriculture*. 1963;3(10):190-197.
19. Murphy J, Riley J. Colorimetric method for determination of P in soil solution. *Anal. Chim. Acta*. 1962;27:31-36.
20. Hedley M, Stewart J, Chauhan B. Changes in inorganic and organic soil phosphorus fractions induced by cultivation practices and by laboratory incubations. *Soil Science Society of America Journal*. 1982;46(5):970-976.
21. Stutter MI, et al. Land use and soil factors affecting accumulation of phosphorus species in temperate soils. *Geoderma*. 2015;257:29-39.
22. Foroughi M, et al. Hillslope position and land-use history influence P distribution in the critical zone, in *Biogeochemistry of the Critical Zone*. 2022;Springer:171-202.
23. Zhu M, et al. Soil P components and soil fungi community traits in poplar shelterbelts and neighboring farmlands in northeastern China: Total alterations and complex associations. *Catena*. 2022;218:106531.
24. McDowell R, et al. The efficacy of good practice to prevent long-term leaching losses of phosphorus from an irrigated dairy farm. *Agriculture, Ecosystems & Environment*. 2019;273:86-94.
25. Maharjan M, Maranguit D, Kuzyakov Y. Phosphorus fractions in subtropical soils depending on land use. *European Journal of Soil Biology*. 2018;87:17-24.

© 2022 Bashir et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/93616>