



Effect of Brown Manuring and Different Levels of Nutrients on Growth and Yield Attributes of Aerobic Rice (*Oryza sativa* L.)

M. Poornima Reddy^{a++*}, S. K. Chaudhary^{a#},
Akshay Kumar Kurdekar^{b†}, S. N. Sandeep^{c†}
and M. Vinay Kumar^{b†}

^a Department of Agronomy, RPCAU, Pusa Bihar, India.

^b Department of Agronomy, University of Agricultural Sciences, Raichur, Karnataka, India.

^c Department of Agronomy, University of Agricultural Sciences, Bangalore, Karnataka, India.

Authors' contributions

This work was carried under the guidance of my guide author SKC Co-authors helped in finalizing the treatments and performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2022/v12i121450

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/88366>

Original Research Article

Received: 15/04/2022

Accepted: 21/06/2022

Published: 17/11/2022

ABSTRACT

A field experiment was conducted during *kharif* season of 2019 to assess the Effect of brown manuring and different levels of nutrients on growth and yield attributes of aerobic rice (*Oryza sativa* L.) at Crop Research Centre, Dr Rajendra Prasad Central Agricultural University, Pusa, Bihar. This experiment was conducted in randomized block design (factorial) with sixteen treatment

⁺⁺ M.Sc. Student;

[#] Professor;

[†] Ph.D. Scholar;

*Corresponding author: E-mail: poornimareddym@gmail.com;

combinations replicated thrice using the rice variety "Abhishek" as a test crop which was planted with a spacing of 20 cm row to row, with the seed rate of 40 kg ha⁻¹. The experimental results indicated that higher values of growth parameters viz. Plant height (109cm), tillers/m² (355), dry matter production (969g/m²) and crop growth rate (12.2g/m²/day) were observed in rice + brown manuring treatment (B₂) as compared to rice without brown manuring (B₁) at harvest. While among the nutrient levels, treatment F₈- 70 kg/ha phosphorous and 50 kg/ha potassium as split recorded significantly higher growth parameters (tillers/m², dry matter production and crop growth rate), yield attributes (panicles/m², grains/panicle), grain yield, while remaining at par with treatment F₄ -60 kg/ha phosphorous and 50 kg/ha potassium as split. However, the interaction effect of both the factors did not significantly influence any of the parameters. Further the Aerobic rice cultivation save 40-45 per cent of water as compared to low land cultivation of rice.

Keywords: *Abhishek variety; aerobic rice; dhaincha; knockdown.*

1. INTRODUCTION

Ever since the beginning of agriculture, Cereals have been the premium source of diet for the man kind as they are rich in carbohydrates. A major cereal crop, Rice (*Oryza sativa* L.) is known to be a highly dominant grain crop worldwide. 21% of human per capita energy along with 15 % protein per capita has been supplied by rice universally, being a constituent of major proportion of daily diet in many South-East Asian countries. 60 per cent of population of earth lives in Asia, which contributes to 92 per cent of rice production in the world with an overall rice consumption of 90%. Countries like India, China, Bangladesh, Indonesia, Thailand and Vietnam collaboratively account to maximum production which is three quarters more compared to world rice production according to IRRI ("International Rice Research Institute"). In conventional puddled transplanted system, huge quantity of irrigation water is utilized for puddling. Although puddling is effective in controlling weeds, improving nutrient as well as water availability and facilitating transplantation and quick establishment of seedlings, it can adversely affect the physical properties of soil like poor structure of soil by breaking capillary pores and destroying soil aggregates that in turn leads to the creation of hard-pan, forming problems in establishing and development of succeeding crops. Considering high labour wages in the traditional rice production, low water availability, high cost of cultivation and low returns, "Aerobic rice" is preferable. Aerobic rice can be successfully raised with half of the water required as against puddled-transplanted rice in summer and entirely on rainfall in wet season with good rainfall distribution. Aerobic rice can save water to the tune of 45 to 55 per cent in summer season and even higher in *kharif* season as compared to continuous submerged crop [1].

Nitrogen fertilization is most crucial in improving the yield and quality of rice crop. Application of phosphatic fertilizers is important in aerobic rice compared to rice in flooded fields because dry aerobic soil can reduce the indigenous supply of phosphorous [2]. Potassium fertilizers applied at panicle initiation stage significantly increases the grain quality & yield [3]. Applying potassium either through KCl or KClO₃ in three equal split doses applied as basal dose, at panicle initiation and flowering stages could be suggested as a strategy of potassium management for yield maximization in aerobic rice [4].

Brown manuring (BM) technique has the advantage of supplementing all nutrients with crop and, is also beneficial for weed control and soil resources development. Procedure of brown manuring literally signifies a 'zero-till' form of green manuring, where a post-emergence herbicide is administered on BM to naturally decompose it without soil incorporation or ploughing till the residues decompose itself on the soil targeting to add organic manure into the soil besides weed suppression by the shade effect produced by brown manure crop [5]. Chlorophyll is lost from the leaves which were subjected to spray of herbicide in order to desiccate hence, leaves tend to change to brown in colour. This method helps the field by adding up to 35 kg/ha Nitrogen, dry matter, managing broad leaved weeds along with developing the yield of grain up to 4 -5 q/ha because of OM addition into soil. On lighter soils prone to erosion this practice shall be preferred. Along with adding organics to soil, BM helps to enhance the soil properties. Hence knowing the beneficial effect of brown manuring, The study was carried out in order to find the combined effect of brown manuring and different levels of nutrients on growth and yield attributes of aerobic rice.

2. MATERIALS AND METHODS

The field experiment conducted during *kharif* period of 2019 at the RPCAU, Pusa, Bihar. The soil of the experimental field was calcareous in nature containing 26% free CaCO₃, sandy loam in texture, alkaline in nature with a pH of 8.3 and EC 0.49 dSm⁻¹. The soil contained low in organic carbon (0.45%), available nitrogen (243kg ha⁻¹) and potassium (112 kg ha⁻¹), while medium in available phosphorus (11kg ha⁻¹). The experiment was conducted in factorial randomised block design with sixteen treatment and replicated three times (Table 1). Abhishek variety of rice was sown according to the date decided in the treatment, planted with a spacing of 20 cm (RR), with the seed rate of 40 kg ha⁻¹. The treatment details are mentioned in Table 1. BM indicates Brown manuring with dhaincha @ 40 kg/ha seed rate and knock down at 25 DAS using a post-emergence herbicide, K_B indicates basal application of 100% Potassium (K), K_S indicates split application of 100% K (60% as basal dose and 40% top dressing during panicle initiation stage of the crop). Along with these treatments the recommended dose of nitrogen (RDN) applied @ 140 kg N/ha. All recommended agronomic practices (weeding, hoeing,

pesticides, irrigation etc.) were kept uniform for all the treatment and were carried out throughout the growing season, when required. During the experiment observations were recorded using the standard procedure for each parameter.

3. RESULTS AND DISCUSSION

3.1 Crop Growth Rate (g/m²/day)

The figures calculated pertaining to Crop Growth Rate (CGR) at all the growth stages (0-25, 25-50, 50-75 DAS and 75 DAS -at harvest) are furnished in the Table 2. A clear data appraisal in from table here under denotes improvement in crop growth rate from 0-25 DAS up to 50-75 DAS which decreased later on as the crop age increased gradually while reaching the physiological maturity stage. At 0-25 DAS, neither brown manuring nor different levels of nutrients indicated their effect on this factor in aerobic rice. It was non-significant irrespective of the treatments applied. At 25-50 DAS, it was observed that significantly higher crop growth rate was found with brown manuring (7.37g/m²/day) compared to rice alone (5.91 g/m²/day). Among the eight nutrient management

Table 1. Details of treatments

Factor 1. System of cultivation		Indicative symbol in the layout
1) Rice alone		B ₁
2) Rice + Dhaincha (BM)		B ₂
Factor 2. Levels of P and K fertilizers		
1) P 60 K _B 40 (P ₂ O ₅ @ 60 kg/ha and K ₂ O @ 40kg/ha as a complete basal dose)		F ₁
2) P 60 K _B 50 (P ₂ O ₅ @ 60 kg/ha and K ₂ O @ 50kg/ha as a complete basal dose)		F ₂
3) P 60 K _S 40 (P ₂ O ₅ @ 60 kg/ha and K ₂ O @ 40kg/ha as split application)		F ₃
4) P 60 K _S 50 (P ₂ O ₅ @ 60 kg/ha and K ₂ O @ 40kg/ha as split application)		F ₄
5) P 70 K _B 40 (P ₂ O ₅ @ 70 kg/ha and K ₂ O @ 40kg/ha as a complete basal dose)		F ₅
6) P 70 K _B 50 (P ₂ O ₅ @ 70 kg/ha and K ₂ O @ 50kg/ha as a complete basal dose)		F ₆
7) P 70 K _S 40 (P ₂ O ₅ @ 70 kg/ha and K ₂ O @ 40kg/ha as split application)		F ₇
8) P 70 K _S 50 (P ₂ O ₅ @ 70 kg/ha and K ₂ O @ 50kg/ha as split application)		F ₈

treatments, F_8 (7.24 g/m²/day) resulted in highest CGR which was statistically on par with F_5 (6.86 g/m²/day), F_6 (7.06 g/m²/day) and F_7 (7.11 g/m²/day). At 50-75 DAS, brown manured plots (16.85 g/m²/day) resulted in markedly lesser CGR compared to rice alone (17.58) but different doses of phosphorous and potassium failed in producing any changes in CGR. Both factors under study gave significant results with respect to CGR at harvest. Significantly greater CGR was recorded with brown manuring treatment whereas among nutrients, F_8 treatment i.e., 70 kg P₂O₅ & 50 kg K₂O as split resulted in highest CGR (12.72 g/m²/day) which was at par with F_6 (11.62 g/m²/day) and F_7 (12.43 g/m²/day) but superior than other treatments. Phosphorous and potassium possess a crucial role in photosynthesis of any plant. Potassium plays a positive part in metabolic activity of the plant that leads to increase in photosynthesis and thereby increases dry matter production. Hence, the growth indices like CGR was higher in treatments where K fertilizer was applied in split as observed by [6].

3.2 Number of Panicles/m² and Panicle Length (cm)

Effect caused by brown manuring and differed levels of nutrients on yield attributes and yield is given in the Table 3. Recorded data exhibited

that brown manuring treatment did not produce significant results in panicle length after a clear perusal. However, slightly longer panicle length was observed with brown manuring compared (21.38cm) to rice without brown manuring (21.15cm). Among the different levels of nutrients none of the treatments significantly affected the panicle length. It was found non-significant however, among eight treatments highest panicle length (21.55cm) was recorded with F_8 treatment while the lowest panicle length (21.05cm) was seen in F_1 treatment with marginal differences through other fertilizer doses. The both brown manuring and nutrient levels resulted in significant Number of panicles produced in a unit area. The no. of panicles produced in brown manuring (285) were significantly superior over rice alone treatment (268). Whereas among the various levels of phosphorous and potassium, F_8 treatment (297) resulted in increasing panicle numbers significantly while being statistically in comparison related to F_4 , F_5 , F_6 and F_7 . Highest number of panicles per m² might be attributed to lot of effective tillers/m² produced as an outcome of BM and Paddy crop receiving additional supply of N, P and K nutrients as for gradual decomposition of brown manure caused higher intake of nutrients also better availability for longer duration resulting in higher yield attributes as observed by [7].

Table 2. Effect of brown manuring and different levels of nutrients on crop growth rate

Treatments	Crop growth rate (g/m ² /day)			
	(0-25) DAS	(25-50) DAS	(50-75) DAS	(75DAS) -At harvest
Factor A: System of cultivation				
B ₁ (rice alone)	0.81	5.91	17.58	11.21
B ₂ (rice+sesbania)	0.79	7.37	16.85	12.20
SEm (±)	0.01	0.09	0.25	0.15
CD (P=0.05)	NS	0.2	0.7	0.4
Factor B: Different levels of phosphorous and potassium				
F ₁ (P60 K _B 40)	0.77	5.94	16.37	10.73
F ₂ (P60 K _B 50)	0.79	6.32	16.95	10.93
F ₃ (P60 K _S 40)	0.78	6.25	16.75	11.73
F ₄ (P60 K _S 50)	0.79	6.38	16.61	12.01
F ₅ (P70 K _B 40)	0.82	6.86	17.52	11.47
F ₆ (P70 K _B 50)	0.82	7.06	17.82	11.62
F ₇ (P70 K _S 40)	0.81	7.11	17.73	12.43
F ₈ (P70 K _S 50)	0.82	7.24	17.93	12.72
SEm (±)	0.02	0.18	0.50	0.30
CD (P=0.05)	NS	0.5	NS	0.8
Interaction(B*F)	NS	NS	NS	NS

Table 3. Effect of brown manuring and various nutrient levels on the yield attributes and grain yield

Treatments	Number of panicles /m ²	Panicle length (cm)	No. of grains in a panicle	Test weight. (g)	grain yield (q/ha)
Factor A: System of cultivation					
B ₁ (rice alone)	268	21.15	92	20	40.64
B ₂ (rice + sesbania)	285	21.38	98	21	42.79
SEm (±)	5.2	0.4	1.8	0.4	0.69
CD (P=0.05)	15.0	NS	5.4	NS	2.0
Factor B: Different levels of phosphorous and potassium					
F ₁ (P60 K _B 40)	258	21.05	90	20	39.55
F ₂ (P60 K _B 50)	261	21.07	93	20	40.37
F ₃ (P60 K _S 40)	259	21.22	96	20	40.96
F ₄ (P60 K _S 50)	270	21.24	96	21	41.56
F ₅ (P70 K _B 40)	279	21.20	96	21	41.79
F ₆ (P70 K _B 50)	288	21.25	96	21	42.57
F ₇ (P70 K _S 40)	293	21.53	98	21	43.16
F ₈ (P70 K _S 50)	297	21.55	100	21	43.79
SEm (±)	10.4	0.83	3.77	0.80	1.39
CD (P=0.05)	30.2	NS	10.9	NS	4.0
Interaction(B*F)	NS	NS	NS	NS	NS

3.3 No. of Grains in a Panicle

A clear appraisal of data (Table 3) indicated that number of filled grains produced per panicle were significantly influenced by the present treatments. Brown manuring produced outstandingly more grain production compared to no brown manuring. Gradual release of nitrogen due to incorporation of dhaincha relatively throughout the primordial initiation, flowering alongside grain filling phases helped to increase the number of spores budding on stigma that resulted in increased amount of spore germination on stigma which in turn enhanced the amount of filled grains in a panicle. The results stated here were found to be in agreement to the outcome of [8] and [9]. Whereas 70kg phosphorous and 50kg potassium in split application resulted in highest number of grains/panicles which was, significantly superior over F₁ viz. 60 kg phosphorous & 40 kg potassium as basal, but the other treatments did not provide significant results. The most elevated estimations of growth characteristics acquired with the use of phosphorus showed signs of improvement in yield qualities which might be likely because of more ingestion and usage of accessible phosphorous for absorption and utilization by plants, by and large improvement of "source-sink relationship", which thus upgraded the yield characteristics of aerobic rice [10]. Utilization of K brought about better plant development and dry matter yield. Adjusted supply of N, P and K was liable for higher

photosynthesis, metabolic action and cell division, which therefore expanded development and yield qualities as detailed by [11]. Comparable discoveries as per the current examination were given by [12].

3.4 Test Weight (g)

Values regarding test-weight did not seem to have any significant variation because of the brown manuring or nutrient management treatments. However slightly higher test weight was observed with brown manuring in rice (21g) whereas rice without brown manure showed lower test weight (20g). Among the nutrient levels, non-significant results were seen although F₈ treatment showed slightly higher test weight than others. The interaction results too remained non-significant.

3.5 Grain Yield (q/ha)

Grain yield obtained through brown manuring (42.79 q/ha) was significantly superior over yield from rice alone plots (40.64 q/ha). Rice grown along with brown manure yielded 5.29% higher grain than rice without brown manuring. Fundamentally more grains could be achieved from B₂-rice with dhaincha contrasted with rice alone, which had significant differences which may have come about because of the higher estimations of yield properties, for example, number of panicles/m², panicle length, grains in panicle and weight of 1000 grains.

Comparable outcomes were recorded by [13]. The residues of brown manure (dhaincha) acted as mulch increasing the availability of moisture which would have contributed to increase in yield attributes and finally leading to enhanced grain & straw yield (Yadav et al., 2011). Comparison of effect of different nutrient levels on grain yield indicated that F₈ treatment gave significant results by producing 43.79q/ha which was significant over F₁ treatment but at par with all others. The difference was to the tune of 8.72%. The vital role taken up by potassium in improving the yield attributes significantly increased the grain yielding capability of the crop. As a consequence of split application of K, a greater number of filled grains in a panicle and higher panicle length were observed. The production of yield attributes and grain yield exhibited a strong positive correlation. Aforesaid findings have similarity to the results given by [14].

4. CONCLUSION

It can be concluded that brown manuring and implementation of 140kg N, 60kg P₂O₅ and 50kg K₂O as split found beneficial in terms of growth and yield attributes for aerobic rice. Along with that Aerobic rice could be one of the best alternative technology that can save 30 to 40 per cent of water compared to traditional lowland cultivation.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Gandhi RV, Rudresh NS, Shivamurthy M, Hittalmani S. Performance and adoption of new aerobic rice variety MAS 946-1 (Sharada) in southern Karnataka. Kar. J. Agric. Sci. 2012;25;25(1).
- Yosef Tabar S. Effect of nitrogen and phosphorus fertilizer on growth and yield rice (*Oryza sativa* L). Int. J. Agron. Plant Prod. 2012;3(12):579-584.
- Wakeel A, Rehman HU, Mubarak MU, Dar AI, Farooq M. Potash use in aerobic production system for basmati rice may expand its adaptability as an alternative to flooded rice production system. Journal of soil science and plant nutrition. 2017; 17(2):398-409.
- Babu A, Hattab O, Aruna L, Mohan R. Effect of sources, split and foliar application of KCl and KClO₃ on uptake of potassium in aerobic rice. J. Pharmacogn. Phytochem. 2018;7(4):1954-1957.
- Nawaz A, Farooq M, Lal R, Rehman A, Hussain T, Nadeem A. Influence of sesbania brown manuring and rice residue mulch on soil health, weeds and system productivity of conservation rice-wheat systems. Land Degrad. Dev. 2017; 28(3):1078-1090.
- Jinger D, Dhar S, Dass A, Sharma VK, Joshi E, Vijayakumar S, Gupta G. Effect of silicon and phosphorus fertilization on growth, productivity and profitability of aerobic rice (*Oryza sativa*) Indian J. Agric. Sci. 2018;88(10):1600-1605.
- Yadav GS, Shivay YS, Kumar D. Effect of mulching and iron nutrition on productivity, nutrient uptake and economics of aerobic rice (*Oryza sativa*). Indian J. Agron. 2011; 56(4):365-372.
- Mohioddin MG, Ramana MV, Sridevi S, Reddy AP. Influence of integrated nitrogen management on yield components and yield of scented rice (*Oryza sativa* L.). The J. Res ANGRAU. 2014;42(2):65-68.
- Chaudhary SK, Marahatta S, Chaudhary M. Performance of dry direct seeded rice and weeds on Sesbania brown manuring as compared to farmers' practice and chemical control method. Int. J. Appl. Sci. Biotechnol. 2018;6(3):265-269.
- Sharma SN, Prasad R, Shivay YS, Dwivedi MK, Kumar S, Kumar D. Effect of rates and sources of phosphorus on productivity and economics of rice (*Oryza sativa*) as influenced by crop-residue incorporation. Indian J. Agron. 2009; 54(1):42-46.
- Vijayakumar, S. Potassium management in aerobic rice (*Oryza sativa* L.) – wheat (*Triticum aestivum* L.) cropping system. Ph.D Thesis, Indian Agricultural Research Institute, New Delhi, India; 2017.
- Awan TH, Manzoor Z, Safdar ME, Ahmad M. Yield response of rice to dynamic use of potassium in traditional rice growing area of Punjab. Pak. J. Agric. Sci. 2015; 44(1):130-135.

13. Samant TK. A study on effect of brown manuring on growth, yield, economics and soil fertility in direct seeded rice (*Oryza sativa* L.). J. Bio. Innov. 2017;6(4):637-643.
14. Arivazhagan K, Revichandran M. Interaction effect of nitrogen and potassium on yield and yield attributes in rice cv. IR 20. Adv. Plant Sci. 2005;20: 425-7.

© 2022 Reddy et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/88366>