



Genitourinary Syndrome of Menopause (GSM)- Pap Test and Gram Findings

**José Eleutério Junior^{a,b*}, Renata Mírian Nunes Eleutério^{b,c},
João Victor da Silva Bezerra^a, Armanda do Carmo Ponce de Leão Lins^a,
Marina Helena da Silva Lopes^a and Ana Katherinne Silveira Gonçalves^d**

^a Postgraduate Program in Pathology, Federal University of Ceará (UFC), Fortaleza, Brazil.

^b Prof. Eleutério Laboratory, Fortaleza, Brazil.

^c Biomedicine Course - Christus University Center, Fortaleza, Brazil.

^d Department of Obstetrics and Gynecology, Federal University of Rio Grande do Norte (UFRN),
Natal, RN, Brazil.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/93446>

Mini-review Article

Received 11 September 2022

Accepted 14 November 2022

Published 22 November 2022

ABSTRACT

The loss of ovarian function, prematurely or due to menopause, causes a decrease of the vaginal epithelium trophy. However, with the end of the menstrual cycle and the consequent hypoestrogenism, the epithelial thickness also decreases, with only the parabasal and basal layers remaining with low glycogen reserves. A thinner epithelial thickness associated with a pH greater than 4.5 and a microbiota without protective bacteria (*Lactobacillus* sp) can lead to an inflammatory condition not necessarily related to the action of a pathogen but mainly the atrophy itself. Atrophic vaginitis is often confused with genitourinary syndrome of menopause (GSM). Laboratory findings do not necessarily coincide with symptoms but could help in diagnosing. Examinations such as bacterioscopy of vaginal smear (Gram) and cytology (Pap test or cytology in liquid medium) have findings that may suggest a picture of intense hypoestrogenism. However, the clinical-laboratory correlation is essential.

Keywords: Genitourinary syndrome of menopause; vaginitis, papanicolaou test; cytology.

1. INTRODUCTION

The loss of ovarian function, prematurely or due to menopause, causes a decrease of the vaginal epithelium trophic. Sometimes the women refer to localized symptoms leading to discomfort and influencing sexual activity [1].

In 2014 the International Society for the Study of Women's Sexual Health and the North American Menopause Society introduced genitourinary syndrome of menopause (GSM) as changes associated with lack of estrogen. This lack of estrogen generates various clinical symptoms such as dryness, itching, friability of the vaginal mucosa and dyspareunia [2]. From a clinical point of view the main goal of treatment is the relief of these and other symptoms that can disrupt the patient's daily life [3]. Sometimes the term GSM is confounding with atrophic vaginitis, but this is only a possible finding in the syndrome, besides urinary and sexual complaints [4]. However, it is essential to understand that the laboratory finding of atrophy with inflammatory infiltrate does not always correspond to complaints and clinical findings [5].

2. PATHOPHYSIOLOGY

The vaginal epithelium originated from the urogenital sinus and the Müllerian duct, is a non-keratinized stratified epithelium that modifies its thickness according to the estrogen stimulus. There are the following layers in the entire thickness of the epithelium: basal, parabasal, intermediate, and superficial. Depending on the menstrual cycle phase and, therefore, on the estrogenic and progestin influence, there is a greater glycogen accumulation in the epithelium's intermediate layer (post-ovulatory phase). The glycogen is metabolized to produce lactic acid and maintain physiological pH in the vagina (<4.5) [6,7].

In women who have a menstrual cycle, the highest level of epithelial thickness occurs in the periovulatory period, when there is also a higher estrogen level. However, with the end of the menstrual cycle and the consequent hypoestrogenism, the epithelial thickness also decreases, with only the parabasal and basal layers remaining with low glycogen reserves. A thinner epithelial thickness associated with a pH greater than 4.5 and a microbiota without

protective bacteria (*Lactobacillus* sp) can lead to an inflammatory condition not necessarily related to the action of a pathogen but mainly the atrophy itself [8].

2.1 Laboratory Exams

Some tests can be used to help diagnose the atrophic vaginitis.

Gram: in cases of women with hypoestrogenism (atrophy) the microbiota study in a Gram-stained smear identifies a microbiota of sparse Gram-positive cocci, sometimes Gram-negative, amid rounded cells (parabasal). This finding differs from oestrogenized women with a well-developed microbiota, with a predominance of gram-positive medical bacilli (*Lactobacillus* sp). However, atrophy with inflammation or atrophic vaginitis is characterized by infiltrating inflammatory cells, predominantly polymorphonuclear leukocytes, in addition to the criteria defined above [9].

Pap test: the Pap smear is a widely used method in secondary prevention of cervical cancer, but it can observe other findings that help diagnose different diseases. But, for a long time, the Pap smear was also used to assess hormonal status. Analyzing the cell types of vaginal smears, it was possible to suggest the action or not of estrogen and progesterone.

Today, cytology reports follow the Bethesda system, which classified atrophy as "non-neoplastic findings" in its latest version [10].

Criteria for diagnosing atrophy and atrophic vaginitis are monolayer sheets of round cells with the high nucleo-cytoplasmic rate (parabasal cells), as well dispersed and isolated cells. Sometimes only nuclei are observed without cytoplasm (autolysis). When abundant inflammatory cells and rounded blue bodies ("blue blobs") are present, we think about extreme atrophy or atrophic vaginitis [11,12].

Liquid-based cytology: the liquid-based cytology, which uses the same polychromatic stain, has slightly different findings. The smear background is clean, parabasal cells are better observed, and autolysis is absent. However, the inflammatory infiltrates and "blue blobs" indicate, as in the conventional cytology (Pap test), atrophic vaginitis [12].

Crothers et al. [13] studied conventional cytology and two methods most used in liquid cytology (Thinprep and Surepath). The authors concluded that in Thinprep, cases were often considered

unsatisfactory. Even Surepath having fewer unsatisfactory cases was more discordant in diagnosing atrophy with inflammation than the conventional smear.




Fig. 1. The aspect of cytological atrophy in vaginal smear (Gram 1000x). We can observe the parabasal cells (rounded), numerous leukocytes and scarce cocaceous microbiota




Fig. 2. Findings of atrophy with inflammation in conventional cytology (Pap 400x). We can see parabasal cells, inflammatory infiltrate, scarce microbiota, filaments, and blue blobs (arrow)




Fig. 3. Atrophy with inflammation in liquid-based cytology (Surepath 400x). We can observe isolated and grouped well-preserved parabasal cells, inflammatory infiltrate, sparse microbiota and blue blobs

3. CONCLUSION

GSM is a syndrome that may have in the morphological laboratory exam, Gram and cytology, a tool to help diagnosis. However, the finding of a picture suggestive of atrophy with inflammation does not necessarily indicate the need for treatment; in these cases, the objective is the patient's well-being and the relief of symptoms. The clinical-laboratorial correlation is vital for a correct diagnosis and adequate management.

CONSENT AND ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Beard MK. Atrophic vaginitis. Can it be prevented as well as treated? *Postgraduate Medicine* 1992, 91:6, 257-260.
DOI: 10.1080/00325481.1992.11701327
2. Phillips NA, Bachmann GA. Genitourinary syndrome of menopause: common problem, effective treatments. *Cleve Clin J Med*. 2018;85:390–398.
DOI: 10.3949/ccjm.85a.15081
3. Angelou K, Grigoriadis T, Diakosavvas M, Zacharakis D, Athanasiou S. The Genitourinary Syndrome of Menopause: An Overview of the Recent Data. *Cureus*. 2020;12(4): e7586.
DOI: 10.7759/cureus.7586.
4. Angelou K, Grigoriadis T, Diakosavvas M, et al. (April 08, 2020) The Genitourinary Syndrome of Menopause: An Overview of the Recent Data. *Cureus* 12(4): e7586.
DOI 10.7759/cureus.7586.
5. Heller DS, Weiss G, Bittman S, Goldsmith L. Does a diagnosis of atrophic vaginitis on Papanicolaou test signify the presence of inflammation? *Menopause*. 2015;22(8): 814-5.
DOI: 10.1097/GME.0000000000000393.
6. Kurita T. Developmental origin of vaginal epithelium. *Differentiation*. 2010;80(2-3):99-105.
DOI: 10.1016/j.diff.2010.06.007. Epub 2010 Jul 17.
7. Godha K, Tucker KM, Biehl C, Archer DF, Mirkin S. Human vaginal pH and microbiota: an update. *Gynecol Endocrinol*. 2018 Jun;34(6):451-455.
DOI: 10.1080/09513590.2017.1407753. Epub 2017 Dec 22
8. Brizzolara S, Killeen J, Severino R. Vaginal pH and parabasal cells in postmenopausal women. *Obstet Gynecol*. 1999 Nov;94(5 Pt 1):700-3.
DOI: 10.1016/s0029-7844(99)00384-1.
9. Kim S, Seo H, Rahim MA, Lee S, Kim YS, Song HY. Changes in the microbiome of vaginal fluid after menopause in Korean women. *Microbiol Biotechnol*. 2021;31(1).
DOI: 10.4014/jmb.2106.06022
10. Wilbur DC, Nayar R. Bethesda 2014: improving on a paradigm shift. *Cytopathology*. 2015;26(6):339-42.
DOI: 10.1111/cyt.12300.
11. Solomon D, Nayar R (Eds). *The Bethesda system for reporting cervical cytology. Definitions, criteria, and explanatory notes*. 3rd Ed. New York: Springer; 2015.
12. Yakoushina TV, Medina IM, Hoda RS. "String of pearls" appearance of blue blobs in postmenopausal atrophy on ThinPrep Pap test. *Diagn Cytopathol*. 2009;37(10): 738-9.
DOI: 10.1002/dc.21008
13. Crothers BA, Booth CN, Darragh TM, Means MM, Souers RJ, Thomas N, Moriarty AT. Atrophic vaginitis: concordance and interpretation of slides in the College of American Pathologists Cervicovaginal Interlaboratory Comparison Program in Gynecologic Cytopathology. *Arch Pathol Lab Med*. 2012;136(11): 1332-8.
DOI: 10.5858/arpa.2011-0441-CP

© 2022 Junior et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/93446>