



An Overview of Lingual Orthodontics: Advantages and Disadvantages

Rajeev Sharma ^{a#} and Deepak Mohan Soni ^{b#*}

^a *PHC Summerkot, Directorate of Dental Health, Shimla, Health and Family Welfare Department, Himachal Pradesh, India.*

^b *CHC Kuthera, Directorate of Dental Health, Shimla, Health and Family Welfare Department, Himachal Pradesh, India.*

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/93182>

Review Article

Received 17 October 2022
Accepted 19 December 2022
Published 22 December 2022

ABSTRACT

The majority of orthodontic issues that can be resolved with conventional labial technique, such as tooth malposition, anteroposterior discrepancies, and pre-prosthetic surgical cases, can be resolved with lingual orthodontics. The lingual surfaces of the teeth appear to be less prone to caries than the buccal surfaces due to differences in surface morphology, plaque retention, salivary flow, and the mechanical cleaning of surfaces by the tongue. Lingual orthodontics is one of the good options for comprehensive treatment of the majority of malocclusions. Despite being a technique-sensitive and demanding orthodontic modality in terms of expertise, special bracket and wire system, complicated system of wire bending, need for indirect bonding, and customization of treatment plan, lingual orthodontics has unquestionably emerged as the most desirable treatment option amongst the latest treatment modalities available because of its most acceptable form of invisible braces, especially in adults and adolescents or more appropriately young adults due to the fact that lingual orthodontics is a treatment option that can be used on both adults and adolescents.

Keywords: Orthodontics; dentistry; orthodontic modality; lingual straight wire technique.

[#] *MDS Orthodontics;*


^{*} *Corresponding author: Email: sonijamesdeepak@gmail.com, sonijamesdeepak@yahoo.com;*

1. INTRODUCTION

Orthodontics is a branch of dentistry which has undergone evolutionary progress with time because of the critical situation of managing the developing malocclusions in growing children or fully developed malocclusions in adults; hindrances in achieving esthetics, functionality & structural balance; not ignoring the foremost objective of fulfilling the patient's desires. The complications encountered in achieving the goals has furthered the research in orthodontics which resulted in the development of various appliances and techniques. The foremost problem faced by orthodontist through its evolution was patient acceptability of the braces because of the unsightly appearance and maneuverability of the braces as some sort of handicapping situation for the patient by kith and kins. Emphasis for development of lingual appliances came from the unesthetic perception of labial fixed orthodontic appliances [1]. Although lingual orthodontic treatment has disadvantages of soft tissue injury and insufficient area for the tongue, the distinct advantage of invisibility and eradication of a feeling of shame in public appearances, surpasses all its other complications which can be managed with some extra efforts in maintaining oral hygiene and improvising upon the eating habits etc. Satisfactory results and the benefit of hidden appliances, convinces most of patients to endorse the lingual orthodontic appliance as beneficial to friends and colleagues [2]. Today lingual technique is not only accepted, rather preferred by adults [3].

For this review, articles were searched in the pubmed database with the keyword "lingual orthodontics". 56 articles were downloaded and scrutinized for important data and finally 21 articles were included. Important excerpts from several books and few other articles were included depending upon the relevance to the topic.

Following areas about lingual orthodontics would be covered as depicted in the flow chart:



2. EVOLUTION OF LINGUAL ORTHODONTICS

It was Pierre Fauchard in 1726, who first proposed that braces could be attached onto the lingual surfaces of teeth [4]. First lingual arch for expansion and alignment of the teeth was created by Pierre Joachim Lefoulon in 1841 [4].

In 1889, John Farrar described "Lingual removable Arch" with some minor modifications in the labial appliance [5]. He described it as a wire of suitable size, to which auxiliary springs can be attached [6].

Dr. Oren Oliver in March 1942, at a pan American congress in New Orleans, took on a demonstration on a labiolingual appliance [7].

Dr William Wilson demonstrated a labio-loop lingual appliance in mid 50's that later developed into the Wilson modular appliance system [7].

As the multibracket labial appliances advanced over time, partial lingual appliances were developed that were used as adjunct to the labial appliance systems such as the Goshgarian (transpalatal bar), Ricketts (Quad-Helix) and Wilson (3D Modular Enhanced Orthodontics).

The idea of Contemporary lingual treatment began simultaneously in 1970's in two separate countries, by Dr Kinya Fujita (Kanagawa, Japan) and Dr Craven Kurz (Beverly Hills, CA, USA) who independently developed their own designs for lingual braces.

The lion's share in development of current lingual orthodontics came from Dr Craven Kurz in 1975, who was instrumental in introducing a bonded edgewise lingual appliance with few inbuilt features for the first time in association with Ormco company and further developed seven generations of lingual brackets with horizontally opening slots, each one more refined and sophisticated than the previous generation [8,9].

Mushroom shaped arch wires were introduced for the first time and promoted in lingual orthodontics by Dr Kinya Fujita from Kanagawa Dental University, Japan. Mushroom archwires however, typically involve intricate wire bending in the form of vertical step bends and first premolar insets. He initiated his endeavours in direction of developing lingual technique in 1968, paradoxically not for esthetic purpose but for protecting his patients who practised martial arts,

from injury by labial brackets and after a decade of hard work and research, in 1978, described the capabilities of the Fujita Brackets, which have an occlusally opening slot [10].

Tae Weon Kim founded the Korean Society of Lingual Orthodontics (KSLO) and developed Korean indirect bonding system. Hee-Moon Kyung founded the Korean Lingual Orthodontics Association (KLOA) and developed the Individual Indirect Bonding Technique (IIBT), the Mushroom Bracket Positioner, as well as the Lingual Straight Wire Technique [11].

Pioneers of lingual orthodontics in the United States were Kelly [9]. (1982), who used labial brackets from Unitek on the palatal/lingual surfaces and Dr. Stephen Paige [12,13]. who introduced Lingual Light Wire technique in 1982 using brackets from Begg's technique on palatal surfaces. Dr. Thomas D. Creekmore [14]. (1989) presented a comprehensive technique and lingual brackets with vertical slot and the Slot Machine which was capable of efficient and precise laboratory procedures.

Weichmann introduced computer aided designing/ computer aided machining (CAD/CAM) into lingual orthodontics in 2002 and brought renaissance in the field by completely customizing lingual brackets and archwires with wire bending through robotic technology. In 2009, 3M Unitek took its copyrights and reintroduced it as Incognito™ lingual appliance [9].

Fillion, in 2010, used the Orapix digital system with a virtual setup to develop a customized straight-wire technique [15].

Many other customized lingual systems have been introduced since then e.g. HARMONY (American Orthodontics), WIN (DW Lingual Systems GmbH) and Indian customized lingual systems: Lingual Matrix and I-Lingual 3D.

Dr Massimo Ronchin and Forestadent (Pforzheim, Germany) developed a self-ligating bracket based on the Begg's technique in 1994. First self-ligating lingual brackets in its pure form were developed by Scuzzo et al. in 2011 [16].




Fig. 1.a. Evolution of the Craven Kurz lingual bracket

Fig. 1 b. 7th generation Curz bracket

- *First generation (1976)*
- *Second Generation(1980)*
- *Third Generation (1981)*
- *Fourth Generation (1982-1984)*
- *Fifth Generation (1985-86)*
- *Sixth Generation (1987-90)*
- *Seventh Generation (1990-present) [20].*

Kairalla et al. [17] established the size and shape of lingual arch forms in 2014 as small, medium, large and extra large and shape similar to a parabola with a slight flattening in the anterior teeth region.

Park et al. [18] introduced a new nomenclature for lingual arch forms in 2015: Narrow, tapering and ovoid.

Various practitioners through out the world have been working upon for development of an efficient lingual system by devising their own bracket system and evolving their techniques.

2.1 Generations of Kurz Brackets

The main features of the Kurz Brackets were a bite plane, a base pad adapted to the anatomic morphology of the lingual surfaces and a preangulated slot according to the torque to be applied on the labial surface [19].

The seventh generation Ormco bracket presented with a larger bracket base, pre-angulated slot and hook, maxillary anterior bite plane now heart shaped with smaller hooks. The premolar brackets have increased width to allow better angulation and rotation control. The molar now comes with either a hinge cap or a terminal sheath [21,22].

2.2 Fujita's Lingual Brackets

Dr. Kinya Fujita developed small sized, low profile brackets in 1979, so that any injury to teeth or soft tissues during physical activities could be avoided. The occlusal slot enabled convenient insertion of archwire into the slot and avoided deformation of archwire at the time of insertion. The grooves for lockpins were parallel to the wire i.e. mesiodistally. An auxiliary groove was also present occlusogingivally for mesiodistal correction.




Fig. 2. Fujita lingual bracket

Currently multiple slots for different wires are present in Fujita brackets in occlusal, lingual and vertical directions i.e. 3 slots in Anterior and premolar brackets and 5 slots in molar brackets so that comprehensive mechanics for different movements of teeth with multiple wires could be carried out simultaneously [23].

3. TYPES OF BRACKET SYSTEMS

Some of the systems are:

1. Conceal brackets by Thomas Creekmore [14,20]. : occlusally opening slots and convenient archwire insertion and removal.
2. STb (SCUZZO- TAKEMOTO bracket) (Sold by Ormco.) [20,24,25]. developed straight wire lingual technique by placing the brackets more gingivally.
3. Forestadent: Low profile 2D-brackets and 3D-brackets for effortless management of elementary problems and severe discrepancies respectively; facilitating three dimensional control of torque and angulations of individual tooth [21].
4. Stealth Brackets (American orthodontics). [21].
Old generation brackets with tiny proportions & large distance between adjacent brackets, comfortable to the patient but below par control of tooth rotations. Rotations had to be managed by an uprighting spring.
5. Philippe self ligating lingual brackets [26, 27].
No slots therefore could not control torque. Direct bonding was employed.
6. Kelly Bracket
A simple labial unitek bracket used on the lingual surface. A twin bracket with a Horizontal insertion slot best for managing tooth rotations.
7. In-ovation L bracket [28].
A self ligating bracket with large distance between adjacent brackets, therefore more comfortable for the patient. Because of the forked design in the base there is better synchronisation with the lingual surface contours.
8. Braces (Incognito) [20,27].
Customized low Profile brackets with good finish. The CAD-CAM technology has enabled the process of patient specific bracket production and precise bracket positioning possible in a single step in the laboratory thus producing accurate movements of the desired teeth only.




Fig. 3a. Conceal brackets have three different arch wire slot widths for the three different tip A-B, torque E-F, and rotation C-F or E-D




Fig. 3b. Conceal brackets A: Lingual insertion B: Occlusal insertion.



Fig. 3c. Stealth Brackets



Fig. 3d. Forestadent



Fig. 3d. Philippe bracket



Fig. 3e. In –ovation L bracket

4. ARCHWIRES IN LINGUAL ORTHODONTICS

Because of the anatomical variations in the morphology of the lingual surfaces, adapting the brackets intimately to the surface becomes a cumbersome process and therefore proper engagement of the archwire into the bracket slot requires intricate wire bending or preformed wires shaped according to the generalized archform on the lingual aspect of the teeth. Hence two types of archwire forms have been developed to be used in routine practice of orthodontics: the straight (ovoid) and mushroom-shaped archwires.

The mushroom shaped archwire is shaped according to the general lingual surface contours having sharp inset between canine and first premolar, whereas a thick base has to be built in the bracket for use with the straight archwire or some surplus composite has to be added under the bracket base to compensate for the different labio-lingual thickness of various teeth. Nevertheless, the difference is very small in a horizontal plane at the junction of crown and root, therefore a "lingual straight archwire" having a configuration of a simple curve in the horizontal plane with no insets has been developed.

Both the types of archwires need to employ different mechanics, because the location of force and/or moment application with respect to the centre of resistance (CR) varies as the thickness of the base in the anterior region, primarily lateral incisors and canines, increases. In addition the first-order bends in the form of insets also changes the manner in which loads are transmitted in the archwire.

Although very little research has been conducted to differentiate between mushroom and straight wires, a comparative study was done by Brandon owen [29]. measuring the force and moments in lingual appliances which concluded that straight wires produce larger forces and moments at the CR. This is because the large inter-bracket distance in the mushroom archwire between the canine and first premolar decreases the stiffness of the archwire and the mushroom shape keeps the bracket-archwire interface closer to the long axis of the teeth, which reduces the moments. Further the thicker bases in straight wires increase the distance of the bracket from the long axis of the tooth which increases the moment arm from the CR, hence producing greater moments. Lateral incisors, first premolars and canines are the teeth most affected by this difference in characteristics of the different archwires.

Mushroom archwires need a large inventory to be hosted by the orthodontist since the different mesiodistal dimensions of anterior teeth displaces the inset bend between canine and first premolar mesially or distally, therefore necessitating different sizes for individual patients. With the straight archwire, the same archwire shape can be used in all the patients for any malocclusion.

Not only it is easier and faster to work with straight archwire but also the mushroom-shaped archwires need arduous, still precise wirebending practice for its construction [30]. which is laborious and time consuming for the orthodontist and if not given proper time, may produce unsatisfactory results as well.




Fig. 4. Standardized arches for the lingual technique: (a) mushroom shape and (b) straight

5. INSTRUMENTS IN LINGUAL ORTHODONTICS

Because of the difficulty in accessing the lingual surface of tooth, maneuverability and dexterity in performing any procedures cannot be achieved with conventional instruments so specially designed instruments have been developed by ETM corporation with longer handles and angulated beaks either at 45°/90° [21].

1. Lingual Ligature Cutter- Angulated 45°(Fig. 5.a)
2. Light ligature plier, Mathieu style plier (Fig. 5.b)
3. Kurz First order bending fork. (Fig. 5.c) it can place first order bends directly in the

mouth, thus eliminating the need to remove wire and ligatures.

4. Kurz Archwire Cutter (Fig. 5.d) its function is same as that of distal end cutting plier.
5. Kurz Mosquito Forceps: curved at 45° (Fig. 5.e) used for attaching elastics and elastic chains.
6. Kurz Ligature wire cutter (Fig. 5.f) angulated at 90°.
7. Direct bond removing Plier (Fig. 5.g)
8. Kurz tongue retractor and Saliva ejector. (Fig. 5.h)
9. Kurz Utility Plier (Fig. 5.i) same as Weingart utility plier.
10. Kurz Second Order Bending Fork. (Fig. 5.j)
11. Module Remover (Fig. 5.k)



Fig. 5.a




Fig. 5.b



Fig. 5.c



Fig. 5.d



Fig. 5.e



Fig. 5.f



Fig. 5.g



Fig. 5.h



Fig. 5.i



Fig. 5.j



Fig. 5.k

6. INDIRECT BONDING AND LABORATORY SETUP IN LINGUAL ORTHODONTICS

Direct bonding of brackets as is done routinely in labial fixed orthodontics, is not a viable option in case of lingual orthodontics since the anatomical diversity of the lingual surface morphology i.e. differences in labio-lingual thickness, height of cingulum, less interbracket space etc. does not allow accurate and precise positioning of brackets onto the lingual surface under direct view. Hence indirect bonding procedures in the laboratory setting have been devised to ensure exactitude in bracket positioning. Furthermore advancements in Laboratory equipment and technology has made it possible to design and produce brackets and archwires and other appliances completely customized for individual patient needs [21,31,32].

Broadly laboratory setups work in two main ways in customization sector, A manual setup working upon patient's dental models and includes systems like BEST, CLASS, and HIRO, etc.,

The second being a completely customized digital lingual setup e.g. Orapix, WIN, HARMONY, Incognito™ and Lingual Matrix, and iLingual III D, which uses computer aided designing (CAD) and computer aided machining(CAM) to scan models or three dimensional images of the patients' dentition to manufacture brackets and archwires and

provides detailed description of analysis of dentition and biomechanical aspects specific for that particular patient.

6.1 Systems with Manual Setup

6.1.1 Torque Angulation Reference Guide (TARG)

Launched by the Ormco Company in 1984 the TARG machine provides a specific Tip and Torque to each bracket which are bonded on to a malocclusion model via medium of a virtual set up, by putting additional composite at the base of each bracket.

6.1.2 Bonding with equal specific thickness (BEST)

Didier Fillon invented Electronic TARG in 1986 which was more accurate compared to original TARG [31]. and DALI (dessin de l'arch linguale informatise) which is an arch wire tracing produced by a computer. The most critical problem in lingual orthodontics that is placing brackets according to the variation in the labiolingual thickness of different teeth, also present in original TARG, was eliminated here with a precise measuring device which provided compensation by enabling precise measurement of the labiolingual thickness of each tooth and the amount of composite needed under the base of the bracket.



Fig. 6 a. Torque Angulation Reference Guide (TARG)




Fig. 6 b. Labial reference gauge for placing brackets (TARG)



Fig. 6c. The TARG machine has several blades, making it possible to prescribe the torque and angulation for each individual tooth




Fig. 7. The Electronic TARG has a precise measuring device that measures the distance (thickness) between the labial surface of the tooth and the slot of the bracket

6.1.3 Customized lingual appliance set up service system (CLASS) [21,31]

The CLASS technique was described by Scott Huge. In this system an ideal diagnostic model is prepared from the malocclusion model of the patient. Using this ideal setup as a template, brackets are bonded onto the malocclusion model keeping in mind to produce an ideal configuration of teeth. Rest of the procedure to prepare transfer trays and the indirect bonding method to transfer the brackets clinically in the mouth is similar to other manually managed indirect bonding methods.

6.1.4 Korean indirect bonding set up (KIS) [33]

Tae Weon Kim was the founder of the Korean Society of Lingual Orthodontics (KSLO). He contrived an indirect bonding system that included a Model Checker, a bracket positioner,

and CRC Ready-Made Core Trays [34]. which were plastic preformed attachments, perfectly conforming to the external surface of the Ormco lingual bracket [35]. Dr. Kim also designed special pliers to be used exclusively in the lingual technique. The model checker confirms the accurate and precise positioning on the setup model.

6.1.5 Slot machine [14]

Designed by Thomas Creekmore, the slot machine helps in conveniently placing both labial and lingual brackets directly on the malocclusion model without the need for an ideal model setup.. The bracket slot horizontal or vertical, is oriented to the machine.

6.1.6 The Lingual bracket jig

This is a ruler developed by Geron. This measurement device greatly facilitates easy and

convenient placing of brackets by either direct or indirect means of bonding. This ruler consists of a universal jig meant for measurement on the posterior teeth and a pack of six different jigs for the measurement of six individual maxillary anterior teeth.

6.1.7 HIRO system

HIRO System was named after its inventor Toshiaki Hiro [36]. and was later adapted and improvised by Kyoto Takemoto and Giuseppe Scuzzo. There is no need for any special equipment. First of all sectioning of teeth is carried out from the malocclusion model and then teeth are arranged according to the ideal alignment anticipated. Archform is prepared using a rigid rectangular arch wire and bending it as per the anatomic contours and positioning the brackets according to it. Transfer trays are fabricated using the Convertible resin core system which in itself manipulates to precisely reposition the brackets.

6.1.8 Hybrid core system

The Hybrid Core System was developed by Matsuno [37]. with the objective of devising a technique for the construction of transfer trays and transfer of brackets. The brackets are firstly covered by silicone and then composite which results in biologically sound trays which are not easily distorted during transfer of brackets onto the teeth. The transfer trays are conveniently

removed from the mouth as silicone easily gets separated from the brackets after bonding.




Fig. 8. a. Lingual Bracket JIG (LBJ) consists of a set of six jigs for the maxillary anterior teeth (canine to , ,canine)which present the most morphologic variations on the lingual side, a millimeter ruler (0.1-mm accuracy),and a wrench for in-out adjustments. The ruler is used to measure the in-out bracket position, compensating f for differences in thickness. The LBJ was used in both direct and indirect bonding techniques (B) High magnification of a jig




Fig. 9. HIRO system

6.2 Completely Digital Customization

6.2.1 Top/Incognito I Braces system (Transfer Optimized Positioning)

Invented by Dr Dirk Wiechmann, *Lingual braces* as they are commonly known, the Incognito I braces are constructed using the CAD-CAM technology which enables complete customization of brackets according to individual case and even archwires are prefabricated with all the features inbuilt according to the biomechanical needs in a particular patient.

The Transfer Optimized Positioning procedure has 4 major characteristics: [38].

1. The brackets are bonded on the malocclusion model.
2. The positioning is done with the TARG Professional.

3. The three-dimensional orientation of the malocclusion model.
4. All brackets are bonded with minimal positioning thickness.

6.2.2 Orapix system

The Orapix system is the latest and most beautiful flower in the bouquet of lingual orthodontics. There is a digital scanner that scans the patient's model or the patient's dentition intraorally three dimensionally and transforms it into a three dimensional data file which consists of multiple well integrated images taken from each and every angle by the scanner. The orthodontist then makes a ideal virtual set up on the computer using the digital model dictating the parameters that are required in a particular case. Based on the information provided by the virtual setup, orthodontist formulates his diagnosis and treatment plan deciding upon the biomechanics required to produce the desired occlusal adjustments [39].






Fig. 10. Conventional lingual bracket (left) and customized bracket




Fig. 11. Incognito Braces



A. Virtual arch segmented into individual units




B. visualizing the contact points



C. Placement of brackets on the virtual set-up and arch form of the straight wire



D. Bonding of a bracket with its jig on the model



E. Automated calculation by 3Txxer software of the slot-vestibular surface distance



F. Bracket bonding using a Memosil transfer tray



G. Individual tooth bonding using positioning extensions



H. Bonded brackets using the Kommon Base technique

Fig. 12. Orapix lingual straight wire technique

According to the diagnosis and treatment planning, the digital software designs the transfer trays and a rapid prototype (RP) machine prepares the transfer trays in resin. Then the technician positions the brackets in the transfer trays and the resin pads on the base of the brackets to complete the lab. work.

6.3 Direct Bonding/ Simplified Technique

Dr. Michael Diamond in 1984 designed a device called as Peri/Reflector which was composed of a mirror, tongue retractor, and saliva ejector.



Fig. 13. Peri/Reflector

The Simplified technique uses the STB brackets and a bracket placement plier and simple tweezers for bracket positioning at a distance of 1.5 to 2mm from the incisal edges of anterior teeth [40].

7. ADVANTAGES AND DISADVANTAGES OF LINGUAL ORTHODONTICS

Esthetics and absolute invisibility of the Appliance is its foremost and paramount advantage when compared to the labial fixed orthodontic appliances, along with its biomechanical efficiency in managing the non-extraction cases and with the advancement of the technique the most difficult cases involving extraction and retraction of anterior teeth can be maneuvered without any difficulty [32]. In addition there are no chances of damage or decalcification of the labial enamel surface which is one of the problems with the labial appliances during or at the completion of orthodontic treatment. There is significant intrusion of the anterior teeth and some extrusion of posterior teeth which becomes very useful in case of reduction of deep bites especially the severe ones [24]. as the position of bracket slot which engages the archwire is very close to the centre

of resistance of the anterior teeth in case of lingual brackets, there is more planned movement of the desired teeth and other unwanted factors as anchorage loss and angulation and torque changes are avoided while attempting space anterior teeth retraction during space closure and also during bite opening [41].

Disadvantages primarily include a lack of direct vision which hampers direct bonding technique and there arises a need for Indirect bonding technique for accurate bracket placement.

[42]. Other problems are occlusal interferences causing frequent bond failures, Speech Distortion, lacerations to the tongue, Gingival irritation, difficult oral hygiene, white spot lesions and increased chair side time and high cost.

8. INDICATIONS FOR LINGUAL APPLIANCES

There are a few situations where lingual brackets come advantageous because of their inbuilt design features and distinct mechanical characteristics such as a bite plane, occlusally opening bracket slots, mushroom shaped arch wires etc.

1. Intrusion of Anterior teeth : The intrusive force in case of lingual brackets passes through the centre of resistance of the teeth as the brackets are placed more close to the centre of resistance on the lingual surface because of the morphological contours and also lower anterior teeth touch the horizontal plane of the lingual brackets. Both these factors produce a bite plane effect on upper teeth effecting light, continuous, forces and hence this force system becomes very effective in intrusion of teeth and therefore deep bite reduction.
2. Expansion of the Maxillary Arch: as the lingual brackets are located between the tongue and lingual/palatal aspect of the teeth, a continuous buccal/labial force is exerted by the tongue over the teeth which is centrifugal and expansile in nature. This force is further sustained and supported by the mushroom shape of the archwires and the reduced interbracket distance.
3. Repositioning of the mandible with orthodontic tooth movements: In the first stage of treatment any temporomandibular problems and pain/related symptoms are sorted out because of the bite plane effect

of the lingual brackets and in the following clinical stage, a new mandibular position is acquired because of the change in occlusion which further stabilizes the occlusion.

4. Distalisation of maxillary molars: molar distalisation through lingual technique is more efficient producing more bodily movement and less distal tipping [21].
5. Cases which are complicated by an existing habit of forceful tongue thrust because of the tongue restriction due to lingually placed brackets.

Ideal Cases [43] for lingual orthodontics include Deep bite in horizontally growing individuals, Class I malocclusion with slight crowding, Class II Division 2 malocclusion, extraction cases with class II malocclusion, midline diastema, Pre-Tooth movement required for ideal alignment in case of planned fixed Prosthesis and Class III malocclusion and others needing Surgery.

Some difficult situations to handle with lingual orthodontics are 1. premolar extraction cases 2. Bilateral/unilateral Crossbites of the posterior teeth. 3. High mandibular plane Angle (vertical facial pattern) cases and 4. Open bite cases [22].

9. COMPARISON OF LINGUAL AND LABIAL APPLIANCES

Scuzzo and Takemoto known for their significant contribution to the lingual orthodontics, took on a study to compare the different orthodontic forces acting on the lingual and labial brackets in all 3 dimensions [21,22]. They summarized the major findings as following:

In the vertical plane the most perceptible and immediate effect is opening of the bite since the horizontal plane of the lingual brackets inferiorly touches with the incisal edges of the mandibular teeth and cause intrusion of the maxillary and mandibular anterior teeth and slight extrusion of the posterior teeth which produces the deep bite reduction effect. Bite opening is advantageous in brachyfacial patients and TMJ cases and tooth movements occur speedily because of posterior disclusion, but in faces with vertically growing and average growth pattern, there is a tendency of increased anterior and posterior vertical height in already long faces due to the bite opening effect. This increased facial height produces anterior open bite, mouth breathing etc., the kind of problems which are difficult to manage clinically once the growth pattern is disturbed due to faulty biomechanics.

Since the lingual brackets are positioned in the center of cingulum there are lesser moments of forces acting on the tooth in vertical, sagittal and horizontal planes when compared to the labial brackets because the distance from the centre of brackets to the centre of resistance of tooth becomes shorter and hence arm of the moment becomes small. Therefore when intrusive forces are applied, the labial vector does not produce labial tipping in normally inclined and already protruded teeth. But in case of retroclined maxillary anterior teeth a clockwise moment is produced by the lingual brackets which increases the lingual inclination of the already retroclined teeth whereas vice versa happens with the labial brackets because of their position on the labial surface [22].

Effect of extrusive forces has also been studied. Extrusive forces in lingual appliance have been examined by Geron and Romano. They concluded that extrusive forces produced labial root movement only when there was more than 20 degrees proclination of the maxillary incisors. Otherwise there is lingual root movement only. Conversely it was observed that there were only labial root movements only with the labial brackets [44].

Anteroposterior plane since there is backward and downward rotation of mandible immediately after appliance installation, due to bite opening, the lingual appliance produces a Class II tendency.

Due to posterior disocclusion with bite opening, Anteroposterior movement of molar becomes easily achievable. Molar anchorage is also secured firmly since there is routine banding of the second molars [42,22].

On application of same amount of retraction force, as it becomes equal to the intrusion force

equals, the net force is directed through the center of resistance in the labial technique, but remains lingual in the lingual orthodontics which produces lingual tipping and a bowing effect vertically similar to the roller coaster effect produced in labial technique with excessive retraction forces (Fig. 14).




Fig. 14. Comparison between the anterior retraction in labial and lingual mechanics

So to avoid these side effects during en masse retraction in lingual orthodontics, the applied force must be within the biological limits of retraction and forces must be concentrated on intrusion vector producing more lingual root torque [21,22].

In the transverse plane the lingual appliance produces forces that tend to promote expansion of the arch, which together with posterior disclusion, cause mesiobuccal molar rotation when spaces are closed using retraction forces. Therefore anchorage in form of transpalatal arches is critical. Retraction is always done on stiffer wires to resist the "bowing effect" due to excessive forces or the failure to manage the force vectors due to the lingual nature of the appliance, both in transverse and vertical plane (Fig 15. a & 15.b) [21,42].



Fig. 15.a. Transverse bowing and mesiobuccal forces




Fig. 15.b. Vertical bowing effect of retraction rotation of molars

10. CONCLUSION

Lingual orthodontics, inspite of being technique sensitive and a demanding orthodontic modality in terms of expertise, special bracket and wire system and complicated system of wire bending , need for indirect bonding and customization of treatment plan; has definitely emerged as the most desirable treatment option in the recent scenario because of its most acceptable form of invisible braces ,especially in adults and adolescents or more appropriately young adults due to the elimination of the feeling of ignominy or the social barrier that orthodontic treatment represented in the past. This, along with improvisation of bracket system and techniques and the research in the coming years, will sure shot be a boon in the hands of skilled orthodontists.

CONSENT AND ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Meier B, Weimer KB, Miethke RR. Invisalign patient profiling. *J Oro Fac Ortho.* 2003;64:352–8.
2. Freitz U, Diedrich P. Weichmann-lingual technique- patients characteristics motivation and acceptance. Interpretation of a retrospective survey. *J Oro Fac Ortho.* 2002;63:227–33.
3. Profitt WR. Treatment for Adults. In:Profitt WR, Contemporary Orthodontics. 4 th edition. St Louis: Mosby Elsevier; 2007.
4. Echarri P. Revisiting the history of lingual orthodontics: A basis for the future. *Semin Orthod* 2006;12:153–59.
5. Gupta A, Kohli V, Hazarey PV. Lingual orthodontics – A review. *J Ind Orthod Soc.* 2005;38:46-54.
6. Kharbanda OP. Orthodontics: Diagnosis and management of malocclusion and dentofacial deformities. 3rd edition. New Delhi: Elsevier health sciences; 2019.
7. Shetty V, Shetty SV, Sarje S. Lingual orthodontics - A review. *Ind J Orthod Dentofac Res.* 2020; 6(2): 44–50
8. Asif Chatoo. A view from behind: a history of lingual orthodontics. *J O.* 2013;4:S2–S7.
9. Echarri P. Revisiting the history of lingual orthodontics: A basis for the future. *Semin Orthod* 2006;12:153-9.
10. Fujita K. Development of the lingual brackets technique. A hygienic and aesthetic approach to orthodontic treatment. *Orthod Clín.* 2001;4(3):176-180.
11. Echarri P. Lingual Orthodontics. Complete technique, step by step. Barcelona, Nexus Ediciones; 2003.
12. Mizrahi E. Lingual orthodontics using the Light Wire technique Part 1. Available:www. Lingualnews.com May 2004;2:1.
13. Mizrahi E. Lingual Orthodontics using the Light Wire technique Part 2. Available:www. Lingualnews.com May 2004;2:2.
14. Creekmore T. Dr. Thomas creekmore on lingual treatment mechanics. *JCO Interviews,* Nov. 1996;30(11):46 1-476.
15. Fillion D. The orapix system. In: Scuzzo G, Takemoto K, editors. Lingual orthodontics: A new approach using STb light lingual system and lingual straight wire. Chicago: Quintessence Publishing; 2010.
16. Scuzzo G, Takemoto K, Takemoto Y, Scuzzo G, Lombardo L. A new self-ligating lingual bracket with square slots. *J Clin Orthod.* 2011;45:682-90.
17. Kairalla SA, Scuzzo G, Triviño T, Velasco L, Lombardo L, Paranhos LR. Determining shapes and dimensions of dental arches for the use of straight-wire arches in lingual technique. *Dent Press J Orthod.* 2014; 19:116-22.
18. Park KH, Bayome M, Park JH, Lee JW, Baek SH, Kook YA. New classification of lingual arch form in normal occlusion using three dimensional virtual models. *Korean J Orthod.* 2015;45:74-81.
19. Grist F. Basic guide to orthodontic dental nursing. John Wiley and Sons. 2010; 155–65.
20. Vasumurthy S, Saritha T. Lingual orthodontics. *Ind J Dent Asso.* 2009;1(1):31-37.
21. Rafi Romano. Lingual orthodontics, first ed. Becker & CO; 2001.
22. Lily MY, Shum et al. Lingual orthodontics - A review. *Hong Kong Dent J.* 2000;1:13-20.

23. Hong RK, Wook Sohn H. update on the fujita lingual brackets. J Clin Orthod. 1999;33:136–42.
24. Reddy A, et al. Indirect bonding with various systems in lingual orthodontics. Annals and Essence of Dentistry. 2012;4(2):98.
25. Scuzzo G, Takemoto K. Invisible orthodontics. Current concepts and solutions in lingual orthodontics. Berlin: Quintessenz Verlags GMBH; 2003.
26. Aldo Macchi et al. Philippe: Self-ligating lingual brackets. J Clin Orthod. 2002;36(1):42-45.
27. Ronchin M. Resolving a Class II malocclusion with molar distalisation. Pract Perio Aesthet Dent. 1994;6:51–8.
28. Navarro. Development of the in ovation L bracket from GAC. Lingual News. 2006;4:2.
29. Owen B, Gullion G, Heo G, Carey JP, Major PW, Romanyk DL. Measurement of forces and moments around the maxillary arch for treatment of a simulated lingual incisor and high canine malocclusion using straight and mushroom archwires in fixed lingual appliances. Euro J Orthod. 2017;39(6):665–672.
30. Hernández-Vázquez RA, Marquet-Rivera RA, Mastache-Miranda OA, Vázquez-López AJ, Cruz-López S, Vazquez feijoo JA. Comparative numerical analysis between two types of orthodontic wire for the lingual technique, using the finite element method. App Bio Biomech. 2021;438:1-15.
DOI: 10.1155/2021/6658039.
31. Buso-Frost L, Fillion D. An overall view of the different laboratory procedures used in conjunction with lingual orthodontics. Semin Orthod. 2006;12(3):203–7.
32. Grist F, Rdn. RDN Fixed appliances – Indirect Bonding and Lingual Orthodontics. Blackwell Publishing Limited; 2010.
33. Kim TW. Indirect bonding system in lingual orthodontics. Kor J Orthod. 2002;32:38-45.
34. Echarri P, Kim TW. Double transfer trays for indirect bonding. J Clin Orthod. 2004;38:8-13.
35. Echarri P. Lingual orthodontics: Patient selection and diagnostic considerations. Sem Orthod. 2006;12(3):160– 166.
36. Hiro T, Iglesia FDL, Andreu P. Indirect bonding technique in lingual orthodontics: the HIRO system. Prog Orthod. 2008;9(2):34-55.
37. Matsuno I, Okuda S, Nodera Y: The hybrid core system for indirect bonding. J Clin Orthod. 2003;37:160-161.
38. Wiechman D. Lingual Orthodontics (part 1): laboratory procedure. J Orofac Orthop. 1999;60: 371-379.
39. Fillion D. Clinical advantages of the Orapix-straight wire lingual technique. Int Orthod. 2010;8: 125-151.
40. Komori A, Morth RCS, Takemoto K, Shimoda T, Miyashita P, Scuzzo G. Precise direct lingual bonding with the kommon base. J Clin Orthod. 2013;47(1):42-9.
41. Poon KC, Taverne AAR. Lingual orthodontics: A review of its history. Aust Orthod J. 1998;15(2):101- 104.
42. Salloum RB, AlRassan L, Almukhaizeem N, Albluwi A, Alfayez R, Shammery DA. Lingual Orthodontics Appliance, An Overview. Int J Oral Dent Health. 2018;4(2): 1-2.
43. Prabhakar R, Vikram NR, Saravanan R, Karthikeyan MK, Rajesh R. Lingual orthodontics. IJMH. 2012;1:16–8.
44. Geron S, Romano R, Brosh T. Vertical forces in labial and lingual orthodontics applied on maxillary incisors-a theoretical approach. Angle Orthod. 2004;74(2):195-201.

© 2022 Sharma and Soni; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/93182>