



# Screening for Biodegradation Potential of Endophytic Bacteria Isolated from the Roots and Leaves of Mangrove Plants (*Avicennia germinans*) (Black Mangrove), *Acrostichum aureum* (Golden Leather Fern) and *Rhizophora mangle* (Red Mangrove)

Somiari, Adaobi Assumpta Abiye <sup>a\*</sup>, Douglas, Salome Ibietela <sup>a</sup> and Nrior, Renner Renner <sup>a</sup>

<sup>a</sup> Rivers State University, Nigeria.

## **Authors' contributions**

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## **Article Information**

DOI: 10.9734/JAMB/2022/v22i730471

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/85551>

Original Research Article

Received 04 February 2022  
Accepted 06 April 2022  
Published 12 May 2022

## **ABSTRACT**

**Aim:** The aim of this study is to screen and determine the biodegradation potential of the endophytic bacteria isolated from roots and leaves of mangrove plants.

**Methodology:** In this work were isolated, identified and screened endophytic bacteria from roots of *Rhizophora mangle*, *Avicennia germinans* and *Acrostichum aureum*; the three major species of mangrove plants found in the Niger Delta. The roots were transported in a sterile bag to the Microbiology Laboratory and treated using standard Microbiological techniques. The organisms isolated include: *Pseudomonas* sp, *Bacillus* sp, *Staphylococcus* sp, *Micrococcus* sp, *Klebsiella* sp, *Azotobacter* sp, *Nitrobacter* sp. and *Nitrosomonas* sp. These organisms were screened for their ability to degrade crude oil by analyzing them using an ultra-violet spectrophotometer at a wavelength of 600nm and through a colorimetric test that involves the use of 2,6, Dichlorophenol-indophenol (DCPIP) as the metabolic activity indicator for a 14 days period.

**Results:** According to the colorimetric test, isolates that were positive hydrocarbon degraders were observed by a colour change from blue to colourless, which include: *Pseudomonas* sp, *Staphylococcus* sp, *Bacillus* sp, *Klebsiella* sp and *Nitrobacter* sp, while those that changed from blue to pink (an indication of a negative reaction) were: *Nitrosomonas* sp, *Bacillus* sp, *Micrococcus* sp, *Staphylococcus* sp, *Azotobacter* sp and *Klebsiella* sp. Amount degraded and percentage biodegradation ranged from *Micrococcus* sp 278.6 (6.2%) < *Azotobacter* sp. 1242.8 (17.8%) < *Nitrosomonas* sp 1392.9 (19.9%) < *Staphylococcus* sp 1543.0 (22.1%) < *Bacillus* sp 17927.8 (25.6%) < *Nitrobacter* sp 1935.7 (28%) < *Klebsiella* sp 3392.9 (48.5%) < *Bacillus* sp 3671.4 (52.5%) < *Pseudomonas* sp 4942.8 (70%).

**Conclusion:** *Pseudomonas* and *Bacillus* species demonstrated the highest potential to degrade crude oil and can therefore be used in bioremediation of water ecosystem impacted by crude oil. Hence, the scientific contribution of this research is related to identifying several culturable groups of bacteria that might be directed to these further biotechnological approaches.

**Keywords:** Endophytic bacteria; mangrove roots; biodegradation.

## 1. INTRODUCTION

Mangrove ecosystems are of great ecological and economic importance as they play various vital roles at the land-sea interface, provide food, breeding grounds and nursery sites for a variety of terrestrial and marine organisms, used for human sustainability and livelihoods (food, timber, fuel and medicine), and they also offer protection against catastrophic events, such as tsunami, tropical cyclones and tidal bores and can dampen shoreline erosion [1].

Mangrove forests occur at the interface of terrestrial and marine ecosystems portraying a rich biological diversity of plants, animals and microorganisms. Microbes are important part of the mangrove environment as they contribute to the productivity of the mangrove ecosystem. They play a very critical role in creating and maintaining this biosphere and also serve as a source of biotechnologically valuable and important products by participating in various steps of decomposition and mineralization of leaf litter; by being able to recycle nutrients, produce and consume gases that affect global climate, destroy pollutants, treat anthropogenic wastes and can also be used for biological control of plant and animal pests. Microorganisms from mangrove environments are a major source of antimicrobial agents and also produce a wide range of important medicinal compounds, including enzymes, antitumor agents, insecticides, vitamins, immunosuppressants and immune modulators [2].

Endophytes are microorganisms that live in plant tissues partly or all their lifecycle without causing any symptoms of disease in the host [3]. These microbes are often bacteria or fungi, found in

various plant organs such as seeds, roots, stems, leaves, flowers and fruits that colonize host tissues similar to a pathogen. Studies have shown that the endophytic association with the host contributes significantly to accelerated seedling emergence, enhanced plant growth, improved resistance against various phytopathogens and abiotic stresses. Endophytic microorganisms can safeguard the hosts against numerous biotic and abiotic factors such as attack of insects, pathogens and herbivores [4].

Various groups of bacteria are found in the mangrove ecosystem where they are known to perform diverse activities including photosynthesis, nitrogen fixation and methanogenesis [5,6,7]. The over-exploitation of mangroves is of today's concern because it threatened the sustainability of the ecosystems. Microorganisms from mangrove ecosystems contain useful enzymes, proteins, antibiotics and salt-tolerant genes, all of which have biotechnological significance [2]. Little is known about bacterial mangrove communities, but these microorganisms may have high biotechnological potential [8]. Recently developed technologies in molecular biology and genetics offer great promise to explore the potential of microbial diversity. The objective of this study is to isolate and screen the endophytic bacteria from the roots of mangrove plants within a crude oil-polluted environment and to determine their biodegradability potential.

## 2. MATERIALS AND METHODS

### 2.1 Description of Study Area

The roots of mangrove plants used in this study were collected from the Creek Road waterfront (Bonny Jetty) in Old Port Harcourt Township,

Rivers State, Nigeria. The area has lots of mangrove species and also was an area that witnesses lots of pollution from wastes and activities from speedboats transporting passengers, carpenters building boats, boats transporting wood, diesel, fuel and even crude oil from illegal oil refineries as their products are sometimes discharged, dumped or even burnt in the river. The location is situated at Longitude 4<sup>0</sup> 45' 10.13976"N and Latitude 7<sup>0</sup> 14.13012 E.

## 2.2 Collection of Materials

The plants collected were placed separately in sterile polyethylene bags and transported first to Department of Plant Science & Biotechnology Laboratory for identification of the plants and then to Microbiology laboratory, of the Rivers State University, Port Harcourt, for Microbiological analysis.



Image 1. *Acrostichum aureum* (golden leather fern)



Image 2. *Rhizophora mangle* (Red mangrove)



Image 3. *Rhizophora racemosa* leaves



Image 4. *Rhizophora racemosa* roots



Image 5. *Acrostichum aureum* leaves



Image 6. *Acrostichum aureum* roots



Image 7. *Avicennia germinans* leaves



Image 8. *Avicennia germinans* roots

### 3. MEDIA USED

Nutrient Agar (TM Media)(g/L);Beef extract (1.50), Yeast extract (1.50), Peptic digest animal tissue (5.00), Sodium chloride (5.00), pH 7.4±0.2 at 25°C.; Phosphate Buffered Saline (g/l);Na<sub>2</sub>HPO<sub>4</sub> (1.44), Sodium Chloride (3), Potassium Chloride (0.2), KH<sub>2</sub>PO<sub>4</sub> (0.24); Mineral Salt Agar (g/l); K<sub>2</sub>HPO<sub>4</sub> (0.5), MgSO<sub>4</sub>·7H<sub>2</sub>O (0.3), NaCl (0.3), MnSO<sub>4</sub>·H<sub>2</sub>O (0.02), FeSO<sub>4</sub>·6H<sub>2</sub>O (0.03), NaNO<sub>2</sub> (0.03), Agar (0.5), Fungusol, Distilled water (200ml).; Burks-N-free Agar; Glucose (10), K<sub>2</sub>HPO<sub>4</sub>(0.41), K<sub>2</sub>HPO<sub>4</sub> (0.52), Na<sub>2</sub>SO<sub>4</sub> (0.05), CaCl<sub>2</sub> (0.2), MgSO<sub>4</sub>·7H<sub>2</sub>O (0.1), FeSO<sub>4</sub>·7H<sub>2</sub>O (0.005), Agar (15.0), Distilled water (1000ml); Winogradsky (*Nitrosomonas*); (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> (2.0), K<sub>2</sub>HPO<sub>4</sub> (1.0), MgSO<sub>4</sub>·7H<sub>2</sub>O (0.5), NaCl (2.0), FeSO<sub>4</sub>·6H<sub>2</sub>O (0.4), CaCO<sub>3</sub> (0.01), Agar (15.0), Distilled water (1000ml); Winogradsky (*Nitrobacter*); KNO<sub>2</sub> (0.1), Na<sub>2</sub>CO<sub>3</sub> (1.0), NaCl (0.5), FeSO<sub>4</sub>·7H<sub>2</sub>O (0.4), Agar (15.0), Distilled water (1000ml).

#### 3.1 Sample Preparation/Processing

The roots of the mangrove plants were treated to obtain only the endophytic bacteria. The roots were first washed under running water thoroughly to remove surface adhering debris. They were washed with sterile distilled water for 5 minutes, surface-sterilized with 70% ethanol for 1 minute, then with 3% sodium hypochlorite (NaOCl solution) for 3mins and rinsed 6 times in sterile distilled water in different containers. Thereafter, they were cut into small pieces, and grounded separately with already sterilized mortar and pestle to make slurries. Ten-fold serial dilution of the plant slurry was made up to 10<sup>6</sup> dilutions. Thereafter, an aliquot of 0.1ml of the appropriate dilutions was inoculated aseptically into different properly dried media

plates; Burk's N-Free Agar(10<sup>3</sup>), Winogradsky Agar (10<sup>3</sup>) (for both *Nitrosomonas* and *Nitrobacter*), Mineral Salt Agar (10<sup>3</sup>) and Nutrient Agar (10<sup>6</sup>). Using vapour phase diffusion method, filter papers were soaked with 2ml of crude oil and placed on the cover of the plate containing Mineral Salt Agar (MSA). The cultured plates were incubated at 37°C for 24hours for the nutrient agar, the modified Winogradsky by [6] used was incubated at 37°C for 3days while the previous Winogradsky agar was incubated for 5-7days and finally, the Mineral Salt Agar was incubated at same 37°C for 5-7days. After incubation, bacterial colonies were differentiated and counted based on their morphological characteristics. Individual colonies were picked randomly and sub-cultured by streaking them onto nutrient Agar plates using the streak plate technique and incubated at 37°C [9,6].

#### 3.2 Screening of Isolates

The microbial isolates were screened twice. Overnight pure cultures were used for the screening. Broth of the organisms was first prepared by transferring the pure culture into properly labeled test tubes containing normal saline (that has been autoclaved and allowed to cool and agitated).

The method described by [10] was applied, 9ml of MSA (Mineral salts agar), tween 8 (surfactant), Dichlorophenol-indophenol (indicator), and 0.5ml of crude oil were put into test tubes. After capping, all the test tubes were autoclaved at 121°C at 15PSI for 15minutes, then allowed to cool and labeled, 0.5ml of the broth (standardized bacterial cell suspension of the respective bacterial isolates) was transferred into the labeled test tubes containing crude oil. The test tubes which served as controls were not

inoculated. All test tubes were incubated for 14 days and their potential to utilize hydrocarbon was determined by observing the colour changes in the test tube on days 1, 7 and 14 and then each tube was analyzed for optical density (OD) using a spectrophotometer on days 1, 7 and 14 respectively to measure the amount of light absorbed by the solution so to determine the organism that has higher potential to degrade hydrocarbon as described by [11]. The optical density of the culture was measured at 600nm. The change in colour from blue to colourless was an indication that hydrocarbon degradation had occurred in those test tubes while those that changed from blue to pink were an indication of a negative result for hydrocarbon degradation. The endophytic bacteria with higher potential to degrade crude oil were used for the remediation study.

Three of the endophytic bacteria that showed greater potential to degrade crude oil were used for the remediation study.

The concentration of absorbance was calculated using the calibration curve for THC (Total hydrocarbon content) as adopted by [6,12] as follows:

$$Y = 0.28 + 0.0609X$$

Y = absorbance

X = mg/ml + conc of oil in xylene

$$X = \frac{Y - 0.0609}{0.28}$$

$$\text{THC} = X \times 1000 \times 0.5$$

Where X = concentration calculated from absorbance

1000 = Value used to convert from g to mg

0.5 = quantity of crude oil used equation 1

### 3.3 Percentage (%) Biodegradation Rate

This was calculated using the formula adopted by [12] as follows.

Step 1: the amount of pollutant remediated equals the initial concentration of pollutant (Day 1) minus the final concentration of the pollutant at the end of the experiment.

Step 2: percentage (%) bioremediation = the amount of pollutant remediated divided by initial concentration of pollutant (Day 1) multiplied by 100.

$$B_c = I_c - F_c$$

$$B_x = I_c - I_o$$

Where,

B<sub>c</sub> = Amount of pollutant degraded

I<sub>c</sub> = the initial concentration of pollutant (Day 0)

F<sub>c</sub> = the final concentration of the pollutant at end of the experiment

B<sub>x</sub> = Actual amount of the pollutant in the test medium

I<sub>o</sub> = the initial concentration of control of crude oil polluted water at Day 0

$$\% \text{Bioremediation} = \frac{B_c}{B_x} \times 100$$

Chart 1. Screening setup

Isolates	Volume of MSA (ml)	Volume of Crude oil (ml)	Volume Broth of Organisms (ml)	Total volume (ml)
Control	9	0.5 (0.45g)	0.5	10
<i>Pseudomonas aeruginosa</i> (MW369466)	9	0.5 (0.45g)	0.5	10
<i>Pseudomonas aeruginosa</i> (MN314747).	9	0.5 (0.45g)	0.5	10
<i>Micrococcus</i> sp	9	0.5 (0.45g)	0.5	10
<i>Azotobacter</i> sp	9	0.5 (0.45g)	0.5	10
<i>Staphylococcus</i> sp	9	0.5 (0.45g)	0.5	10
<i>Brevibacillus brevis</i>	9	0.5 (0.45g)	0.5	10
<i>Staphylococcus</i> sp	9	0.5 (0.45g)	0.5	10
<i>Bacillus</i> sp	9	0.5 (0.45g)	0.5	10
<i>Nitrosomonas</i> sp	9	0.5 (0.45g)	0.5	10
<i>Bacillus amyloliquefaciens</i>	9	0.5 (0.45g)	0.5	10
<i>Nitrobacter</i> sp	9	0.5 (0.45g)	0.5	10
<i>Klebsiella</i> sp	9	0.5 (0.45g)	0.5	10
<i>Klebsiella pneumonia</i>	9	0.5 (0.45g)	0.5	10

#### 4. RESULTS AND DISCUSSION

The mangrove plant samples used were identified in the Department of Plant Science & Biotechnology laboratory as: *Avicennia germinans* (Black mangrove), *Rhizophora mangle* (Red mangrove) and *Acrostichum aureum* (golden leather fern).

Eight (8) bacterial genera were isolated from the mangrove plant which include; *Pseudomonas* sp, *Micrococcus* sp, *Nitrobacter* sp, *Nitrosomonas* sp, *Staphylococcus* sp, *Azotobacter* sp, *Klebsiella* sp and *Bacillus* sp.

The results of the screening using a spectrophotometer to determine the organisms with the highest biodegradation potential are presented below in Table 1. According to the

results, *Pseudomonas aeruginosa* (MN314747) degraded the hydrocarbon most at a percentage (%) value of 71% (4942.8), followed by *Brevibacillus brevis* at 59% (4100), then *Bacillus amyloliquefaciens* at a value of 53% (3671.4), followed by *Klebsiella pneumoniae* at a value of 49% (3392.9), then *Pseudomonas aeruginosa* (MW369466 at 35% (2450), then *Klebsiella* sp with a degradation value of 34.1% (1957.1), followed by *Nitrobacter* sp with a value of 28% (1935.7), followed by *Bacillus* sp with a degradation value of 26% (1792.8), then *Staphylococcus aureus* with a degradation value of 21.1% (1473.1), then *Nitrosomonas* sp with a value of 20% (1392.9), followed by *Azotobacter* sp with a degradation value of 18% (1242.8), then *Staphylococcus* sp with a value of 17.3% (1210.7) while *Micrococcus* sp had the lowest biodegradation potential at a value of 6% (434.6).

**Table 1. Test for biodegradation potential**

Isolates	Day 1	Day 7	Day 14	Amount degraded (mg/L)	% degraded
Control	6993.6	5672.1	5429.3	1564.4	23.5%
<i>Pseudomonas aeruginosa</i> (MW369466)	4993.6	3036.4	2543.6	2450	35%
<i>Pseudomonas aeruginosa</i> (MN314747).	6036.4	3050.7	1093.6	4942.8	71%
<i>Micrococcus</i> sp	4122.2	3842.9	3687.6	434.6	6%
<i>Azotobacter</i> sp	4529.2	4065	3286.4	1242.8	18%
<i>Staphylococcus</i> sp	5978.5	5075.6	4767.8	1210.7	17.3%
<i>Brevi bacillusbrevis</i>	5993.6	4865	1893.6	4100	59%
<i>Staphylococcus</i> sp	5922.7	4927.5	4449.6	1473.1	21.1%
<i>Bacillus</i> sp	5779.2	6350.8	3986.4	1792.8	26%
<i>Nitrosomonas</i> sp	6107.9	5229.3	3715	1392.9	20%
<i>Bacillus amyloliquefaciens</i>	6272.1	4315	2600.7	3671.4	53%
<i>Nitrobacter</i> sp	5908.5	4321.6	3951.4	1957.1	28%
<i>Klebsiella</i> sp	5707.9	4993.6	3322.1	2385.8	34.1%
<i>Klebsiella pneumoniae</i>	5186.4	6536.4	1793.5	3392.9	49%

**Table 2. Screening using surfactant and indicator**

Isolates	Reaction	Colouration
<i>Pseudomonas aeruginosa</i> (MW369466)	+ve	Colourless
<i>Pseudomonas aeruginosa</i> (MN314747).	+ve	Colourless
<i>Micrococcus</i> sp	-ve	Pink
<i>Azotobacter</i> sp	-ve	Pink
<i>Staphylococcus</i> sp	-ve	Pink
<i>Brevi bacillusbrevis</i>	+ve	Colourless
<i>Bacillus</i> sp	-ve	Pink
<i>Staphylococcus aureus</i>	+ve	Colourless
<i>Nitrosomonas</i> sp	-ve	Pink
<i>Bacillus amyloliquefaciens</i>	+ve	Colourless
<i>Nitrobacter</i> sp	+ve	Colourless
<i>Klebsiella</i> sp	-ve	Pink
<i>Klebsiella pneumoniae</i>	+ve	Colourless

#### 4.1 Screening Using Surfactant and Indicator

For the screening using tween 80 (surfactant) and Dichlorophenol-indophenol (indicator), the results are presented in Table 2; the colour change from blue to colourless indicates a positive result of hydrocarbon degradation and those that changed from blue to pink is an indication of a negative result. According to the results, *Pseudomonas aeruginosa* (MN314747), *Brevibacillus brevis*, *Bacillus amyloliquefaciens*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Nitrobacter* sp and *Staphylococcus aureus* were positive as they turned colourless by day 14, while *Azotobacter* sp, *Staphylococcus* sp, *Micrococcus* sp, *Bacillus* sp and *Nitrosomonas* sp turned pink by day 14 showing a negative result.

#### 4.2 Evaluating Hydrocarbon Utilizing Potential of the Endophytic Bacteria

Screening of the isolates was done using two different methods to evaluate their hydrocarbon utilizing potential. The readings from the spectrophotometer were used to determine the isolates with the highest biodegradation potential by subtracting the reading of Day 14 from the reading on Day 1 using the formula adopted by [6]. All the isolates showed the ability to degrade hydrocarbon as it has been reported that more than 79 bacterial genera e.g. *Pseudomonas*, *Staphylococcus*, *Acinetobacter*, *Streptococcus*, *Enterobacter*, etc have been found to play vital roles in petroleum hydrocarbon degradation [13,14,15,16]. According to the results, the amount degraded and percentage degradation rate are shown in Figs. 1 and 2 respectively. The

rate of crude oil degradation is as follows: *Pseudomonas aeruginosa* (MN314747), which showed the highest hydrocarbon degradation potential with a value of 4942.8mg/l at 71% > *Bacillus amyloliquefaciens* with a value of 4100mg/l at 59% > *Brevibacillus brevis* with a value of 3671.4mg/l at 53% > *Klebsiella pneumoniae* with a value of 3392.9mg/l at 49% > *Pseudomonas aeruginosa*(MW369466) with a value of 2450mg/l at 35% > *Nitrobacter* sp with a value of 1957.1 at 28% > *Bacillus* sp 1792.8 (26%) > *Staphylococcus* sp 1473.1 (21.1%) > *Nitrosomonas* sp. 1392.9 (20%) > *Azotobacter* sp 1242.8 (18%) > *Staphylococcus* sp 1210.7 (17.3%) > *Micrococcus* sp 434.6 (6%). This confirms or emphasizes the thought of [17] that *Pseudomonas* and *Bacillus* species are more adapted to survival and biodegradation in marine environments. Obi et al, 2016 [18] also reported *Pseudomonas* sp as the best degrader when compared with others in their study because *Pseudomonas aeruginosa* is a typical strain for rhamnolipid production and can utilize crude oil as the sole carbon source [19].

The surfactant and indicator were also used to identify the hydrocarbon degraders by monitoring the colour change at the end of the 14 days. Seven of the thirteen isolates (*Staphylococcus* sp, *Pseudomonas aeruginosa* (MW369466), *Klebsiella pneumoniae*, *Bacillus amyloliquefaciens*, *Brevibacillus brevis*, *Pseudomonas aeruginosa* (MN314747), *Nitrobacter* sp) showed a positive reaction as they turned colourless while six (*Micrococcus* sp, *Klebsiella* sp, *Azotobacter* sp, *Staphylococcus* sp, *Bacillus* sp and *Nitrosomonas* sp) showed a negative reaction as they turned pink [20-26].





Fig. 1. Amount of crude oil degraded by the isolates



**Fig. 2. Percentage (%) degradation of the isolates**

The screening using surfactant and indicator were compared against the screening with a spectrophotometer and the organisms with the highest degradation rate showed a positive result with a colour change from blue to colourless. The five (5) isolates that showed a higher degradation rate changed to colourless confirmation of the hydrocarbon-degrading ability. They were then picked for further study [27-29].

## 5. CONCLUSION

Results of this research revealed that, the mangrove plants roots and leaves contains high numbers of active endophytic bacteria [30-33]. The organisms isolated include: *Pseudomonas* sp, *Bacillus* sp, *Staphylococcus* sp, *Micrococcus* sp, *Klebsiella* sp, *Azotobacter* sp, *Nitrobacter* sp. and *Nitrosomonas* sp.

Most of the organisms identified in this study possess catabolic abilities and showed the ability to degrade crude oil. It was observed that, the mangrove plants (roots and leaves) harbour bacterial genera that may play important role in the nitrogen cycle and may also bring about bioremediation of polluted environment. Hence, a consortium of these bacteria can be used to clean up oil spills in hydrocarbon polluted environment.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Carugati L, Gatto B, Rastelli E, Martire M, Coral C, Greco S, Danovaro R. Impact of mangrove forests degradation on

2. biodiversity and ecosystem functioning. *Science Rep.* 2018;8:13298.
2. Thatoi H, Behera BC, Mishra RR, Dutta SK. Biodiversity and biotechnological potential of microorganisms from mangrove ecosystems: A review. *Ann Microbiol.* 2013;63:1-19.
3. Anyasi R, Atagana HI. Endophyte: Understanding the Microbes and its Applications. *Pakistan Journal of Biological Sciences.* 2019;22(4):154-167.
4. Compant S, Vacher C. Endophyte biotechnology: Potential for agriculture and pharmacology. 2019;32-41
5. Holguin G, Vazquez P, Bashan Y. The role of sediment microorganisms in the productivity, conservation, and rehabilitation of mangrove ecosystems: An overview. *Biology and Fertility of Soils.* 2001;33:265-278.
6. Nrior R, Ogbonna DN, Alabo AE. Biodegradation of drilling fluid used in upstream sector of the Nigeria Petroleum Industry in Marine Water Environment. *International Journal Waste Resources.* 2017 7(4):302-306.
7. Das S, Lyla PS, Khan SA. Spatial variation of aerobic culturable heterotrophic bacterial population in sediment of the Continental slope of western Bay of Bengal. *Ind J Mar Sci.* 2006;36:51-58.
8. Castro RA, Quecine MC, Lacava PT, Batista BD, Luvizotto DM, Marcon J, Ferreira A, Melo IS, Azevedo JL. Isolation and enzyme bioprospection of endophytic bacteria associated with plants of Brazilian mangrove ecosystem. *Springer Plus.* 2014;3:382. Available: <https://doi.org/10.1186/2193-1801-3-382>



9. Gupta RM, Kale PS, Rathi ML, Jadhav NN. Isolation, Characterization and Identification of Endophytic Bacteria by 16S rRNA partial Sequencing technology from Roots and Leaves of *Prosopis cineraria* Plant. Asian Journal of Plant Sciences & Research. 2015;5(6):36–43.
10. Ebilayefa G, Ariole CN. Molecular characterization of hydrocarbon degrading endophytic fungi from *Eicchhorniacrassipes*. International Journal of Maritime and Interdisciplinary Research (IJMIR). 2020;1(1):6.
11. Okpokwasili GC, Okorie BB. Biodegradation potentials of microorganisms isolated from car engine lubricating oil. Tribology International. 1988;21:215-220.
12. Nrior R, Otuogha IM. Enhanced Biodegradation of Degreaser using *Pseudomonas* and *Bacillus* Species in Fresh Water Ecosystem. Current Journal of Applied Science and Technology. 2019;35(2):1-10.
13. Tremblay J, Yergeau E, Fortin N, Cobanli S, Elias M, King TL, et al. Chemical dispersants enhance the activity of oil- and gas condensate-degrading marine bacteria. ISME J. 2017;11: 2793–2808. DOI: 10.1038/ismej.2017.129
14. Margesin R, Labbé D, Schinner F, Greer CW, Whyte LG. Characterization of hydrocarbon-degrading microbial populations in contaminated and pristine alpine soils. *Applied Environmental Microbiology* 2003;69:3085–3092. DOI: 10.1128/AEM.69.6.3085-3092.2003
15. Varjani SJ, Upasani VN. Biodegradation of petroleum hydrocarbons by oleophilic strain of *Pseudomonas aeruginosa* NCIM 5514. Bioresources Technology. 2016;222: 195–201.
16. Xu X, Zhai Z, Li H, Wang Q, Han X, Yu H. Synergetic effect of bio-photocatalytichybrid system: g-C<sub>3</sub>N<sub>4</sub>, and *Acinetobacter*, sp. JLS1 for enhanced degradation of C16 alkane. Chem. Engineering Journal. 2017;323:520–529. DOI: 10.1016/j.cej.2017.04.138
17. Odokuma LO. Bioremediation of crude oil polluted tropical mangrove Environment. Journal of Applied Science and Environmental Management. 2003;7(2):23-29.
18. Obi LU, Atagana HI, Adeleke RA. Isolation and characterization of crude oil sludge degrading bacteria. Springer Plus. 2016;5 1-13.
19. Prakash B, Irfan M. *Pseudomonas aeruginosa* is Present in Crude Oil Contaminated Sites of Barmer Region (India). J Bioremed Biodegrad. 2011;2:129.
20. Anjum N, Chandra R. Endophytic bacteria: optimization of isolation procedure from various medicinal plants and their preliminary characterization. Asian Journal of Pharmaceutical and Clinical Research. 2015;8:233–238.
21. Araujo WL, Marcon J, Maccheroni JW, van Elsas JD, van Vuurde JWL, Azevedo JL. “Diversity of endophytic bacterial populations and their interaction with *Xylella fastidiosaincitrus* plants,” Applied and Environmental Microbiology. 2002; 68(10):4906–4914.
22. Cheesbrough M. District Laboratory Practices in Tropical Countries Second Edition Cambridge: Cambridge University Press, ELBS. 2006;7:312-315.
23. Deivanai S, Bindusara AS, Prabhakar G, Bhore SJ. Culturable Bacterial Endophytes isolated from Mangrove tree (*Rhizophora apiculata* Blue) enhance seedling growth in Rice. Journal of Natural Science, Biology & Medicine. 2014;5(2):437-444.
24. Dias ACF, Andreote FD, DiniAndreote F, Lacava PT, Sá AL, Melo IS, Azevedo JL, Araújo WL, Diversity and biotechnological potential of culturable bacteria from Brazilian mangrove sediment. World J. Microbiol. Biotechnol. 2009;25:1305–1311.
25. Duke NC. Mangroves. In: Hopley, D. (Ed.), Encyclopedia of Modern Coral Reefs. Structure, Form and Process. Springer, Dordrecht, The Netherlands. 2011;655-663.
26. Holt JG, Krieg NR, Sneath PHA, Staley JT, Williams ST. *Bergey’s Manual of Determinative Bacteriology*, Williams and Wilkins, Baltimore, Maryland, USA. 1994;151–157.
27. Kadafa AA. Environmental impacts of oil exploration and exploitation in the Niger Delta of Nigeria. *Global Journal of Science, Frontier Research Environmental Earth Science*. 2012a;12(3):1–11.
28. Kathiresan K, Selvam MM. Evaluation of beneficial bacteria from mangrove soil. Bot. Mar. 2006;49:86–88.
29. Nair DN, Padmavathy S. Impact of Endophytic Microorganisms on Plants, Environment and Humans. The Scientific World Journal; 2014. Article ID:250693

30. Numbere A, Maduike E. Investigation of the anti-bacterial properties of mangrove fern, *Acrostic humaureum* in the Niger Delta, Nigeria. *African Journal of Biotechnology*. 2021;20(4):142-149.
31. Ramírez-Elías MA, Ferrera-Cerratoa R Alarcóna A, Almaraz JJ, Ramírez-Valverde G, de Bashanc LE, Esparza-García FJ, García-Barradas O. Identification of culturable microbial functional groups isolated from the rhizosphere of four species of mangroves and their biotechnological potential. *Applied Soil Ecology*. 2014;82:1–10.
32. Savitri WD, Wirjaputra MV, Hardjo PH. Isolation ad Characterization of Endophytic Bacteria from the leaf explants of *Avicennia marina* (Forsk); 2016.
33. Yaish MW, Antony I, Glick BR. Isolation and characterization of endophytic Plant growth-promoting bacteria from date palm tree (*Phoenix dactylifera* L.) and their potential role in salinity tolerance. *Antonie Van Leeuwenhoek*. 2015;107:1519–1532.

© 2022 Somiari et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/85551>