



IPM – An Ecofriendly Approach to Manage the Pink Boll Worm (*Pectinophora gossypiella*) in *Bt* Cotton of Mancherial District, Telangana State

Nagaraju Alugoju ^{a*}, Rajeshwar Malavath ^a,
Shivakrishna Kota ^a, I. Thirupathi ^a, U. Sravanthi ^a
and Sathish Kumar Bollaveni ^a

^a PJTSAU-KVK, Bellampalli, Mancherial, Telangana, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i242638

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/94884>

Original Research Article

Received: 12/10/2022

Accepted: 20/12/2022

Published: 23/12/2022

ABSTRACT

A Front Line Demonstration (FLD) on Integrated Pest Management (IPM) practices for the management of pink boll worm, *Pectinophora gossypiella* (Saunders) in *Bt* cotton has been conducted in different villages in various mandals of Mancherial district in Telangana state during *Kharif* (June — December) season of 2018–19, 2019–20 and 2020–21, respectively to create awareness about the IPM practices among the farming community. The adoption of IPM practices like installation of pheromone traps at 45 DAS, removal & destruction of rosette flowers, spraying of Azadiractin 1500ppm and need-based application of selective insecticides etc. were carried out. The results revealed that the lowest per cent rosette flowers (6.63, 6.52 and 2.84)

*Corresponding author: E-mail: nagarajualugoju.entomologist@gmail.com;

and green boll damage (5.71, 8.91 and 13.39) was recorded in IPM demonstrated plots over farmers practice (per cent rosette flowers (11.69, 8.51 & 13.39), green boll damage (8.57, 10.46 and 21.84) with increased yields of 7.83, 9.98 and 9.85 during corresponding *Kharif* (June – December), 2018–19, 2019–20 and 2020–21. Further, the demonstration plots registered with higher cotton yield of 2079, 1785 and 1817 kg ha⁻¹ as compared to 1928, 1623 and 1654 kg ha⁻¹ under farmers practice. The Cost- Benefit ratio of 1.89, 1.90 and 1.80, respectively in the technology demonstrated plots whereas in farmers practice the recorded Cost - Benefit ratio of 1.55, 1.64 and 1.45 during corresponding *Kharif* (June – December), 2018–19, 2019–20 and 2020–21. Hence, need to popularize the IPM practices through FLD among the farming community to alleviate the gap between the improved technology and farmer's practice.

Keywords: Cotton; integrated pest management; pink boll worm; percent rosette flower damage; percent boll damage; yield and economics.

1. INTRODUCTION

Cotton (*Gossypium spp.*) is an important and oldest commercial fiber crop in India as well as in the state of Telangana. Popularly known as white gold, playing a key role in economic affairs of the nation. In India the crop is cultivated in an area of 13.4 mha with a production of 36.5 mbales with a productivity of 460.0 kg ha⁻¹. The area of cotton in Telangana state is 21.2 lakh hectares whereas production is 54.0 lakh bales with a productivity of 432.0 kg ha⁻¹ [1]. In India, annually Rs. 3,39,660 million worth of yield loss [2]. In India, since its introduction, the cultivation of *Bt* cotton has increased greater than before [3] and is instigated by 166 recorded insect pest species on cotton [4]. Among them Pink bollworm, *Pectinophora gossypiella* (Saunders) is one of the key destructive insect pests of cotton [5] with an extensive range across India [6] leading to significant loss to the crop [7] by damaging squares, flowers and bolls [8-10]. In Telangana state the reduction in yield ranges from 35–90% [11]. Large-scale cultivation of *Bt* cotton can impose a continuous and intense selection pressure on bollworms leading to the development of resistance to toxins [12]. Recently, in India, the development of resistance in Pink bollworm to Cry 1 Ac and Cry 2 Ab toxins has been reported [6]. During the recent past, Pink Boll Worm is appearing early stage of the crop (45–60 DAS) on BG-II hybrids in Central and South India [13]. Unless extension initiatives to manage PBW were implemented on war footing, the situation might have further lead to yield losses and had a surging effect on textile industry and Indian economy. The adoption of Integrated Pest Management helps the farming community in reducing the usage of chemical pesticides there by reducing cost of cultivation as well as increasing yields [14]. Timely implementation of the IPM interventions by

educating the farmers helps to reduce the cost of production [15]. To overcome these lapses KVKs acts as a Knowledge and Resource Centre at district level to demonstrate the technologies [16] and the output of the research is disseminated to farmers through conduction of frontline demonstrations about the developed technologies [17]. For this IPM technologies need to be practiced in cluster approach to manage the pest [18]. Hence, the following integrated pest management module under front line demonstrations has been validated in the field conditions for effective dissemination of new technologies in the different villages of various mandals in Mancherial district during *Kharif* (June – December), 2018–19, 2019–20 and 2020–21 to reduce the incidence of pink bollworm & cost on plant protection practices and to study the yield and economic impact of the technology.

2. MATERIALS AND METHODS

The present study was carried out as frontline demonstrations (FLDs) on Integrated Pest Management (IPM) of pink boll worm in *Bt* cotton was conducted in farmers fields to demonstrate the impact of IPM technology in the different adopted villages of various mandals of Mancherial district over the three years of *Kharif* (June – December), 2018–19, 2019–20 and 2020–21 by the Krishi Vigyan Kendra, Bellampalli, Mancherial, PJTSAU. In this study, 25 farmers were selected for the demonstration of the technology in the three consecutive years as per the approved technical programme of work during the State Level Technical Programmes at University level The improved technology i.e., Integrated Pest Management Practices were imposed, consisting of deep summer ploughings, erection of pheromone traps at 45 DAS @ 4acre for monitoring of the pest or

8 per acre for mass trapping and control of the pest, need based spraying of Azadiractin 1500 ppm @ 5ml lt⁻¹ with Surf/Sandovit @ 1ml or g lt⁻¹ of water as a prophylactic measure at 40–45 DAS of the crop, if the pest reaches ETL (trap catches 8/day/trap for 3 consecutive days or 10% rosette flowers or 10% damaged green bolls) then spray with Thiodicarb 75 WP @ 1.5g lt⁻¹ or Profenophos 50 EC @ 2ml per litre and termination of the crop before 200 days or by end of the December etc. Whereas, the farmers practice includes indiscriminate spraying of different insecticides like Acephate 75SP @ 300 – 400gacre⁻¹, Ampligo 150ZC (80–100 mlacre⁻¹), Chlorantraniliprole 18.5SC (60–80mlacre⁻¹) and Synthetic pyrethroids, Lambdacyhalothrin 5EC (250 mlacre⁻¹) during the cropping period from vegetative stage to end of the crop growth period. The regular field visits were taken up for recording the data on the following observations such as per cent rosette flowers, per cent green boll damage, adult moth catches/trap/week (Monthly Average), cotton yield, yield attributes and cost benefit ratio etc. These studies also provide information about the favourable periods for pest build-up that help in the management of the pest. The weather parameters viz., Maximum and Minimum Temperatures (°C) and Rainfall (mm) were recorded on monthly basis from August to December during 2018–19, 2019–20 and 2020–21, respectively. Observations on the incidence of pink bollworm in green bolls were made during the field visits at regular intervals. For this purpose, selected 30 cotton green bolls randomly with optimal size were collected to cut open the bolls for the careful examination of pink bollworm damage.

3. RESULTS

The observations were recorded on intensity of pink boll worm (*Pectinophora gossypiella*) consequently in three years during *Kharif* (June – December), 2018–19, 2019–20 and 2020–21.

3.1 Rosette flowers and Green Boll Damage

The statistics presented in the Table 1 showed that the per cent rosette flowers found to 6.63%, 6.52% and 2.84% in the demonstration of IPM module as compared to 11.69%, 8.51% and 8.20% in farmers practice during *Kharif* (June – December), 2018–19, 2019–20 and 2020–21, respectively. Whereas the percent pink bollworm

infestation in green boll found to 5.7%¹, 8.91% and 13.39% in IPM module as compared to 8.57%, 10.46% and 21.84% in farmers practice. In all the three years, the demonstration plots showed significant differences in the percentage damage in rosette flowers and green bolls against farmers practice.

3.2 Adult Trap Catches

Studies on adult moth catches of pink bollworm in pheromone traps installed in the demonstrations indicated that the early catches of adults noticed during October, 2018–19, 2019–20 and 2020–21 with a peak adult emergence in the months of November and December of last two years.

3.3 Yield Impact

The information regarding the impact of technology demonstrated in terms of escalation in yield has been presented in Table 3. The data showed that the yield of cotton improved by 7.83% 9.98% and 9.85% in the demonstration of IPM module as compared to farmer's practice. In all the three years, the demonstration plots showed significant differences in the yields against farmers practice.

3.4 Economic Impact

The total cost of cultivation incurred, gross and net returns and B:C ratio were to assessed to study the economic impact of technology of IPM module and farmer practice. The data (Table 3) revealed that the yield of IPM module demonstrated field was 2079, 1785 and 1817 kg ha⁻¹ whereas in the farmer practice, the yield was 1928, 1623 and 1654 kgha⁻¹ during *Kharif* (June – December), 2018–19 and 2019-20 respectively. The economic analysis results revealed that the cotton crop recorded higher returns from demonstration as 103950, 92713 and 89917 Rs ha⁻¹ as compared to 96400, 85208 and 81873 Rs ha⁻¹ in farmers practice during *Kharif* (June – December), 2018–19, 2019–20 and 2020–21, respectively. The B:C Ratio in IPM module was high 1.89, 1.90 and 1.80 when compared to farmer practice 1.55, 1.64 and 1.45 during *Kharif* (June – December), 2018–19, 2019–20 and 2020–21, respectively.

Table 1. Percent of rosette flower, green boll damage, green boll locule damage, open boll damage, open boll locule damage during Kharif (June – December), 2018–19, 2019–20 and 2020–21


Particulars	Percent Rosette Flowers (%)			Percent Green Boll Damage (%)		
	2018–19	2019–20	2020–21	2018–19	2019–20	2020–21
IPM Module	6.63	6.52	2.84	5.71	8.91	13.39
Farmer Practice	11.69	8.51	8.20	8.57	10.46	21.84

Table 2. Seasonal incidence of pink boll worm, *Pectinophora gossypiella* in cotton in relation to meteorological condition during Kharif (June – December), 2018-19, 2019–20 and 2020-21

Month	2018–19								2019–20								2020–21							
	RF (mm)	Temperature (°C)		Demonstration		Farmer's Practice		RF (mm)	Temperature (°C)		Demonstration		Farmer's Practice		RF (mm)	Temperature (°C)		Demonstration		Farmer's Practice				
		Max	Min	Avg. Trap Catches	% Flower Damage	% Boll Damage	% Flower Damage		% Boll Damage	Max	Min	Avg. Trap Catches	% Flower Damage	% Boll Damage		% Flower Damage	% Boll Damage	Max	Min	Avg. Trap Catches	% Flower Damage	% Boll Damage	% Flower Damage	% Boll Damage
August	531.5	30.0	23.2	0.11	8.88	0.00	10.37	0.00	469.2	30.6	23.6	2.25	7.50	12.50	10.58	10.67	323.1	29.7	22.8	2.50	0.00	0.00	8.00	0.00
September	107.5	32.6	23.0	0.00	4.87	4.31	8.78	5.35	284.3	30.7	23.3	0.89	7.66	8.65	6.13	6.63	130.0	30.9	23.1	3.42	6.00	6.46	12.77	14.46
October	7.4	34.3	20.6	2.90	13.44	9.21	13.01	9.12	123.0	31.4	22.3	2.27	9.16	5.13	9.76	12.08	87.1	31.0	21.1	6.05	4.63	12.84	8.42	22.11
November	0.0	33.1	17.9	5.12	4.71	7.78	8.68	11.38	8.1	31.1	18.8	3.18	7.04	8.99	5.98	7.40	12.0	30.3	18.1	15.40	2.40	20.00	6.40	32.80
December	28.9	28.6	15.1	1.00	1.25	7.25	17.62	17.01	1.1	28.7	16.4	6.95	1.25	9.28	10.12	15.55	8.8	29.5	15.4	35.75	1.20	27.67	5.4	39.85
Average	135.1	31.7	20.0	1.826	6.63	5.71	11.69	8.57	177.2	30.5	20.9	3.108	6.52	8.91	8.51	10.47	112.2	30.3	20.1	12.62	2.84	13.39	8.19	21.84

Table 3. Economic analysis of on-farm trial on cotton during *Kharif*(June – December), 2018–19, 2019–20 and 2020-21

Particulars	Yield (Kgha ⁻¹)			Per cent increase in yield over check			Cost of cultivation Rs. ha ⁻¹			Gross returns (Rs. ha ⁻¹)			B:C Ratio		
	2018–19	2019–20	2020–21	2018–19	2019–20	2020–21	2018–19	2019–20	2020–21	2018–19	2019–20	2020–21	2018–19	2019–20	2020–21
IPM module	2079	1785	1817	7.83	9.98	9.85	54795	48723	50153	103950	92713	89917	1.89	1.90	1.80
Farmer practice	1928	1623	1654	-	-	-	61940	52048	56486	96400	85208	81873	1.55	1.64	1.45



Graph 1. Graphical Representation of Average Month wise Trapcatches in relation with temperature in Kharif (June – December), 2018 - 2020

4. DISCUSSION

IPM module showed positive results with respect to yield and economics of cotton. It was marked from the results that B:C Ratio of cotton crop in IPM module was higher as compared to farmer practice in both the years. Because of non-adoption of IPM module for pink bollworm management in cotton crop resulted in lower B:C Ratio in farmer practice. Thus, promising B:C Ratio and higher net returns in IPM module showed the economic sustainability of the demonstrated technology and influenced the farmers on the utility of technology provided at actual farming situation. Similar results were also reported by Shankar et al., [18], Undhad et al., [19], Surulivelu, [20] and Verma, [21].

5. CONCLUSION

In IPM module, documented higher cotton yield with gross returns 103950, 92713 and 89917 Rs ha⁻¹ which was about 7.83%, 9.98% and 9.85% higher than the non- IPM module with 96400, 85208 and 81873 Rs ha⁻¹ during *kharif* 2018–19 2019–20 and 2020-21, respectively. The IPM based practices were found effective in comparison to farmer practice. From the above study, it can be concluded that by adopting IPM based pink boll worm management strategies in *Bt* cotton can be efficiently managed instead of practicing chemical control measures. Similar results on yield improvement and reduced cost of cultivation on cotton through FLD has also been reported by Patel et al., [22] and Dhaka et al., [23].

ACKNOWLEDGEMENT

The authors are thankful to Director of ATARI Zone-XandPJTSAU Hyderabad for providing financial support and Associate Director of Research, R.A.R.S., Polasa,Jagtial, PJTSAUfor providing the necessary facilitiesin carrying out the present investigation.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Anonymous. Directorate of economics and statistics, department of agriculture, cooperation and farmers Welfare, Ministry

- of Agriculture and Farmers Welfare, Govt.of India. 2020; 1–6.
Available: <http://eands.dacnet.nic.in>
Accessed on 23th November, 2021.
2. Dhaliwal GS, Vikas J, Dhawan AK. Insect pest problems and crop losses: changing trends. Indian Journal of Entomology. 2010;37(1):1–7.
3. Henneberry TJ, Naranjo SE. Integrated management approaches for pink bollworm in the Southwestern United States. Integrated Pest Management Review. 1998;3:31-52.
4. Puri SN, Murthy KS, Sharma OP. Integrated pest management for sustainable cotton production. In: Sundaram, V. (Ed.), Handbook of cotton. Indian Society of Cotton Improvement, CIRCOT, Mumbai. 1999;233–245.
5. Balakrishnan N, Vinodkumar B, Sivasubramaian. Bioefficacy of Kina dongold against sucking pests of cotton. Madras Agricultural Journal. 2010; 97:88-91.
6. Naik VC, Kumbhare S, Kranthi S, Satija U, Kranthi KR. Field-evolved resistance of pink bollworm, *Pectinophora gossypiella* (Saunders) (Lepidoptera: Gelechiidae), to transgenic *Bacillus thuringiensis* (Bt) cotton expressing crystal 1Ac (Cry1Ac) and Cry2Ab in India. Pest Management Sciences. 2018;11:2544-54.
7. Deore JS, Borikar PS, Yadav GA, Dhumal MS. Efficacy of newer insecticides against bollworm complex in cotton. Pestology. 2010;34(2):12–17.
8. Ghosh SK. G.M.Crops: Rationally irresistible. Current Science. 2001;81(6) :655–660.
9. Amin AA, Gergis MF. Integrated pest management approaches strategies for controlling certain cotton key pests in middle Egypt. Agronomy Research. 2021;4:121–128.
10. Roopsingh M, Kumar P, Bharat LM, Bhagwan Singh M. Management of pinkbollworm, *Pectinophora gossypiella* (Saunders) using mating disruption pheromone (PB Rope L) in cotton. Acta Scientific Agriculture. 2021;5(3):24–31.
11. Naik VCB, Subbireddy KB, Sandhya K, Nagrare VS, Sujit K, Nandini GN, Waghmare VN. Pink bollworm, *Pectinophora gossypiella* (Saunders) (Lepidoptera: Gelechiidae) survival on transgenic cotton in India. Egyptian Journal of Biological Pest Control 2021;31(40):1–7.

12. Hardee DD, Van Duyn JW, Layton MB, Bagwell RD. Bt cotton continued effectiveness by managing resistance of tobacco budworm and bollworm. USDA-ARS- 154. United States Department of Agriculture -Agricultural Research Service, Stoneville, MS; 2001.
13. Sivarama Krishna M, Rama Reddy Y, Chandrayudu E. Validation of pink bollworm *Pectinophora gossypiella* (Saunders) management strategies in Bt cotton. Journal of Entomology and Zoology Studies. 2020;8(5):2064-2067.
14. Ajanta B, Tanwar RK, Kumar A, Singh SP, Kumar R, Kanwar V. Evaluation of pest management practices against sucking pests of Bt cotton. Indian Journal of Agricultural Sciences. 2019;89(1):124-129.
15. Raghava NV, Punnarao P. Impact of frontline demonstrations on groundnut production technology in Guntur District of Andhra Pradesh. Agriculture Update. 2013; 8(1-2): 283-290.
16. Sharma RK, Bhati DS, Sharma SK. Impact of frontline demonstrations on rapeseed mustard growers. Journal of Progressive Agriculture. 2017;8(1):115-118.
17. Singh DK, Gautam US, Singh RK. Study on yield gap and level of demonstrated crop production technology in sagar district. Indian Research Journal of Extension Education. 2007; 7(2&3): 94-95.
18. Shankar M, Aariff Khan MA, BalazziiNaaiik RVT, Sumalini K. Extension interventions for enlightening tribal farmers for enhancing cotton production in Nalgonda district, Telangana. International Journal of Bio-resource and Stress Management. 2022;13(4):365-371.
19. Undhad SV, Sharma PS, Prajapati VS. Impact of frontline demonstrations on integrated approaches against the management of pink bollworm in Bt-Cotton. Gujarat Journal of Extension Education. 2018;29(2):184-186.7.
20. Surulivelu T. Sustainable IPM for irrigated cotton. Model training course on "Cultivation of long staple cotton (ELS)" December 15-22. Central institute for cotton research, Regional station, Coimbatore. 2007;84-89.
21. Verma SK. Population dynamics of pink bollworm *Pectinophora gossypiella* (saunders) in cotton crop. International Journal of Pure & Applied Bioscience. 2017;5(2):801-806.
22. Patel MM, Jhaharia AK, Khadda BS, Patil LM. Frontline Demonstration: An Effective Communication Approach for Dissemination of Sustainable cotton production technology. Indian Journal Extension Education. 2013;21:63-67.
23. Dhaka BL, Meena BS, Suwalka RL. Popularization of improved maize production technology through frontline demonstration in south eastern Rajasthan. Journal of Agricultural Science. 2010;1(1): 39-42.

© 2022 Alugoju et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/94884>