



Impact of Cluster Front Line Demonstration on Productivity and Yield Gap of Lentil in Kokrajhar District of Assam

Bhupen Kumar Baishya^{a++}, Priyanka Bora^{a++*} and Manoj Kumar Bhuyan^{b#}

^a *Krishi Vigyan Kendra, Kokrajhar, India.*

^b *Krishi Vigyan Kendra, Nalbari, India.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJAEES/2022/v40i121793

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/94622>

Original Research Article

Received: 12/10/2022

Accepted: 20/12/2022

Published: 26/12/2022

ABSTRACT

The Cluster Frontline Demonstration (CFLD) on lentil var.WBL-77 was conducted by Krishi Vigyan Kendra (KVK), Kokrajhar to find out the grain yield, technological gap, extension gap, and economics during 2020-21 and 2021-22. In both year, lentil variety WBL-77 was used in the demonstration plot and considering local variety as check in the farmer's plot. The result of the study revealed that the yield of lentil was found to be 10.0 q/ha and 9.50 q/ha, respectively under the demonstration plot as compared to 7.0 q/ha and 6.50 q/ha, respectively under the farmer's plot and the yield increased by 42.85 and 46.15 percent, respectively during the year, 2020-21 & 2021-

⁺⁺Subject Matter Specialist;

[#]Senior Scientist & Head;

^{*}Corresponding author: E-mail: borapriyanka91@gmail.com;

22. The technology gap, extension gap and technology index were found to be 2.0 q/ha & 2.5 q/ha, 3.0 q/ha & 3.0 q/ha, 16.6 % & 20.83%, respectively during 2020-21 & 2021-22. The net return (Rs. 31314.00 /ha & Rs.34074.00 /ha, respectively during 2020-21 & 2021-22) and Benefit Cost ratio (2.59 & 2.87, respectively during 2020-21 & 2021-22) were found to be highest under demonstration plot in both the years. The results depict that the seed yield of lentil in the Kokrajhar district could be enhanced by using lentil var. WBL-77 over local variety and CFLD could be an effective tool to minimize yield gaps.

Keywords: Lentil; cluster frontline demonstration; yield gap; B-C ratio.

1. INTRODUCTION

Pulses are good sources of proteins, vitamins & minerals, a second important constituent of the Indian diet therefore; it plays an important role in the nutritional security of India. Besides, being legumes they can fix atmospheric nitrogen, maintains soil sustainability, nutritious animal fodder, act as mulch and check soil erosion. In Assam, pulses are grown in an area of 146.4 thousand hectares with a production of 107.5 thousand tons and productivity of 735.0 kg/ha [1]. Lentil is an important pulse crop. In Assam, production of most of the pulse crops is less than the requirement except for blackgram and pea & also the present production can meet only 20% of the present requirement therefore, it is very important to grow *rabi* pulse like lentil after harvesting of paddy to reduce the gap between demand and supply. There could be various reasons for the low productivity of lentil in the state like the use of local varieties, imbalanced use of fertilizer, delayed sowing, no proper crop management, etc. To combat the issue of low productivity of lentil in Kokrajhar district, different trials were conducted on varietal performance, integrated nutrient management, water management and weed management etc. But, CFLD could be a better way to facilitate the bridge between extension specialists and farmers to overcome the yield gaps. Because, it is conducted under the supervision of KVK scientists to demonstrate the production potentiality of location-specific newly released technologies with proper crop management practices on large area (cluster) in different locations and facilitates speedy dissemination of technology through extension activities like group discussion, field days etc. Hence, CFLD on lentil variety WBL-77 was conducted in Kokrajhar district to increase the productivity of lentil and reduce the extension and technology gap.

2. MATERIALS AND METHODS

The Cluster Frontline Demonstration on lentil (var. WBL-77) was conducted by Krishi Vigyan Kendra, Kokrajhar during *rabi* seasons of two consecutive years, *i.e.* 2020-21 and 2021-22. In this programme 22 demonstrations in 10 ha and 23 demonstrations in 10 ha were carried out during 2020-21 and 2021-22, respectively where, the number of demonstrations was equal to the number of farmers. Firstly, group meeting was held and discussed regarding low yield of lentil in Kokrajhar district. Then farmers were aware to importance of improved technology by KVK scientist which could increase the yield of the crop. After that selected farmers were trained regarding use of improved lentil var. WBL 77, timely sowing, application of balanced dose of fertilizer, water management, weed management etc which they will have to perform in demonstration plot. In CFLD, inputs are provided by the KVK and the adjoining farmer's field is treated as a check or farmer's practice. The brief about the technologies adopted in the farmer's field is given in Table 1. After harvesting the crop the data were collected from both the demonstration and farmer's plot, analyzed and the cost of cultivation, net income, and benefit-cost ratio were calculated. The extension gap, technology gap, and technology index were calculated by using the formulas given by Kadian et al. [2] and Samui et al. [3].

1. Technology gap = Potential yield – Average demo yield.
2. Extension gap = Average demo yield – Average farmer's practice yield.
3. Technology index = $\{(\text{Potential yield} - \text{Average demo yield}) \times 100\} / \text{Potential yield}$.

Table 1. Package of practices followed in demonstration and farmers plot under CFLD on lentil

Particulars	Demonstration plot	Farmer's plot
Variety	WBL-77	Local
Seed rate	30 kg/ha	38-40 kg/ha
Seed treatment	Seed treatment with <i>Rhizobium</i> culture @ 50 g/kg of seeds & PSB @ 50 g/kg of seeds	No seed treatment
Fertilizer dose	Balanced fertilizer dose N:P2O5:K2O @ 10:35:15 Kg/ha	Imbalance use of fertilizer
Method of sowing	Line sowing	Broadcasting
Spacing	25 cm x 5-7 cm	-
Plant protection	Need based application	Nil

3. RESULTS AND DISCUSSION

3.1 Yield

The data (Table 2) depicted that the average seed yield of lentil was found to be 10.0 q/ha and 9.5 q/ha during 2020-21 & 2021-22, respectively under demonstrated plot whereas, under the farmer's plot the seed yield was 7.0 q/ha and 6.5 q/ha during respective years. This resulted in 42.85 and 46.15 percentage increases in seed yield during 2020-21 & 2021-22, respectively as compared to the farmer's plot. The increase in yield could be due to integrated crop management approaches like optimum seed rate and spacing which maintains optimum plant population and reduces the competition for space, sunlight, nutrition and moisture. Secondly, seed treatment with *Rhizobium*/ PSB increases root nodulation to fix atmospheric nitrogen & PSB improves the phosphorus uptake from soil. Thirdly, need-based plant protection protects against disease and pests and finally increases the yield attributing characters. This indicated that the productivity of lentil in the Kokrajhar district could be enhanced by creating awareness among farmers and adopting the improved technologies. The findings conform with the findings of Singha et al. [4], Reja et al. [5] and Deka et al. [6].

3.2 Technology Gap, Extension Gap and Technology Index

The technology gap (Table 2) was found to be 2.0 q/ha & 2.5 q/ha during 2020-21 and 2021-22,

respectively. This could be due to variation in soil fertility, weather conditions etc. The data depicted that the extension gap of 3.0 q/ha was observed in both years. This emphasizes the need to get farmers to be aware of the improved agricultural techniques to enhance production and mitigate the extension gap. The technology index was found to be 16.60 and 20.83 percent during the years, respectively. The technology index indicates the viability or feasibility of the improved technology in the farmer's field. The lower the difference between potential and demonstration yield (technology index) means the technology is more feasible or profitable for the farmers. The results are similar to the findings of Singha et al. [4], Singh et al. [7] and Basumatary et al. [8].

3.3 Economics

The data of economic analysis of lentil production revealed that the net return (Rs. 31314.0/ha in 2020-21 and Rs. 34074.0/ha in 2021-22) and benefit-cost ratio (2.59 in 2020-21 and 2.87 in 2021-22) were found higher under demonstration plot as compared to farmers plot. This could be due to higher yield obtained under demonstration plot as compared to farmer's practice. In this study, a higher benefit-cost ratio under the demonstration plot indicates that the farm income can be enhanced by cultivating lentil variety WBL-77 with improved technologies; which uplift the livelihood of the farming community. The results were conformed with the findings of Mandal et al. [9] and Bezbaruah et al. [10].

Table 2. Productivity, technology gap, extension gap and technology index of CFLD on Lentil

Year	Area (ha)	No. of farmers	Seed yield (q/ha)			% increase over control/farmers practice	Technology gap (q/ha)	Extension gap (q/ha)	Technology index (%)
			Potential	Demonstration	Control				
2020-21	10	22	12	10.00	7.00	42.85	2.0	3.0	16.60
2021-22	10	23	12	9.50	6.50	46.15	2.5	3.0	20.83

Table 3. Total cost and return of lentil production

Year	Gross cost (Rs./ha)		Gross return (Rs./ha)		Net return (Rs./ha)		B:C	
	Demonstration	Farmer's Practice	Demonstration	Farmer's Practice	Demonstration	Farmer's Practice	Demonstration	Farmer's Practice
2020-21	19686	19466	51000	35700	31314	16234	2.59	1.83
2021-22	18176	16000	52250	35750	34074	19750	2.87	2.23

4. CONCLUSION AND RECOMMENDATION

The CFLD conducted on lentil revealed that the lentil variety WBL-77 gave a higher yield and net return under the recommended package of practices in the demonstration plot as compared to the local variety in the farmer's plot (check) which implies that the productivity of lentil could be enhanced by adopting improved technologies and proper crop management practices. Moreover, scientists or extension specialists should play a vital role in getting farmers to be aware of the innovations by showing the technologies at the forefront under the umbrella of CFLD. Hence, CFLD could be an effective way to mitigate the yield gap and makes the country self-sufficient.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Barman N, Hazarika JP, Talukdar U, Deka N. Supply-demand projection and gap analysis for major pulses in Assam. *Economic Affairs*. 2020;65(4):619-625.
2. Kadian KS, Sharma R, Sharma AK. Evaluation of frontline demonstration trials on oilseeds in Kangra valley of Himachal Pradesh. *Annals Agri Res*. 1997;18(1):40-43.
3. Samui SK, Maitra S, Roy DK, Mandal AK, Saha D. Evaluation on front line demonstration on groundnut. *Journal of the Indian Society of Coastal Agricultural Research*. 2000;18(2):180-183.
4. Singha AK, Deka BC, Parisa D, Nongrum C, Singha A. Yield gap and economic analysis of cluster frontline demonstrations (CFLDs) on pulses in Eastern Himalayan Region of India. *Journal of Pharmacognosy and Phytochemistry*. 2020;9(3):606-610.
5. Reja H, Mandi S, Kundu M, Nath R, Goswami SB. Performance of different lentil varieties in new alluvial zone of West Bengal. *The Bioscan*. 2017;12(3):1673-1676.
6. Deka P, Baishya BK, Bhagwati G, Bhuyan MK, Nath RK. Yield gap analysis of rapeseed (*Brassica campestris* var. Toria) in Kokrajhar district of Assam, India. *Asian Journal of Agricultural Extension, Economics & Sociology*. 2021;39(10):425-428.
7. Singh M, Govind H, Deokaran M. Yield gap minimization in Lentil (*Cicer arietinum* L.) under Frontline Demonstration conducted in Indo-Gangetic Plains of Eastern India. *Journal of Krishi Vigyan*. 2022;10(2):336-341.
8. Basumatary M, Deka RS, Bezbaruah R, Hoque J. Productivity and profitability of toria through Frontline Demonstrations (CFLD) in Morigaon district. *Asian Journal of Agricultural Extension, Economics and Sociology*. 2022;40(9):202-208.
9. Mandal S, Ray P, Mandal S, Ankure P. Frontline Demonstration on lentil using improved varieties for increasing productivity under lateritic soil of West Bengal. *Legume Research*. 2019;42(3):426-429.
10. Bezbaruah R, Deka RS. Impact of Cluster Frontline Demonstration on productivity and profitability of greengram in Morigaon district of Assam. *Journal of Krishi Vigyan*. 2020;9(1):164-169.

© 2022 Baishya et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/94622>