

Neonatal Tetanus are We Winning the War?

Lucy E. Yaguo Ide^{1*}, Tochi A. Uchenwa-Onyenegecha¹ and Alice R. Nte¹

¹Department of Paediatrics and Child Health, University of Port Harcourt Teaching Hospital, Port Harcourt, P.M.B 6173, Rivers State, Nigeria.

Authors' contributions

This work was carried out in collaboration between all authors. Authors LEYI and ARN did the study design and wrote the protocol. Authors LEYI and TAUI did the statistical analysis and literature search while analysis of the study was done by author LEYI. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJTDH/2016/19046

Editor(s):

(1) Shankar Srinivasan, Department of Health Informatics, University of Medicine & Dentistry of New Jersey, USA.

Reviewers:

(1) Anonymous, Mayo Clinic College of Medicine, USA.

(2) Gregory Edie Halle-Ekane, University of Buea, Cameroon.

Complete Peer review History: <http://sciencedomain.org/review-history/11443>

Original Research Article

Received 23rd May 2015
Accepted 21st July 2015
Published 17th September 2015

ABSTRACT

Aims: To determine if we are winning the war against deaths from Neonatal tetanus.

Study Design: Retrospective descriptive study

Place and Duration of Study: Department of Paediatrics, University of Port Harcourt Teaching Hospital (UPTH) between 1995 to 2014.

Methodology: Hospital records on Neonatal tetanus cases treated at the University of Port Harcourt Teaching Hospital from 1st January 1995 to 31st December 2014 were reviewed. The records were retrieved and reviewed for age, sex, presenting complaint, mother's immunization status, mother's educational status, examination findings, duration of hospitalization and outcome of illness.

Results: Three hundred and thirteen neonates comprising 182(58%) males and 129(41%) females were treated, 2(0.006%) did not have any record of their sex. Most of the mothers were unimmunized and delivered outside health facilities. One hundred and sixty two (51.7%) of the newborns with neonatal tetanus in this study died.

Conclusion: There appears to be an upward trend in mortality of Neonatal tetanus cases as the previous review of neonatal tetanus in UPTH between 1995 to 2009 gave a mortality of 50% while

*Corresponding author: Email: lucyaguolucy@gmail.com

the data for cases managed from 2009 to 2014 gave a higher mortality rate of 68.75% and as such we are not winning the war, the battle rages on.

Keywords: Neonatal tetanus mortality winning the war.

1. INTRODUCTION

Tetanus is a devastating disease in the neonatal period. Infection mostly results from unhygienic birth practices that expose the umbilical cord to the tetanus organism: *Clostridium tetani* [1].

The neonatal period represent the most vulnerable time for a child's survival. Global neonatal mortality rate in 2013 was estimated to be 20 per 1000 live births and that 2.8 million babies die before 28 days [2]. Neonatal tetanus (NNT) though a preventable disease has remained a major cause of morbidity and mortality among neonates worldwide. Global evidence has revealed tetanus as the highest mortality contributor among children after measles in vaccine preventable diseases [3]. Neonatal tetanus kills a baby every 9minutes [4]. WHO estimated that 49,000 newborns died from NNT in 2013 [5]. Low income countries such as Nigeria and India account for most of the deaths from NNT [6]. Over all, in Nigeria tetanus is responsible for 4% of the neonatal deaths [7]. The percentage of neonatal deaths caused by tetanus varies considerably in various states in Nigeria with the highest of 19% reported in Yobe state and the lowest of 1% in Ekiti, Lagos and Osun States [8]. In Rivers state, neonatal deaths caused by tetanus were 3% [8].

The prevalence of neonatal tetanus has remained high in many low income countries because a high proportion of deliveries still take place at home or in places where hygienic condition may be poor. In Nigeria, the percentage of deliveries at home has remained very high, only 36% of births in Nigeria are delivered in a health facility (as compared with 35% in 2008), 63% are delivered at home [8]. Northwest had the highest proportion of deliveries at home with 88% followed by North east 79% and the lowest was South east 20%. In Rivers State 49.6% delivered at home and only 49% in a health facility [8]. Overall, only 38% of deliveries were assisted by a skilled provider, 22% by a traditional birth attendant, 23% by relatives or other persons and 13% unassisted [8].

Elimination of NT was redefined by 1995 as less than one case per 1000 live births in every

district in every country. The target date was however postponed to 2000 because of slow implementation of the recommended maternal tetanus elimination strategies, in 1999, it was reviewed by UNICEF, WHO and UNFPA and the initiative was reconstituted with a 2005 target date for maternal tetanus elimination which was later shifted to [5]. Nigeria today is still one of the high risk countries that have not yet achieved the maternal and neonatal tetanus elimination goal yet [5]. Every case of neonatal tetanus in Nigeria indicates a major failure of public health practice because it can be completely prevented by immunizing females before or during pregnancy, ensuring clean delivery, clean severance of the umbilical cord and proper care of the cord after delivery. Immunisation of pregnant women or women of child bearing age (15-49 years) with at least 2 doses of tetanus toxoid is estimated to reduce mortality from NNT by 94% [6]. The percentage of women who received 2 or more tetanus toxoid (TT) injection during their last pregnancy was 48% and 53% of women had their last birth protected against neonatal tetanus [8]. The proportion of these women who received 2 or more TT injection during the pregnancy for their last live birth varies considerably across zones ranging from a high of 82% in Southeast to a low of 27% in the Northwest. In Rivers state, 76.5% of women received 2 or more TT injections during their last pregnancy and the percentage of women whose last birth was protected against neonatal tetanus was 80.3% [8].

Unfortunately, the unnecessary deaths caused by this highly preventable disease has continued in most developing countries including Nigeria despite the availability of effective preventive measures. In 1998 in Uganda, 3433 tetanus cases were recorded in newborns, of these 2043 died but after a major public health effort Uganda in 2011 was certified as having eliminated tetanus [9]. In Nigeria it is possible to eliminate tetanus with sustained public health effort.

This study therefore aims to review the trend and outcome of neonatal tetanus in University of Port Harcourt from 1st January, 1995 to 31st December, 2014 with a view of highlighting the morbidity and mortality status over the study

period and continued existence of neonatal tetanus and as such highlight possible ways to achieving maternal tetanus elimination.

2. MATERIALS AND METHODS

The cases of neonatal tetanus managed in University of Port Harcourt Teaching Hospital from 1st January 1995 to 31st December 2014 were reviewed. Ethical approval for the study was obtained from the Hospital ethics Committee. All cases were diagnosed by Paediatricians using the clinical features of the disease.

The case notes of these patients were retrieved from the records department, details obtained were socio demographic data of patients, presenting complaints, history of the illness, onset interval, incubation period, mother's educational and immunization status, examination findings, management, duration of hospitalization and outcome of illness. The Paediatric department of the hospital has a unit for management of tetanus cases (community and social paediatrics unit). The unit has three consultant paediatricians, two resident doctors, and one or two house officers. There are no nurses attached solely to this unit but we have two or more nurses on each shift covering the children medical ward with a capacity for admitting 36 patients, (nurse patient ratio 1:18) There is a two bedded tetanus room (a side room at the end of the children's medical ward) with a suction machine an oxygen concentrator, an incubator and other consumables for the management of tetanus patients. We do not have a paediatric intensive care unit for the management of these patients. Although the hospital has an intensive care unit, tetanus patients are not managed in the intensive care unit of the hospital as they cannot afford to pay for it (fifty thousand naira per night equivalent of 250 dollars per night) All cases were admitted into the tetanus room of the children's ward They all received anti tetanus serum (ATS), IV antibiotics and spasms were controlled with a combination of phenobarbitone, chlorpromazine and diazepam.

Data was analysed using SPSS 20.0 and presented using simple frequencies and in tables.

3. LIMITATIONS

There were repeated strikes by the health workers in 2014 which may have been the reason why only one case of neonatal tetanus was recorded in 2014 as most cases would have presented to private health facilities to seek medical care. Poor record keeping system in our hospital resulting in a lot of missing data and the values used in this study are actually an underestimate of the true picture.

4. RESULTS

A total of three hundred and thirteen cases of neonatal tetanus were managed in UPTH over the study period. All cases of neonatal tetanus were studied. There were 182 (58.2%) males and 129(41.2%) females (M:F 1.4:1) and 2(0.006%) did not have any record of their sex. Table 1 shows the gender characteristics of the cases and outcome. Table 2 shows the educational status of the mothers of the neonates studied. The educational status of the mothers was not indicated in most of the records (213:68%), 39(12.4%) had primary education, 60(19.1%) had secondary education and 1 (0.3%) had tertiary education.

The age distribution of cases ranged from 2-28 days with mean age of 7.82±3.26. Data on the outcome of 22 children was missing (7.0%), 90(28.8%) of children survived, 39(12.5%) were either discharged against medical advice or absconded from the hospital while 162 (51.7%) died. Figs. 1 and 2 shows the yearly distribution of cases and case fatality rates per year respectively.

The annual case fatality rates showed no consistent pattern, the percentage of deaths ranged from 20% in 2003 to 100% in 2013 and 2014. The duration of hospitalization ranged from <1 day to 35 days.

Table 1. Gender characteristics of cases and outcome

Characteristics	Cases studied (%)		Outcome	
	Total percent	Survival (%)	Parental discharge (%)	Death (%)
Sexes				
Male	182(58.2)	54(29.6)	23(12.6)	93(51.1)
Female	129(41.2)	35(27.1)	16(13.2)	68(52.7)
Sex not indicated	2(0.006)	1(50)		




Fig. 1. Yearly distribution of cases

The earliest recovery time was 7 days during which period only 5(1.6%) children recovered, most deaths occurred within the first 7days. Forty –six (14.6%) of mothers received ANC in a health facility. The place of delivery was not indicated in most of the records (166:53%), most deliveries were outside a health facility, only 29(9.2%) delivered in a health facility, 69(22%) in a TBA’s home or at home, 47(15.0%) in a prayer house. Data on cord care for 204(65.1%) patients was missing, cord care with methylated spirit was done by 54(17.2%), 59(18.8%) used Robb (pain relief ointment), 17(5.4%) Vaseline, 8(2.6%) herbs. The immune-zation status of the mothers was largely unknown with 20(6.4%) claiming to have received 1 dose of TT and 28(8.9%) received two or more doses of TT. Among these patients, 40.0% and 60.7% respectively died thus receiving at least a dose of TT increases the chances of survival.

Table 2. Educational status of mothers

Educational status	Total cases	Percentage (%)
Not indicated	213	68
Primary education	39	12.4
Secondary education	60	19.1
Tertiary education	1	0.3

5. DISCUSSION

There were more males than females in our study with male: female ratio of 1.4:1, this is similar with the report from other studies in Nigeria [10,11]. The reason for this finding is not clear but may partly explained by the preference given to the male child in the society that makes parents to seek for medical attention promptly. This contrasts with the study by Babatunde et al [12] and Peterside et al. [13] who reported more females than males with neonatal tetanus in Ado-Ekiti and Bayelsa respectively. Emodi et al. [14] in Enugu however reported equal sex ratio among patients with neonatal tetanus. The findings in our study have shown that neonatal tetanus has remained a major cause of morbidity and mortality in Nigeria despite over two decades of commitments by the public health community globally to the elimination of tetanus. In this review, a high proportion of women delivered outside a health facility, only 5.1% delivered in a hospital and 4.1% in a maternity this is similar to the report by other authors in Nigeria [11,15].

Most of the mothers were unimmunized or incompletely immunised and this may be attributed to the fact that very few mothers (14.6%) received antenatal care in a health

facility which corroborates with other studies in Nigeria [12,15]. Most mothers in our study had low level of education, 19.1% completed secondary school and 0.3% post-secondary education which has been a consistent finding by most authors in Nigeria [1,16-18]. Osaghae et al. [1] in his study illustrated that there was a vicious cycle of low level of maternal education contributing to lack of antenatal care, low immunization, delivery outside a health facility, poor cord care and other risk factors for neonatal tetanus. These prevalent conditions have persisted for several years and are the reasons for continuous existence of NNT in Nigeria thus indicating that not much impact has been made in terms of combating this preventable disease and therefore a pointer to the failure of our health care system. In developed countries with better female education, adequate vaccination coverage and neonatal care, the numbers of cases of neonatal tetanus reported were rather small when compared to developing countries [6,19].

The case fatality rates noted in this review showed that there appears to be an upward trend in the mortality rates of neonatal tetanus in the recent years though the number of cases in the recent years reduced; from the previous review between 1995-2009, the mortality rate was 50% while in the recent years between 2009, the

mortality rate increased to 68.75% and as such we are not winning the war the battle rages on. We may attribute this increase to infrastructural factors and ignorance. These include absence of a quite dark room for managing these patients during the period between 2009 and end of 2013 and also to late presentation and ignorance about immunization on the side of the mothers. These factors have been addressed as we now have a quite dark room for managing patients with tetanus, also we have a periodic health talk on tetanus where we talk about its aetiology, pathophysiology, clinical presentation, and how to prevent it on the local television that reaches every local government area of Rivers state where we are practicing. These health talks are anchored by consultants in the community paediatrics unit that care for tetanus patients. We believe this has impacted on the community as we are yet to record any case of neonatal tetanus for 2015. The mortality rate in this review falls within the range of 30-70% mortality found by other workers in Nigeria [12-15]. This calls for a reappraisal of our control programme on tetanus in Nigeria. The majority of the mortalities in our review occurred within the first seven days of admission, this is consistent with the findings in other studies in Nigeria [13,15] suggesting that the first week of hospitalization is a critical period.




Fig. 2. Yearly case fatality rates

6. CONCLUSION AND RECOMMENDATION

There is an upward trend in the mortality of tetanus suggesting a failure in the health system in combating neonatal tetanus. The reason for this upward trend in the mortality may have been because the tetanus cases managed from 2009 to 2011 were nursed in an open ward which was noisy and as such worsening spasms and therefore mortality rate. There is also an epileptic supply of vaccines in Nigeria. The government should ensure regular supply of tetanus vaccines and Immunisation campaigns for women of child bearing age (15-49 years) should be strengthened to improve the immunisation coverage for tetanus in the country. We can borrow a leaf from Uganda public health effort in achieving tetanus elimination in Nigeria.

There is need to promote female education and increase the level of public awareness on tetanus and the associated risk factors. The health facilities should be made more accessible and affordable pregnant mothers and neonates by offering them free or subsidized health care services.

CONSENT

It is not applicable.

DISCLAIMER

This manuscript was presented in the conference ""NATIONAL NEWBORN CONFERENCE"" available link is <http://newbornhealthnigeria.com/wp-content/uploads/2014/10/LIST-OF-ABSTRACTS-REAL2.docx>

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Osaghae DO, Amuabunos EA. Social factors in neonatal tetanus: focus on maternal educational status. *J Med Med Sci.* 2012;3:208-211.
2. Neonatal Mortality-UNICEF Statistics; 2015. Available: <http://www.data.unicef.org/child-mortality/neonatal> 05/03/2015
3. World Health Organization: Vaccine-Preventable diseases: Monitoring System 2010 Global Summary. Immunization, Vaccines and Biologicals. Available:http://whqlibdoc.who.int/hq/2010/WHO_IVB_2010_eng.pdf 05/03/2015
4. The Eliminate Project: Kiwanis eliminating maternal and neonatal tetanus. Available:www.kiwanis.org/kiwanis/programs/eliminating-maternal-neonatal-tetanus#.VTv-nMqYzIU (05/03/2015).
5. WHO/Maternal and Neonatal Tetanus (MNT) Elimination; 2015. Available:www.who.int/immunization/diseases/MNTE-initiative/en/ (05/03/2015).
6. Blencowe H, Lawn J, Vandelaer J, Roper M, Cousens S. Tetanus toxoid immunization to reduce mortality from neonatal tetanus. *Int J Epidemiol.* 2010; 39:102-109.
7. Nigeria State Data Profiles, Federal Ministry of Health; 2013.
8. National Population Commission (NPC) [Nigeria] and ICF International. 2014. Nigeria Demographic and Health Survey 2013. Abuja, Nigeria, and Rockville, Maryland, USA: NPC and ICF International.
9. Tetanus Wikipedia: The free Encyclopedia. Uganda announces elimination of Maternal and Neonatal Tetanus. Available:<http://www.en.wikipedia.org/wiki/Tetanus> (05/03/2015).
10. Grange AO. Neonatal tetanus in Lagos metropolis. *Niger J Paediatr.* 1991;18:12-22.
11. Onalo R, Ishiaku HM, Ogala WN. Prevalence and outcome of neonatal tetanus in Zaria, North western Nigeria. *J Infect Dev Ctries.* 2011;5:255-259.
12. Babatunde OA, Adebara VO, Atoyebi OA, Oluwagbemi OA. Neonatal tetanus in Federal Medical Center, Ido-Ekiti, South west Nigeria in a five year retrospective study. *GJMS.* 2014;4:42-47.
13. Peterside O, Duru C, George B. Neonatal tetanus at the niger delta university teaching hospital: A 5 year retrospective study. *Internet J Pediatr Neonatol* 2012; 14:(2).
14. Emodi IJ, Ikefuna AN, Obichukwu C. Incidence and outcome of neonatal tetanus in Enugu over a 10year period. *SAJCH.* 2011;5:117-119.

15. Alhaji MA, Bello MA, Elechi HA, Akuhwa RT, Bukar FL, Ibrahim HA. A review of neonatal tetanus in University of Maiduguri Teaching Hospital North eastern Nigeria. Niger Med J. 2013;54:398-401.
16. Oyedeji OA, Fadero F, Joel-Madewase V, Elemile P, Oyedeji GA. Trends in neonatal and post neonatal tetanus admissions at a Nigerian Teaching Hospital. J Infect Dev Ctries. 2012;6:847-53.
17. Orumabo RS, Mbagbaw LT. Neonatal Tetanus in Port Harcourt. Niger J Paediatr. 1986;13:115-20.
18. Osinusi K, Dawodu AH, Sodeinde O, Adeyokunu AA. Neonatal Tetanus in Ibadan. Niger J Paediatr. 1986;13:121-5.
19. CDC- Surveillance of tetanus–Chapter 16-Vaccine; 2014. Available:www.cdc.gov/vaccines/pubs/surv-manual/chpt16-tetanus.html (05/03/2015).

© 2016 Ide et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://sciencedomain.org/review-history/11443>