



Dry season Irrigation Regime Effects on Water Use, Rootzone Moisture and Yield of Cacao (*Theobroma cacao* L.) in a Rainforest Zone of Nigeria

Agele, Samuel Ohikhena ^{a*}

^a Plant Physiology & Ecology Group, Department of Crop, Soil & Pest Management, Federal University of Technology, Akure, Nigeria.

Author's contribution

The sole author designed, analysed, interpreted and prepared the manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i183293

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/96942>

Original Research Article

Received: 05/01/2023

Accepted: 07/03/2023

Published: 17/07/2023

ABSTRACT

The humid tropics is characterized by wet-dry seasonal transitions, irrigation has the potential for building adaptation and resilience to climate stress for enhancing crop performance. A field trial was conducted to investigate the effects of dry season irrigation regimes on water use, root zone moisture dynamics and yield of cacao in a rainforest zone of Nigeria. Irrigation amounts computed as EPan x Pan coefficients were based on cumulative class A Pan evaporation. Irrigation treatments were coded as IrT1 (EPan x 1,0), IrT2 (EPan x 0.7) and IrT3 (EPan x 0.5). Irrigation water was applied 5-days interval and discharged via point source emitters (2.8 l/h discharge rate) on drip lines laterally installed per row of trees. Mean irrigation requirements were 4.49 mm (9.81:9.6 l/tree/day), 3.14 mm (7.06:6.8 l/tree/day) and 2.44 mm (5.49: 4.8 l/tree/day) while total seasonal water applied were 121.19, 84.83 and 60.59 l/tree for IrT1, IrT2 and IrT3 respectively. Mean soil moisture contents and cacao evapotranspiration (ET_c) were 52, 45 and 28 % and 4.54, 3.19 and 2.32 mm/day while evaporation from soil area wetted by emitters (EW_z) were 5.65, 2.82 and 0.19 mm/day for respective IrT1, IrT2 and IrT3. The deficit irrigation strategies (IrT2 and 31

*Corresponding author: E-mail: soagele@futa.edu.ng, ohiagele@yahoo.com;

IrT3) imposed soil moisture deficit stress on cacao and produced lower pod and bean yields, it however enhanced water use efficiencies (25 and 44 %) and 30 and 50 % water savings. The study established suitable Pan coefficients for scheduling irrigation for cacao yield enhancement and drought (climate stress) amelioration.

Keywords: Cacao; Pan evaporation; crop coefficient; water use; seasonal transition; climate stress.

1. INTRODUCTION

Cacao (*Theobroma cacao* L.) is an important perennial fruit tree with an estimated annual world production of 3.2 million tonnes [1]. Within the cocoa-growing belt of West Africa, sale of cocoa beans is a major foreign exchange earner, the cocoa sector employs millions of smallholder farmers (small farm sizes ranging from 0.5 to 5.0 hectare (ha) and contributes about 70-100 % of their annual household incomes. In Nigeria, the main cocoa-producing areas are concentrated in the rainforest of the southern part of the country where an estimated 1.45 million hectares is cultivated. The productivity is 250 kg/ha, a yield level that is lower than those from Cote d'Ivoire and Indonesia (bean yields ranging from 600 to 1000 kg/ha respectively). In the smallholder cocoa farms of West Africa, farm sizes are small ranging from 0.5 to 5.0 hectare using low external inputs. The perennial fruit tree species of the rainforest of Nigeria: Cacao (*Theobroma cacao*), Coffee (*Coffea* spp) and Kola (*Kola* spp.) are characterized by deciduous growth habit but are cultivated under rainfed conditions [2,3].

Global warming and drought and other climate-related disasters are tied to the changing climate, extreme and variability of the weather [4,5]. The frequency and severity of drought event including (1.5 to 2 °C) warming are expected to increase in the near future as result of the decrease of regional precipitation and the increase in evapotranspiration driven by global warming [6,7,8]. Among natural hazards, drought ranks first in terms of the number of people directly affected [7]. The changing climatic events has implications for agriculture food security, economies, and welfare of the society and ecosystems [9]. Projections of climate change have pointed to an increase in mean temperature (about 2 °C by 2080) and potential/evapotranspiration, decrease in precipitation and crop (actual) evapotranspiration under future scenarios and uncertainties [7,8]. This implies future yield decreases which can be associated with enhanced heat and water stress under future climate conditions. Thus, climate Change (temperature and rainfall) scenarios for the

rainforest of Nigeria which have been variously constructed using climate models have indicated variabilities in rainfall pattern (amount, distribution, onset and cessation dates) and elevated maximum and minimum temperatures. These projected climatic changes will exacerbate soil moisture and thermal stresses during the dry season with implications for crop performance [7,8, 9].

Cocoa is cultivated as a rainfed crop, and it is highly sensitive to soil and weather conditions of low rainfall, soil and air moisture deficit and temperature stresses [2,10,3]. The changing growing environmental conditions (marginal soils and extreme weather events) impose constraints on cacao growth and productivity. In order to alleviate the constraints imposed by changing growing environmental conditions (marginal soils and extreme weather events) on cacao productivity, it is imperative to develop climatic-stress adaptive strategies for the fruit tree-based agroforestry systems of the rainforest tropics in the wake of changing climate/weather conditions (climate change and weather variabilities).

The FAO Penman-Monteith equation, is accepted worldwide as the standard method for estimating reference evapotranspiration (ET_o). The reference evapotranspiration (ET_o) is a measure of the evaporative demand of a given environment and thus crop consumptive water use which is the sum of evaporation from soil and plant transpiration from the field [11-14]. The procedures to calculate ET_o from radiation, wind, humidity and temperature data are presented in the FAO Paper No. 56. The standard procedure for estimating evapotranspiration is documented in the FAO I&D No. 56, where a list of K_c values for each crop and developmental stage is provided. This K_c approach has been used to obtain reference evapotranspiration (ET_o) and crop consumptive water use (ET_c) for arable, trees and vines [15,16]. The ratio between ET and ET_o, is defined as a crop coefficient (K_c). Thus, if K_c is known, the ET_c is calculated as:

$$ET_c = K_c ET_o \quad (1)$$

The FAO-56 dual crop coefficient approach [15,17] also describes the relationship between crop evapotranspiration (ET_c) and reference evapotranspiration (ET_o) by separating the single K_c into the basal crop (K_{cb}) and soil water evaporation (K_e) coefficients. In the FAO-56 single crop coefficient approach, the effect of both crop transpiration and soil evaporation are integrated into a single crop coefficient (K_c). However, crop evapotranspiration (ET_c) estimation is more accurate by dual crop coefficient approach than the single crop coefficient approach, the dual crop coefficient approach uses more parameters and take soil practices and crop characteristics into consideration [18,13, 19]. In the dual approach a daily basal crop coefficient (K_{cb}), representing primarily the plant transpiration, and a daily soil evaporation coefficient (K_e) are considered separately according to the equation:

$$K_c = K_{cb} + K_e \quad (2)$$

$$ET_c = (K_{cb} + K_e) ET_o \quad (3)$$

K_{cb} is a transpiration coefficient and K_e is an evaporation coefficient. The K_{cb} and K_e are the basal crop and the soil evaporation coefficients. For the dual K_c method, K_s applies only to transpiration (Tr) and provides actual transpiration (Ta). The values of K_s are obtained using its relationship with measured water stress indicators. The FAO procedure for estimating crop consumptive use requirements provides a list of K_c values for each crop and developmental stage. In the K_c approach both crop transpiration and soil evaporation are timely averaged into a single coefficient (K_c) commonly used to obtain the ET_c for various crops. The FAO No. 56 Publication offers the option of differentiating E from Tr by using a dual crop coefficient approach. The estimates of crop water requirements (ET_c) are derivable from the product of potential evapotranspiration of a reference crop (ET_o) using a crop factor (k_c). Based on these relations:

$$ET_c = ET_o * k_c \quad (4)$$

The crop coefficient k_c is estimated as:

$$k_c = ET_o/ET_c \quad (5)$$

The crop coefficient (K_c) is based on a theoretical understanding of the processes of transpiration and evaporation from a tall crop,

and assumes full crop cover or frequent wetting of the soil surface. Allen et al. [15] suggested a K_c value of 1.0–1.05 for a cocoa crop with a complete canopy.

Information is inadequate on the actual water use (ET_c) of cocoa on the field [20]. Estimated values of ET_c ranging from 3 to 6 mm/day during rains and less than 2 mm/day in the dry season have been reported for cocoa [11]. Field data (based on the sap flow method) suggest ET_c rates of less than 2 mm/day for cocoa crop with a complete canopy, this appear to be low compared with potential ET_o estimate of 3–5 mm d⁻¹ using Penman equation [11]. In a simulated El Niño drought experiment reported by Moser et al. [21] in Indonesia, there were no significant differences in cocoa leaf, stem and branch wood, or fine root biomass production between the rainfed control treatment and the one in which rain through-fall was reduced by 70–80% (for dry soil profile to permanent wilting point during the year). The combined average rate of water use by both cocoa and *Gliricidia sepium* (measured using heat dissipation sap flux sensors) was 1.3 mm. d⁻¹ in the protected plots reduced rain through-fall (rainfall reduction between 70–80 %) and 1.5 mm d⁻¹ in the control (70% of which was from the cocoa trees). These crop water use values were described as low [22]. Over a consecutive 18-day period, transpiration average was equivalent to 1.31 mm d⁻¹, (about 10 litres per tree per day) compared with a Penman potential ET_o estimate of 3–5 mm d⁻¹ which equates to a crop factor (K_c) of about 0.3 [11,21]. The reports of cocoa water use and yield production when grown as mixed crop with coconut and once a week irrigation during November-December, once every six days during January-March and once in four to five days during April-May with 175 l water tree⁻¹. Maximum yields were obtained when cocoa was drip irrigated with 20 l tree⁻¹ day⁻¹ and the total irrigation amount was 175 l per tree. Assuming a planting density of 1600 trees ha⁻¹ (2.5 m × 2.5 m) these figures equate to water use values of 5.6–7.0 mm d⁻¹ or 3.9–4.8 mm d⁻¹, and 2.2 or 3.2 mm d⁻¹ at 1100 trees ha⁻¹ (3 m × 3 m) under drip irrigation. No estimates of the yield benefits are given or the total quantity of water to be applied over a season. Based on field trials by Diczbalis et al. [23], the annual irrigation requirement was estimated as 470 mm, with peak weekly requirements of about 200 l tree⁻¹ (1250 trees ha⁻¹) while dry bean yields of between 1.5 and 2.7 t ha⁻¹ was achieved from young trees.

The annual total rainfall in the cocoa growing regions of Nigeria is about 1500 mm (less than 2000 mm). The rainfall distribution pattern is bimodal from April to July and September to November. There is a short dry period from July to August during which the relative humidity is still high with over cast weather conditions. There is a main dry season from November to February-March. The four to six months of dry weather results in soil water deficit and since irrigation is not part of the farming system, causing seedling mortality [24]. In bearing plants, the existence of the short dry season during main crop pod filling can affect bean size if it is sufficiently severe. In adult plantings, water deficits result in lower yields and an increase in the level of mirid (capsid) damage. In the rainforest cocoa growing belt of west Africa, fruit trees in plantations (cacao, kola, coffee, citrus species and oil palm) are seldom irrigated especially during the terminal drought situation of the dry season. Few studies had addressed the responses of cacao to dry season irrigation especially, the effects of irrigation on root zone moisture dynamics, tree water use, growth and yield in the premise of unfavourable weather constituted by soil moisture deficit and high temperature stresses of the dry season.

Given the changing environment regimes (soil and weather/climate) and increasing worldwide demand for cocoa, it is important to develop sustainable production systems based on sound agronomic practices such as irrigation to ameliorate the extreme weather conditions (hydrothermal stresses), improve its productivity and extend frontiers of its production to marginal weather and soil conditions. Few studies had addressed these features in tropical trees and very little is known about cacao the responses of cacao to dry season irrigation in the premise of unfavourable weather constituted by soil moisture deficit and high temperature stresses. In addition, information is inadequate on water use of cocoa and dynamics of soil moisture extraction as affected by irrigation regimes. Experiments were designed to examine the effects of regulated dry season irrigation on root zone moisture dynamics tree water use, and bean yield of cacao in a rainforest zone of Nigeria.

2. MATERIALS AND METHODS

2.1 Experimental Site and Conditions

An experiment was conducted on the Research and Experiment Station of the Department of

Crop, Soil and Pest Management, Federal University of Technology Akure, Nigeria. Akure is located in the rainforest zone of south west Nigeria on latitude: $7^{\circ} 18' N$, Longitude: $5^{\circ} 8' E$ and 350 m abs. Five to six years old fruiting cacao trees which had been previously irrigated during dry season from seedling establishment (April, 2013) till date were used.

The cocoa-growing rainforest belt of southern Nigeria, is characterized by wet and dry season transition, and the seasons have variable weather conditions. The annual rainfall range from 1500 to over 2000 mm distributed in a bimodal pattern within seven to eight months duration and 3 to 4 months of dry season. The dry season is a terminal drought situation characterized by inadequate rainfall,, soil moisture, high vapour pressure deficit and temperatures stresses and very clear sky (high intensity of solar radiation) [24, 3]. In the rainforest cocoa growing belt of west Africa, fruit trees in plantations (cacao, kola, coffee, citrus species and oil palm) are seldom irrigated especially during the terminal drought situation of the dry season.

2.2 Soil Characteristics and Moisture Determination

The soil of the site of experiment is sandy-clay-loam with relatively high water holding capacity. Available soil water in the upper 0.60 m of the soil depth is 187 mm. the percent and volumetric soil water contents at field capacity and permanent wilting point are 21 and 10 % respectively. Mean bulk density was 1.25 g cm^{-3} .

Soil samples were taken using soil Auger for water content measurement within the top soil layer (0 - 30 cm) by gravimetric method. Core samples were taken for bulk density and porosity measurement. Soil moisture content would attain field capacity in two days since the soil is sandy clay to silty clay loam [25]. The samples were taken two days after and just before the next irrigation. The difference in moisture content between the two sampling periods was taken to be the moisture used. That is, the evapotranspiration by the crop for that period. Since it was assumed that drainage was negligible (no drainage), the moisture change was principally attributed to evapotranspiration. Soil moisture depletion (SWD) was obtained from the differences in soil moisture contents (changes in soil moisture contents: ΔS) measured between two measurement period.

Soil moisture contents were determined weekly at 20 cm depths from soil samples taken with augers and core samplers.

2.3 Irrigation Strategies

Cacao trees were drip-irrigated based on levels of cumulative pan evaporation. Irrigation treatments were based on the restoration of cumulative Pan evaporation (EPan) using variable Pan coefficients (Kcp) of 100, 70 and 50 % [15, 26, 25]. The Pan coefficients (100, 70 and 50 % Kcp; the relative water deficit of 0, 0.3 and 0.5) indicated zero, high and low water stress conditions and respectively. Irrigation amount was calculated using Pan evaporation and Pan coefficients (Kcp1: 1.0; Kcp2: 0.7, and Kcp3: 0.5) according to Doorenbos and Pruitt (1975) and Allen et al. [15] as:

$$I_r = A * EPan * Kcp \quad (6)$$

where I_r is the amount of applied irrigation water (mm), A is the plot area, $EPan$ is the cumulative evaporation at irrigation interval (mm) and Kcp is the plant-pan coefficient.

Irrigation treatments were coded as $EPan * 100 Kcp$ (IrT1), $EPan * 70 \% Kcp$ (IrT2) and $EPan * 50 \% Kcp$ (IrT3) while irrigation was fixed at 5 days-interval for the three irrigation treatments. Irrigation treatment IrT3 had the maximum water deficit which was used to determine stressed baseline while IrT1 suggest adequate irrigation to meet full crop water requirements (the non-crop water stress baseline). Irrigation water was applied using gravity-drip irrigation system at 4.8, 6.8 and 9.6 l/tree/day at each irrigation via point source emitters of 2l/h discharge rate which were installed on laterals per row of cacao tree spaced at 3 x 3 m. One drip lateral served each plant row and an inflow meter was installed at the control unit to measure total flow distributed to all replications in each treatment. Irrigation buckets were suspended on 3.5 m high tank stands to provide the required hydraulic heads [25, 3].

Total amount (volume) of irrigation water applied per treatment was calculated using equation:

$$V = P * A * EPan * Kcp \quad (7)$$

where, V , is the volume of irrigation water (L); P , wetting percentage (taken as 100 % for row crops); A , is plot area (m^2); $EPan$ is the amount of cumulative evaporation for the irrigation interval (5-days) and Kcp Pan coefficients (1.0,

0.7 and 0.5). This corresponded to 7.14 mm (1.93 l/day), 10.7 mm (2.90 l/day), 14.28mm (3.86 l/day) for the respective 0.5, 0.7 and 1.0 Kcp . In order to attain good plant stand, a pre-treatment total of 135 mm of irrigation water was applied equally to all treatment plots in several applications, this replenished soil water in the 0.60 m profile depth to field capacity across treatments. Following the pre-treatments of 4.82 l/day for 5 days, differential irrigation treatments commenced on 13th December, 2017 and was terminated May 9th, 2018. The amount of water applied per irrigation and seasonal irrigation amount varied from a maximum of 4.82 l/day and 127500 mm (DI_1 level) to a minimum of 1.93 l/day and 20400 mm (DI_4 level). Irrigations continued until one week before the final harvest. Actual crop evapotranspiration (ET_c) of cacao under the irrigation amounts was calculated with the water balance equation (Equation 1) [25].

$$ET + I + P + \Delta S - D_p - R_f \quad (8)$$

where, ET , is actual crop evapotranspiration (mm); I , the amount of irrigation water applied (mm); P the precipitation (mm); ΔSW , changes in the soil water content (mm); D_p , the deep percolation (mm); R_f , amount of runoff (mm). Since the amount of irrigation water was controlled, deep percolation and run off were assumed to be negligible. Daily crop evapotranspiration was estimated using the pan evaporation data, pan factor and crop coefficient [12, 25]. Data for Pan evaporation ($EPan$) used for the experiment were obtained from measurements with Class-A Pan (121 cm in diameter and 25.5 cm in depth) from the Meteorological Station, Department of Meteorology & Climate Science, FUTA) located near the plots.

Deep percolation was considered as zero because there was no high underground water problem in the area. If available water in the root zone (0–90 cm) and total applied water amount by irrigation were above the field capacity, it would be assumed that water amount above field capacity leaked into the deeper soil zones and was called deep percolation (D_p : available total water amount at 0–90 cm soil depth before irrigation + applied irrigation water field capacity) [12]. Total water requirement (WR) was determined using the relation:

$$WR = A \times B \times C \times D \times E \quad (9)$$

where: WR = Water requirement (l per day /plant) A = Open Pan evaporation (mm/day) B =

Pan factor (1.0, 0.7 and 0.5), C = Spacing of plant (m²), D = Crop factor (factor depends on plant growth, value for fully grown cacao = 1.13 but for cacao in the early fruiting stage, 0.83 was adopted). Water requirements (WR) were 9.63, 6.75 and 4.8 l/plant/day for the respective IrT1, IrT2 and IrT3 irrigation treatments.

Irrigation water requirement is determined using average season wise pan evaporation data for the area. The total water requirement (TWR) of the farm plot was obtained using the relation. Therefore, the total water requirement (TWR) of the farm plot is:

$$TWR = WR \times \text{No. of Plants} \quad (10)$$

Maximum allowable deficit (MAD) for cacao (50% of available water storage capacity of the soil (AWC) Gross irrigation requirement (GIR) of an orchard or vineyard, the computed ET_c, which is considered as the net irrigation requirement (NIR), should be divided by the application efficiency (AE).

$$GIR = NWR/AE \quad (11)$$

Yield and crop water use were deployed to evaluate appropriate the efficiencies of irrigation management practices among the different irrigation strategies adopted.

2.4 Orchard Water Use Efficiencies

Water productivity (Irrigation water use efficiency (IWUE) and crop water use efficiency (WUE)) was determined based on the methods of Sezen et al. [26] and Agele et al. [25] as:

$$IWUE = \frac{\text{Biomass weight (Y)}}{\text{total irrigation water applied (Ir)}} \quad (12)$$

$$WUE (\text{crop}) = \frac{\text{Biomass weight (Y)}}{\text{Cumulative seasonal Etc}} \quad (13)$$

where IWUE is the irrigation water use efficiency (t.ha⁻¹ mm), EY is the economical yield (t.ha⁻¹), Ir is the amount of applied irrigation water (mm).

Cacao water requirement was determined using FAO-56 single and dual crop coefficient models approach. The aim was to analyze the capacity of the FAO-56 single and dual crop coefficient models to assess cacao evapotranspiration and water requirements (estimating adequacy of irrigation amount for cacao). The FAO-56 dual crop coefficient approach [15] describes the

relationship between crop evapotranspiration (ET_c) and reference evapotranspiration (ET₀) by separating the single K_c into the basal crop (K_{cb}) and soil water evaporation (K_e) coefficients, while in the FAO-56 single crop coefficient approach, the effect of both crop transpiration and soil evaporation are integrated into a single crop coefficient (K_c). Cacao orchard irrigation was scheduled from transpiration and evaporation coefficients (K_{cb}, K_e). FAO I&D No. 56 publication offers opportunity for differentiating E from Tr by using a dual crop coefficient approach, according to the equation:

$$ET_c = (K_{cb} + K_e) ET_0 \quad (14)$$

where K_{cb} is a transpiration coefficient and K_e is soil evaporation coefficient.

K_{cb} is basal crop coefficient (k_{cb} = ET_c/ET₀)

$$K_c = k_{cb} + k_e \quad (15)$$

and then,

$$ET_c = (k_{cb} + k_e) ET_0 \quad (16)$$

2.5 Size of Cacao Canopies

Tree canopies may be characterized using two parameters: canopy volume (m³ of tree volume/ of ground surface: m³/ m²) and leaf area density (m² of leaf area /m³ of tree volume). Tree canopy can be measured with a measuring rod once the tree shape has been approximated as a sphere, an ellipsoid, or a truncated inverted cone. As an alternative to the measurements or calculations of the radiation actually intercepted by the tree, a simple parameter that is easy to determine is the degree of ground cover. The ground cover (normally expressed in percentage) is obtained by measuring the shaded area outlined from the horizontal projection of the tree canopy

The ground cover (normally expressed in percentage) was obtained by measuring the shaded area outlined from the horizontal projection of the tree canopy

$$A = \frac{\pi d^2}{4} \quad (m^2) \quad (17)$$

d = diameter of shaded area by cacao canopy (2m), A is per cent ground cover by cacao canopy; Tree spacing is 3x3 m (9 m²); d₁ (areal canopy area), d₂ (height bt d₁ and d₃); d₃ (projection of canopy area on ground, d₁ > d₃).

$$\text{Canopy volume} = \frac{1}{3}\pi d^2 \left[\frac{d_1 x^2}{4} + \frac{d_3 x^2}{4} + d_1 d_3 \right] \quad (18)$$

$$K_{ext} = 0.52 + 0.00079dp - 0.76e^{-1.25DAF} \quad (23)$$

$$DAF = 2 - 0.53(Vu - 0.5) \quad (24)$$

2.6 Soil Surface Evaporation as Affected by Irrigation

Soil evaporation from surface not wetted by emitters (Edz: the rest of the soil surface outside the emitter wetting pattern) and surface evaporation from the soil wetted by the emitters (emitter wetted zone (EWz) were estimated.

The equations derived from the study on olive orchard ET by Orgaz et al. [27] was adapted for estimating values of Ewz and Wdz.

Evaporation (EDz) from soil not wetted by emitters (EDz)

$$Edz = K_s, e E_{To} \text{ (mm/day)} \quad (19)$$

Where G is ground cover fraction of tree canopy, is monthly rainfall amount, wz is fraction of soil surface wetted by drip emitters (E_{To} = reference ET). EWz of 0.6E_{To} for dense crop cover/plant density; EWz is the of soil surface kept wet by emitters, and Cover crop coefficient varies from 0.25 to 0.8.

Evaporation (EDz) from soil not wetted by emitters (EDz)

$$Edz = K_s, e E_{To} \text{ (mm/day)} \quad (20)$$

Where G is ground cover fraction of tree canopy, is monthly rainfall amount, wz is fraction of soil surface wetted by drip emitters (E_{To} = reference ET)

Cover crops/weed cover transpiration (Tr cc).

Weed cover up to 2 m in a 3 m row spacing

2.7 Transpiration (Tr) of Cacao Orchard

Cacao is deciduous (partly evergreen in some cases), its crop coefficient (K_c) and Tr were determined using the methods of Orgaz et al. [27].

$$K_c, Tr = (QdF_1) F_2 \quad (21)$$

$$Qd = 1 - e^{-k_{ext} Vu} \quad (22)$$

where K_{ext} is radiation extinction coefficient.

where DAF must be > 2); Vu = Vo(dp/10000), Vo = 1/6xD²H

E (exponent = 2.718), H (height of canopy; m); D is average canopy diameter, m); Vo is canopy volume ; m³/tree); Vu is canopy volume as amount on ground cover; m³/m²); DAF is leaf area density; dp is tree density; number/ha), Fi = 0.07 for tree density greater than 300 trees/ha), F2 is monthly coefficient of Tr which is about 0.7 to 1.0 from wet to dry seasons

$$Qd = 1 - e^{-k_{ext} Vu}$$

$$K_c, Tr = (QdF_1) F_2 \quad (25)$$

$$ET_c = E_{To} K_c K_{r,t} \quad (26)$$

K_{c,t} is empirical coefficient relating the ET of an orchard of incomplete cover to a mature orchard of full canopy cover. In addition, K_{r,t} relates to horizontal projection of tree shade/canopy (ground cover per cent; Orgaz et al. [27]), and K_{r,t} is about 0 to 70% of G.

2.8 Cacao Orchard Transpiration (Tr) was Determined as

$$Tr = K_c, Tr E_{To} \quad (27)$$

Where K_c, Tr is transpiration coefficient which varies bt 0.75 to 1.0 seasonally until leaf senescence onset for a fully wetted orchards (sufficient soil moisture situation) (K_c,Tr decrease at senescence and recovers at the onset of rainfall.

3. RESULTS AND DISCUSSION

3.1 Weather Conditions during Period of Study

The late (minor) rainy season (mid August to December) is characterized by high cloud overcast (overcast sky), low air temperatures and higher relative humidity compared with the major rainy season (April to mid August) and the dry season (Fig. 1). On the average, the rainy season had higher mean relative humidity averaged (71 %) and lower air temperatures (32.8 °C) compared with the dry season (December to March). Also, higher air temperature and VPD and lower relative humidity

were found for the unshaded open sun cacao compared with the shaded plants.

A low pressure gravity-drip system was deployed to deliver water to cacao rootzone which alleviated moisture stress during the dry season. Across sampling dates, different amounts of irrigation water were applied based on cumulative Pan evaporation (EPan)*Pan coefficients for the respective irrigation treatments IrT1 (EPan.kc:1.0), IrT2 (EPan.Kc:0.7) and IrT3 (EPan.Kc: 0.5). Irrigation amounts on monthly averages were 1009.88, 706.91 and 504.94 mm while seasonal totals were 2116.5, 8482.95 and 6059.25 mm for IrT1; IrT2 and IrT3 treatments. Well irrigated treatment (IrT1, EPan.kc:1.0) had highest amount delivered and lowest by IrT3 (EPan.kc:0.5) (Fig. 2). The deficit irrigations (IrT2 and IrT3) delivered 79 and 68 % respectively water to cacao rootzone compared with the well irrigated condition. Peak and significantly higher values of irrigation amount were delivered at DOY 45, 60, 75 and 90, periods which coincided with highest EPan values (> 5 mm/day). The irrigation treatments (IrT1, IrT2 and IrT3) affected soil moisture contents within cacao root zone. Soil moisture contents adequately reflected the irrigation water delivered across measurement dates (Fig. 3). Highest soil moisture contents were obtained for well irrigated (IrT1) and lowest for deficit irrigation (IrT3) treatment. For the respective deficit irrigation treatments (IrT2 and IrT3: 0.7 and 0.5 EPan coefficients), average soil moisture contents were 61, 48 and 42 % for IrT1, IrT2 and IrT3 irrigation treatments (Fig. 3). Highest soil moisture contents and crop evapotranspiration (ETc) were obtained from well irrigated plots (IrT1: EPan*kcp (1.0) (9.6 l/tree/day) followed by IrT2 (EPan*kcp (0.7) (6.8 l/tree/day) and lowest for IrT3 EPan*kcp (0.5) (4.8 l/tree/day). The deficit irrigation treatments (IrT2 and IrT3) had lower soil moisture contents (14.7 and 11.8 %) which equated to 30 and 50 % water savings.

Declines in soil moisture contents were obtained from DOY 345 to DOY 60, followed by increasing trends in soil moisture from DOY 75 till end of measurement (DOY 150). Declining trends in the values of soil moisture contents may be attributed to the increasing intensities in climatic demand (high vapour pressure deficits). Unfavourable weather of high temperatures and soil evaporation and low atmospheric humidity would enhance soil moisture depletion thus the low soil moisture status [28,9]. Increases in moisture were observed from DOY 75 till end of

measurement (DOY 150) can be attributed to rainfall received following its commencement (Mid March). In general, the observed trends in the status of rootzone moisture is attributable to the prevailing weather conditions denoted by increasing intensities of climatic demand (vpd) and temperatures during periods (DOY 345 to 60) of experiment.

3.2 Cycle of Soil Water before and after Irrigation

Soil water contents were measured using soil samples within the 0 - 20 cm soil profile depth before and one day after each irrigation. Soil moisture contents across measurement days ranged between wilting point (140 mm) before irrigation and field capacity (260 mm) after irrigation (data not shown). For the low and high water stress conditions (IrT2 and IrT3), soil moisture was often close to wilting point before each irrigation. For the deficit irrigation treatments (IrT2 and IrT3: 0.7 and 0.5 Pan coefficients), available water fell below 50% more often than not during the period of study. Because much more water was applied under high Pan coefficients (Kcp 1.0), soil moisture contents of well watered treatments (IrT1) was higher compared with deficit irrigation (IrT2 (0.7 Kcp: and IrT3 0.5 Kcp) treatments. The well watered treatment (IrT1), most times, maintained soil moisture within field capacity range.

In general, based on the values of soil moisture, the stored water within crop rootzone profile was used up between irrigation cycles. This is attributable to the intensities of climatic stress (high temperatures and vapour pressure deficits) which presumably enhanced soil evaporation and the rapid depletion of water stored in the soil profile. Soil moisture content immediately following irrigation gradually decreased towards next irrigation event, this situation confirms the inability of soil moisture reserve to satisfy cacao water demand during the dry season which was consistent with earlier reports of Famuwagun et al. [24] and Charles et al. [3].

Soil moisture depletions over two measurement days were deployed to determine cacao water use (ETc). Cacao water use (ETc) differed across measurement dates and irrigation treatments (Fig. 4a). Average cacao evapotranspiration (ETc) were 139, 97 and 63 mm/day for the respective IrT1 (IrT1 (Kc:1.0), IrT2 (Kc 0.7) and IrT3 (Kc 0.5). Cacao evapotranspiration (ETc) for the deficit irrigation

treatments (IrT2: 0.7 and IrT3:0.5 EPan coefficients) were about 45 and 70 % less than values for adequate irrigation (IrT1) treatment. Peak ETc values were obtained at DOY 45, 60 and 75 possibly due to high EPan (> 5 mm/day), lowest for DOY 120 and 135 with increases afterwards. The increases in cacao water use (ETc) from DOY 135 afterwards are attributable the commencement of rainfall and associated replenishment of soil moisture,

lowering of temperatures (air and soil) and high atmospheric humidity (declining atmospheric demand). The well watered treatment (IrT1) had highest ETc and the more deficit irrigation (IrT3) had least cacao water use (Fig. 4a). The mean ETc across measurement dates were 5.07, 3.55 and 2.63 mm/day for IrT1, IrT2 and IrT3 irrigation treatments for the period of experiment (December to May).




Fig. 1. Weather variables during period of experiment (temperature, rainfall, humidity, vapour pressure deficit and Pan evaporation)




Fig. 2. Time course of irrigation water applied during the period of experiment (December to May). The three irrigation treatments were coded as: IrT1 (EPan*1.0 kcp), IrT2 (EPan *0.7kcp) and IrT3 (EPan * 0.5 kcp). EPan is Pan evaporation and kcp is Pan




Fig. 3. Dynamics of soil moisture contents as affected by irrigation treatments. The three irrigation treatments were coded as:IrT1 (EPan*1.0 kcp), IrT2 (EPan *0.7kcp) and IrT3 (EPan * 0)

In addition to single crop coefficient ($k_c = 1.31$), cacao water requirement (ET_c) was also computed using the dual (k_r : 1.04) crop coefficient (Fig. 4b). Means of cacao water use for dual crop coefficient across measurement dates were 5.2, 3.7 and 2.8 mm/day for IrT1, IrT2 and IrT3 irrigation treatments for the period of experiment (December to May). The time course of cacao water use estimated using both the single and dual crop coefficients are presented in

Fig. 5a, b and c. Results showed similar trends in cacao ET_c for both methods and irrigation treatments while values were higher for the dual coefficient compared with the single k_c (Fig. 5a, b and c). The decreasing order of ET_c for single k_c and dual k_c were $IrT1 > IrT2 > IrT3$. Crop evapotranspiration (ET_c) increased with increases in the volume of irrigation water applied, this modified values of ET_c obtained for both the single and dual k_c approaches.




Fig. 4a. Cacao water use (evapotranspiration; ET_c) Fig. 4a. Time course of cacao water use (evapotranspiration; ET_c). The three irrigation treatments were coded as:IrT1 (EPan*1.0 kcp), IrT2 (EPan *0.7kcp) and IrT3 (EPan * 0.5 kcp). EPan is Pan evaporation




Fig. 4b. Time course of cacao water use (evapotranspiration;ETc). The three irrigation treatments were coded as:IrT1 (EPan*1.0 kcp), IrT2 (EPan *0.7kcp) and IrT3 (EPan

The FAO-56 dual crop coefficient approach [15] which describes the relationship between crop evapotranspiration (ETc) and reference evapotranspiration (ETo), separates the crop coefficient (Kc) into the basal crop (Kcb) and soil water evaporation (Ke) coefficients. In the single crop coefficient approach, the effect of both crop transpiration and soil evaporation are integrated into a single crop coefficient (Kc) while in the dual coefficient approach, a daily basal crop

coefficient (representing plant transpiration: Kcb) and daily soil evaporation coefficient (Ke) in the form of $Kc = Kcb + Ke$). However, crop evapotranspiration (ETc) estimation is more accurate by dual crop coefficient approach than the single crop coefficient approach, the dual crop coefficient approach uses more parameters and take soil management practices and crop characteristics into consideration [15].




Fig. 5a. Time course of cacao water use (ETc) by single and dual crop coefficients (kc) : kc (1.13), kr t (1.04) for well irrigation treatment (IrT1; EPan*100 kcp)




Fig. 5b. Time course of cacao water use (ETc by single and dual kc: kc (1.13), kr t (1.04) IrT1:1.0 EPan and EPan (IrT1; EPan*0.7 Kcp)




Fig. 5c. Time course of cacao water use (ETc by single and dual kc: kc (1.13), kr t (1.04) IrT1:1.0 EPan and EPan (IrT1; EPan*.0.5 Kcp)

The magnitudes of cacao ETc (single and dual crop coefficients obtained from the respective irrigation treatments followed from the irrigation water delivered (Fig. 5a, b and c). The irrigation regimes affected soil moisture contents and thus, its availability to meet crop water use. The values of cacao water use obtained from the respective irrigation treatments would have followed from the irrigation water delivered. Irrigation under well watered treatment increased tree water use and soil moisture status compared with deficit irrigation treatments (IrT2 and IRT3) which is

consistent with reports on citrus by Yang et al. [29]. The magnitude of cacao ETc obtained in this study are within the range of those reported in literatures [20]. Cacao water use (ETc) values ranging from 3 to 5 mm/day during rains and less than 2 mm/day in the dry season has been reported under irrigation regime of 10 litre/tree/day [11], Kohler et al. [22] obtained cacao ETc of 2 mm/day and Moser et al. [21] obtained 1.3 – 1.5 mm/day in Indonesia. Cacao transpiration average of 1.31 mm d⁻¹ (about 10 litres per tree per day) and ETc computed with a

Penman potential ETo of 3 – 5 mm d⁻¹ have been reported [21]. This value equates to crop factor (Kc) of about 0.3 [11,21]. Field data (based on the sap flow method) suggest ETC rates of

less than 2 mm/day for cocoa crop with a complete canopy, this appear to be low compared with potential ETo estimate of 3–5 mm d⁻¹ using Penman equation [11].




Fig. 6a. Ratio of cacao water use (ETc) to Pan evaporation (EPan) as affected by irrigation regime. The irrigation treatments were coded as: IrT1 (EPan*1.0 kcp), IrT2 (EPan *0.7kcp) and IrT3 (EPan *0.5kcp). EPan is Pan evaporation



Fig. 6b. Ratio of cacao water use (ETc) to irrigation water applied. The irrigation treatments were coded as: IrT1 (EPan*1.0 kcp), IrT2 (EPan *0.7kcp) and IrT3 (EPan *0.5 kcp). EPan is Pan evaporation




Fig. 7. Time course of monthly evaporation from wetted (EDz) and non-wetted (EWz) within cacao field. Monthly evaporation from wetted (EDz) and non-wetted (EWz) soil within cacao field during period of experiment (December to May)

The $ET_c/EPan$ ratio denotes the proportion of climatic water demand satisfiable by crop water use (ET_c). Among the irrigation treatments. The proportions of Pan evaporation ($EPan$) to cacao water use (ET_c) denoted as $ET_c/EPan$ ratio, differed across measurement dates and irrigation treatments. The means of $ET_c/EPan$ ratios across measurement dates were 1.016, 0.714 and 0.492 for the respective IrT1, IrT2 and IrT3 treatments (Fig. 6a). $ET_c/EPan$ curves were similar but the ratios ranged from 1.16 to 0.50 which indicated that both climatic demand ($EPan$) and cacao water consumption (ET_c) were high during the dry season at the site of study. Although a weak relationship, linear regression equation was fitted to the $ET_c/EPan$ trends as: $Y = 0.011x + 0.94$, $R^2 = 0.32$). The ratio of water use (ET_c) to irrigation water applied denotes the proportion of irrigation water applied used for crop evapotranspiration. Trends of ET_c to irrigation were similar among irrigation treatments and measurement dates but values differed among irrigation treatments (Fig. 6b). The mean values were 1.016, 0.714 and 0.492 for the respective IrT1, IrT2 and IrT3 treatments (Fig. 6b). Generally, the ratios ranged from 1.13 to 0.27, 0.79 to 0.19 and 0.57 to 0.14 which indicated differences in the ability of irrigation water applied to satisfy climatic demand ($EPan$) driven trends of cacao water

consumption (ET_c). When soil water is readily available to a crop, the rate of water evaporation from an Evaporation Pan is proportional to the rate of crop water use (Doorenbos and Kassam, 1979) [15]. Doorenbos and Kassam (1979) found positive linear and significant logarithmic correlation ($P < 0.01$) between ET_c and $EPan$ while Smajstrla et al. [30] obtained significant logarithmic correlation ($P < 0.01$) between ET_c and $EPan$. These reports confirmed the established close relationship between plant water consumption and Pan evaporation.

3.3 Soil Evaporation from Cacao Orchard

The ET_c from an orchard is more complex. In addition to tree Tr , there could be Tr losses from cover crop or from weeds, and there are E losses from the soil. Under irrigation conditions, there are two E components that may differ in their rates: one is the E from the soil areas wetted by the emitters, and the other is the E from the rest of the soil surface which is only wetted by rainfall. Soil evaporation was respectively estimated for the wetted zone (Edz) and the non-wetted zone (Ewz) during the period of experiment. The mean values for soil evaporation for the wetted (Edz) were 5.65, 2.82 and 0.19 mm/month for the respective IrT1, IrT2

and IrT3 treatments. The seasonal totals soil evaporation for the wetted (Edz) and the non-wetted (Ewz) zones were 234.29 and 33.94 mm respectively. Based on the cumulative seasonal totals, soil evaporation for the wetted zone (Edz) was averagely 7 times compared with the non-wetted zone (Ewz) (Fig. 7). Irrigation replenished soil moisture depletion while cacao canopy produced cover to soil and a more favourable microclimate around the canopy spread. This is appear to explain the magnitudes of soil evaporation from the wetted (Edz) and the non-wetted (Ewz) zones within cacao field [8]. Studies also indicated that the conditions at the soil surface due to (i) the percentage of soil surface wetted via irrigation, (ii) the irrigation intervals and (iii) the soil exposure to light determine the dynamics of Tr and Es in orchards [31,8].

In this study, drip irrigation was deployed to replenish moisture depletion from cacao root zone. There were spatial variations in the degree of wetting within the orchard; some areas are frequently wetted by the emitters while the rest of the soil surface remains dry in the absence of rainfall. The drip lines were placed near the trees while the wetted areas are shaded by the cacao canopy. The effects of orchard canopy and drip irrigation appeared adequate to alleviate radiation-limited soil water evaporation (E). Measurements and models suggest that E from the soil surface in orchards, which are wetted frequently (every 1-2 days) by emitters is equivalent to about 60 percent of the ETo from the wet areas [31, 32]. As a first approximation, the quantification of E from the wetted spots in a drip-irrigated orchard can be made using a semi-empirical model of Bonachela [31] of the relation:

$$Es = 0.6 ETo \quad (28)$$

Total pod and bean yields were highest for IrT1 (35.4 and 2.29 t.ha⁻¹) followed by IrT2 (22.1 and 1.37 t.ha⁻¹) and lowest (10.3 and 1.03 t ha⁻¹) for IrT3. Bean yields decreased by 60 and 40 % under IrT3 and IrT2 compared with IrT1 (Table 1). Deficit irrigations however produced 30 and 50 % water savings compared to well watered treatment (IrT1). Water productivity was affected by irrigation regimes. Water use efficiencies values ranged between 0.3 and 0.04 t/mm for the ratio of yield to evapotranspiration (Y/ETc) and 0.16 to 0.19 kg/mm for ratio of yield to irrigation amount (Y/Irrig) respectively (Table 1).

The yields of pods and beans were significantly higher in IrT1 treatments compared with IrT2

while they were lowest significantly for IrT3. However, Carr [20] and Charles et al. [3] had reported that fruit yield does not only depend on irrigation amount but a function of other management practices adopted and soil properties such as infiltration rate, and water holding capacity.

Irrigation effect was significant both on number and weight of pods and beans in cacao. Evaluation of irrigation amount and frequency should not only consider fruit yield and yield components, but also consider WUE [32, 3]. Other studies have reported irrigation effects on biomass, pod and bean yields of cacao. Dicbalis et al. (2010) working in Australia examined effects of seasonal irrigation requirement of 470 mm and 200 l/tree for weekly irrigation and obtained resultant bean yields of 1.5 – 2.7t/ha. Based on field trials by Diczbali et al. [23], the annual irrigation requirement was estimated as 470 mm, with peak weekly requirements of about 200 l tree⁻¹ (1250 trees ha⁻¹) while bean yields of between 1.5 and 2.7 t ha⁻¹ were obtained as achieved from young trees.

Cocoa is cultivated as a rainfed crop but sensitive to weather extremes of low rainfall, soil moisture deficit and high temperature stresses had been variously reported in the literature [2, 10, 33,34, 3]. Global warming (including 1.5 to 2 °C), drought and other climate-related disasters are tied to the changing climate, extreme and variability of the weather [5, 4]. These situations would drive decreases of regional precipitation and increase in evapotranspiration driven [6, 8, 9]. These conditions may enhance yield decreases associated with enhanced heat and water stress under present and future climate conditions. The climate models of the rainforest of Nigeria have been variously constructed [35]. The results indicate that the projected climatic changes will exacerbate soil moisture and thermal stresses with implications for crop performance [9]. As precipitation becomes more variable and unpredictable in addition to the expected increased warming due to changing climate, development of water-saving management practices for sustainable agriculture now and in the future is envisaged [6,8,9]. Establishing the optimal irrigation scheduling is important in the development of water-saving practices for sustainable cacao production and climate stress alleviation in the wake of the hydrothermal (extreme heat and water deficits) stresses envisaged under future climate.

Table 1. Summary of measured soil and cacao variables as affected by irrigation regimes. Irrigation treatments were: IrT1: EPan*1.0 kcp, IrT2: EPan *0.7 kcp and IrT3: Epan * 0.5 kcp

Irrigation	Seasonal irrigation	Scm (%)	ETc Single (kc:1.13)	ETc Dual (kr t: 1.04)	ETc/EPan ratio	Seasonal ETc	No.Pods/plant	Pod wgt /plant (kg)	No. Beans/plant	Bean Wgt/Plant (g)	Bean wgt (kg/ha)	WUE (Irr)	WUE (ETc)	EWz
IrT1	33858.2	21.4	5.07	5.2	0.92	139.1	15.4	4429	114	396.5	4.41	0.0117	0.032	5.65
IrT2	32705.3	17.3	3.55	3.7	0.73	97.3	12.3	3125	102	334.3	3.72	0.0142	0.043	2.82
IrT3	16929.4	14.4	2.63	2.8	0.56	62.7	9.8	2673	91	308.1	3.42	0.0182	0.055	0.19
LSD		4.1	1.8	1.6	0.21	17.3	2.7	112.3	5.4	23.8	0.25	0.003	0.005	1.33

(0.05)

Abbreviations: ETc::crop water use/evapotranspiration), ETc (single): ETc using single coefficient, ETc (dual): ETc using dual coefficient, WUE: water use efficiency, EWUE(Irrig): WUE computed as ratio of bean yield to irrigation amount(t/mm/ha), WUE (ETc): WUE computed as the ratio of bean yield to crop water use (t/mm/ha), EWz (surface evaporation from soil area wetted by emitters (emitter wetted zone). Following data analysis, the statistical significance of irrigation treatment on the measured variables was indicated by the LSD while the significance of differences between treatment means were explored at $P < 0.05$. Irrigation requirement was based on cumulative class A Pan evaporation and Pan coefficients (kcp)

The site of study in the rainforest zone of Nigeria is characterized by bi-modal rainfall pattern and the wet-dry season transition. The rainfall distribution pattern is bi-modal from April to July and September to November. The dry season which span December of a year to April of the other, is a terminal drought situation characterized by inadequate rainfall, soil moisture deficit, high vapour pressure deficit and temperatures and very clear sky (high intensity of solar radiation [36, 24, 3]. Such unfavorable weather condition will enhance hydrothermal stresses, evapotranspiration, leaf senescence, branch and twig die-back and even tree mortality [24, 3, 37-43].

4. CONCLUSIONS

The variable irrigation based on variable Pan coefficients (1.0, 0.7 and 0.5) affected dynamics of root zone moisture, cacao water use (ET_c) and yields. Study established suitable Pan coefficients for scheduling irrigation during the terminal drought situation of the dry season in a rainforest zone of Nigeria. Relative to adequate irrigation (I_rT1), deficit irrigations (I_rT2 and I_rT3) produced lower soil moisture, water use (ET_c) pod and bean yields which were however accompanied by 25 to 44 % increases in water use efficiencies and 30 and 50 % water savings. The water saving advantage of deficit irrigation strategies can be scaled up for adoption. The low pressure gravitydrip irrigation system alleviated climate stress during the dry season and improved cacao performance. Findings will be of relevance to adaptation strategy in the premise of the changing climate, extreme and variability of weather while measured cacao water use (ET_c) will find use for improving irrigation models for fruit trees.

DISCLAIMER

This paper is an extended version of a preprint document of the same author.

The preprint document is available in this link: <https://assets.researchsquare.com/files/rs-1306540/v1/8bb25b4b-d50d-4356-b5de-64c49b057038.pdf?c=1643900011>.

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

1. Food and Agriculture Organization (FAO:2012) FAOCLIM: Environment and natural resources service. Working Papers. 2012;5. FAO, Rome
2. Opeke LK. Tropical commodity crops. Spectrum Books Ltd, Ibadan, Nigeria. 2006;213.
3. Charles EF, Agele SO, Aiyelari OP, Famuwagun IB, Faboade E. Shade and irrigation effects on growth, flowering, pod yields and cacao tree survival following 5 years of continuous dry season irrigation. *International Journal of Environment and Climate Change*. 2019;10(7):54-64. DOI:<https://doi.org/10.9734/ijeccc/2020/v10i730211>
4. United Nations Framework Convention on Climate Change (UNFCCC) 2004. the 10th Conference of the Parties (COP10). Buenos Aires, Argentina. Available:<https://unfccc.int/>
5. Intergovernmental Panel on Climate Change (IPCC). *Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Core Writing Team, R.K. Pachauri and L.A. Meyer (eds.)]. IPCC, Geneva, Switzerland. 2014;151.
6. Sheffield J, Wood EF, Michael L, Roderick ML. Little change in global drought over the past 60 years. *Nature*. 2012;491:435–438.
7. Trenberth KE, Dai A, van der Schrier G, Jones PD, Barichivich J, Briffa KR, Sheffield J. Global warming and changes in drought. *Nature Climate Change*. 2014;4:17–22.
8. Tombesia S, Frionia T, Ponia S, Palliottib A. Effect of water stress “memory” on plant behavior during subsequent drought stress. *Environmental and Experimental Botany*. 2018;150(2018):106–114.
9. Agele S. Global warming and drought, agriculture, water resources, and food security: impacts and responses from the tropics. In: *Handbook of Climate Change Management*. (Leal Filho W, Luetz J, Ayal D. eds.) Springer, Cham; 2021. DOI:https://doi.org/10.1007/978-3-030-22759-3_183-1
10. Zuidema PA, Peter A, Leffelaar PA, Gerritsma W, Mommer L, Niels PR, Anten NPR. A physiological production model for

- cocoa (*Theobroma cacao*): model presentation, validation and application Agricultural Systems. 2005;84(2005):195–225.
11. Penman HL. Natural evaporation from open water, bare soil and grass. Proc R Soc Lond A. 1948;193:120–145.
 12. Doorenbos J, Pruitt WD. Crop water Requirements (revised). Food and Agriculture Organization of the United Nations, Irrigation and Drainage Paper 24, Rome. FAO; 1977.
 13. Ventura F, Spano VF, Duce P, Snyder RL. An evaluation of common evapotranspiration equations. Irrigation Science. 1999;18:163-170.
 14. Allen RG, Pereira LS, Smith M, Raes D, Wright J. FAO-56 dual crop coefficient method for estimating evaporation from soil and application extensions. J. Irrig. Drain. Eng. 2005;131:2–13.
 15. Allen RG, Pereira LS, Raes D, Smith M. Crop evapotranspiration: Guidelines for computing crop water requirements. FAO Irrigation and Drainage Paper 56; FAO—Food and Agriculture Organization of the United Nations: Rome, Italy. 1998;300.
 16. Ferreira MI. Stress coefficients for soil water balance combined with water stress indicators for irrigation scheduling of woody crops. Horticulturae. 2017;3:38. DOI:10.3390/horticulturae3020038
 17. Allen RG, Pereira LS. Estimating crop coefficients from fraction of ground cover and height. Irrig. Sci. 2009;28:17–34.
 18. Monteith JL. Principles of environmental physics. Arnold, London. 1973;291.
 19. Zhang B, Liu Y, Xu D, Zhao N, Lei B, Rosa RD, Paredes P, Paço TA, Pereira LS. The dual crop coefficient approach to estimate and partitioning evapotranspiration of the winter wheat—summer maize crop sequence in North China Plain. Irrig. Sci. 2013;31:1303–1316.
 20. Carr MCV. The water relations and irrigation requirements of cocoa (*Theobroma cacao* L.): A review. Expl Agric. Cambridge University Press. 2011;47(4):653–676. DOI:10.1017/S0014479711000421
 21. Moser A, Leuschner C, Hartel D, et al. Response of cocoa trees (T cacao) to a 13-month dessication period in Sulawesi, Indonesia. Agroforestry Systems. 2010;79:171-187.
 22. Kohlerlscher M, Schwendenmann L, Holscher D. Throughfall reduction in a cacao agroforest.: tree water use and soil water budgeting. Agric & Forest Meteorol. 2010;150:1079-1089.
 23. Diczbalis Y, Lemm C, Richards N, Wicks C. Producing Cocoa in Northern Australia. Australian Government, Rural Industries Research and Development Corporation Report 09/092; 2010.
 24. Famuwagun B, Agele S, Aiyelari P. Shade effects on growth and development of cacao following 2 years of continuous dry season irrigation. International Journal of Fruit Science. 2017;18(7):1-24.
 25. Agele SO, Anifowose AY, Agbona AI. Irrigation scheduling effects on components of water balance and performance of dry season fadama-grown pepper in an inland-valley ecosystem in a humid tropical environment. International Journal of Plant and Soil Science. 2014;4(2):171-184.
 26. Sezen SM, Celikel G, Yazar A, Tekin S, Kapur B. Effect of irrigation management on yield and quality of tomatoes grown indifferent soilless media in a glasshouse. Scientific Research and Essay. 2010;5:41-48.
 27. Orgaz F, Testi L, Francisco J, Villalobos LJE, Fereres E. Water requirements of olive orchards-II: Determination of crop coefficients for irrigation scheduling Irrigation Science. 2006;24(2):77-84. DOI: 10.1007/s00271-005-0012-x
 28. Agele SO, Iremiren GO, Ojeniyi SO. Evapotranspiration, water use efficiency and yield of rainfed and irrigated tomato in the dry season in a humid rainforest zone of Nigeria. International Journal of Biology & Agricultural Sciences. 2011;13:469-476.
 29. Yang Q, Wang Y, Jia X, Zheng Y, He S, Deng L, Ma Y, Xie R, Yi S, Lv Q. Fruit yield and quality response of Newhall navel orange to different irrigation regimes and ground cover in Chongqing Three Gorges Reservoir area. Scientia Horticulturae. 2018;241(15):57-64. DOI:10.1016/j.scienta.2018.06.083
 30. Smajstrla AG, Locascio SJ, Weingartner DP, Hensel DR. Subsurface drip irrigation for water table control and potato production. Applied Engineering in Agriculture. 2000;16(3):225-229. DOI: 10.13031/2013.5147
 31. Bonachela S, Villalobos SJ, Fereres E. Soil evaporation from drip-irrigated olive orchards. Irrigation Science. 2001;20:65-71.

- DOI:<https://doi.org/10.1007/s002710000030>.
32. García-Tejera O, López-Bernal A, Orgaz F, Franciscol JI, Villalobos FJ. Analysing the combined effect of wetted area and irrigation volume on olive tree transpiration using a SPAC model with a multi-compartment soil solution. *Irrig Sci*; 2017. DOI:10.1007/s00271-017-0549-5
 33. Daymond AJ, Hadley P. The effects of temperature and light integral on early vegetative growth and chlorophyll fluorescence of four contrasting genotypes of cacao (*T. cacao*). *Annals of Applied Biology*. 2004;145:257-262.
 34. Daymond AJ, Hadley P, Machado RCR, Ng E. Canopy characteristics of contrasting clones of cocoa (*T. cacao* L.). *Hort Science*. 2002;37:799-801.
 35. Akinseye FM, Agele SO, Traore PCS, Adam M, Whitbread AM. Evaluation of the onset and length of growing season to define planting date—a case study for Mali (West Africa) *Theor Appl Climatol*; 2015. DOI 10.1007/s00704-015-1460-8
 36. Agele S, Famuwagun B, Ogunleye A. Effects of shade on microclimate, canopy characteristics and light integrals in dry season field-grown cocoa (*Theobroma cacao* L.) seedlings. *Journal of Horticulture*. 2016;11(1):47 – 56.
 37. Ertek A. Importance of pan evaporation for irrigation scheduling and proper use of crop-pan coefficient (Kcp), crop coefficient (Kc) and pan coefficient (Kp). *African Journal of Agricultural Research*. 2011;6(32):6706-6718. DOI: 10.5897/AJAR11.1522
 38. Intergovernmental Panel on Climate Change (IPCC). Special Report on global warming on 1.5 °C (SR 14); 2018. Available:<http://www.ipcc.ch>
 39. Locascio SJ, Smajstrla AG. Water application scheduling by pan evaporation for drip-irrigated tomato. *J. Amer. Soc. Hort*. 1996;121:63-68.
 40. López-López M, Espadafor M, Testi L, Lorite IJ, Orgaz F, Elías F. Water use of irrigated almond trees when subjected to water deficits. *Agricultural Water Management*. 2018; 195(C):84-93.
 41. Pereira LS, Allen RG, Smith M, Raes D. Crop evapotranspiration estimation with FAO 56: Past and future. *Agric. Water Manag*. 2015;147:4–20.
 42. Testi L, Villalobos FJ. New approach for measuring low sap velocities in trees. *Agric. For. Meteorol*. 2009;149(3-4):730-734.
 43. United Nations Framework Convention on Climate Change (UNFCCC). UNFCCC Process: Convention, Kyoto Protocol and the Paris Agreement; 1994. Available:<https://unfccc.int>

© 2023 Agele; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/96942>