



Influence of Different Spacings under Varying Fertility Levels on Protein Content, Protein Yield and Available NPK in Soil of Summer Greengram (*Vigna radiata* L.)

Priya R. Chaudhari ^{a++*}, K. I. Patel ^{b#},
Karuna M. Chaudhari ^{a++} and Yonika Saini ^{c++}

^a Department of Agronomy, NMCA, NAU, Navsari, Gujarat, India.

^b ARS, Kholwada, Siddhpur, SDAU, S. K. Nagar, Gujarat, India.

^c Department of Agronomy, Agricultural University, Kota, Rajasthan, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2024/v36i44473

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/113719>

Original Research Article

Received: 24/12/2023

Accepted: 29/02/2024

Published: 04/03/2024

ABSTRACT

The present study highlights the influence of different spacings under varying fertility levels on protein content, protein yield and available npk in soil of summer greengram (*vigna radiata* l.). Due to its importance in Indian agricultural and ability to provide vital amino acids and protein, food beans are a staple of the Indian diet. In the summer of 2020, a field experiment on loamy sand soil was conducted at the Agronomy Instructional Farm, Chimanbhai Patel College of Agriculture,

⁺⁺ Ph. D. Scholar;

[#] Associate Research Scientist;

^{*}Corresponding author: E-mail: priyachaudhari2908@gmail.com;

Sardarkrushinagar Dantiwada Agricultural University, Sardarkrushinagar. Twelve treatment combinations comprising of four spacings S₁ (30 × 10 cm), S₂ (30 × 20 cm), S₃ (45 × 10 cm) and S₄ (45 × 20 cm) and three fertility levels viz., F₁ (75% RDF), F₂ (100% RDF) and F₃ (125% RDF) were evaluated in factorial randomized block design with three replications. The results revealed that significantly higher protein yield (256.10 kg/ha) under 30 × 10 cm spacing as compared to other treatments. Among the fertility levels significantly higher protein content (24.17%), protein yield (260.27 kg/ha) and Higher value of available N and P₂O₅ status in soil after harvest was observed in treatment (F₃) 125% RDF of summer greengram. Thus, from the foregoing results of one year experiment, it can be concluded that for securing higher seed yield, stover yield, protein yield and higher available N, P₂O₅ in soil after harvest of summer greengram should be sown at 30 cm × 10 cm spacing and fertilized with 100% RDF (20-40-00 N-P₂O₅-K₂O kg/ha) under loamy sand soil.

Keywords: Greengram; protein content; yield; spacing; goldengram; fertilizers; organic carbon; foliar sprays.

1. INTRODUCTION

"Pulses are wonder present of nature to the living universe and are the real gateway of sustainable agriculture. Food legumes constitute an important dietary ingredient of Indian diet as they supply protein and essential amino acids and play significant role in Indian farming" [1]. "Pulses are the cheapest source of quality protein for the human beings" [2]. "The protein hunger is common problem in India, where majority of the people have vegetative diet. Greengram native is India and central Asia. Greengram is cultivated in the countries of India, Burma, Srilanka, Pakistan, China, Fiji, Queens land and Africa. Greengram (*Vignaradiata*) is commonly known as moog, goldengram, mung. It's belonging to family Leguminosae" [1]. "It is third important pulse crop after chickpea and pigeonpea, cultivated throughout India for its multipurpose uses as vegetable, pulse, fodder and green manure crop. Its seed is more palatable, nutritive, digestible and non-flatulent than other pulses grown in the world. Spacing plays an important role in contributing to the high yield because thick plant population will not get proper light for photosynthesis and high infestation of diseases" [3]. "On other hand very low plant population will also reduce the yield" [3]. "Due to this reason normal plant population is essential for high yield. Advantage of optimum spacing under irrigated conditions is due to reduced competition for light because when the moisture is lacking, light is no longer only limiting factor and advantage of uniform spacing is lost" [4]. "It is the most important non-monetary input, which can be manipulated to attain the maximum production per unit area" [5]. "Fertilizer plays important role for obtaining crop production" [6]. For greengram, 20 kg nitrogen (N), 40 kg phosphorus should be given at sowing time. It is advisable to use

fertilizers on the basis of soil test and recommendations, normally 100 kg DAP/ha is sufficient for one hectare the fertilizer should be applied by drilling either at the time of sowing or just before sowing in such way that they are placed about 2-3 cm below the seed.

Keeping facts in view the present study entitled "Influence of different spacings under varying fertility levels on protein content, protein yield and available npk in soil of summer greengram (*Vigna radiata* L.)" will be taken with the following objective:

1. To study the effect of different spacings and fertility levels on protein content and protein yield of summer greengram
2. To find the effect of different spacings and fertility levels on available nutrients in soil of summer greengram

2. METHODOLOGY

The field experiment was conducted at Agronomy Instructional Farm, Chimanbhai Patel College of Agriculture, Sardarkrushinagar Dantiwada Agricultural University, Sardarkrushinagar during summer season of year 2020. The experiment field was fairly leveled and uniform. "The soil of the experimental plot was loamy sand in texture and slightly alkaline having pH 7.4. The soil was low in organic carbon (0.20 %) and available nitrogen (137 kg/ha), medium in available phosphorus (31.24 kg/ha) and high in potassium (282 kg/ha) status based on the chemical analysis of soil. Twelve treatments were laid out in Randomized Block Design with factorial concept with three replications. The four levels of spacings (S₁-30 × 10 cm, S₂-30 × 20 cm, S₃- 45 × 10 cm and S₄-45 × 20 cm) and three levels of fertility levels

[F₁-75% (15-30-00 N-P₂O₅-K₂O kg/ha) RDF, F₂-100% (20-40-00 N-P₂O₅-K₂O kg/ha) RDF and F₃-125% (25-50-00 N-P₂O₅-K₂O kg/ha) RDF]. The greengram variety GM 6 was sown on 4th march 2020". [7]

3. RESULTS AND DISCUSSION

The results obtained from the present investigation as well as relevant discussion have been summarized under following heads.

3.1 Effect of Spacings on Quality Parameters and Nutrient Studies

3.1.1 Protein content and Protein yield

The results from Table 1 revealed that protein content different spacings found non-significant effect on protein content in seed of greengram.

However numerically higher protein content (23.33 %) was found in 45 × 20 cm spacing (S₄) followed by 30 × 20 cm spacing (S₂: 23.18 %). The lowest protein content (22.22 %) was noted with 30 cm × 10 cm spacing (S₁). The reason may be more competition between plants for growth resources like nutrient, moisture, light under narrow spacing that reduced protein content in seed. Similar results observed by Panchal [8], Patel [9], Patel [1] and Patel et al. [10] in mothbean and incase of protein yield in Table 1 showed that 30 × 10 cm (S₁) spacing gave significantly higher protein yield (256.10 kg/ha) but it was remained statistically at par with 45 × 10 cm (S₃) and 30 × 20 cm (S₂). This might be due to increased seed yield under 30 × 10 cm (S₁) spacing. The lowest protein yield (189.54 kg/ha) was recorded under 45 × 20 cm spacing (S₄). Similar results were observed by Panchal [8] and Patel [1].



Fig. 1. 45 cm × 20 cm + 125% RDF

Table 1. Effect of spacings and fertility levels on protein content in seed and protein yield of summer greengram

Treatments	Seed yield (kg/ha)	Protein content (%) in seed	Protein yield (kg/ha)
Spacings (S)			
S ₁ : 30cm × 10cm	1133	22.22	256.10
S ₂ : 30cm × 20cm	927	23.18	216.01
S ₃ : 45cm × 10cm	1037	22.33	231.86
S ₄ : 45cm × 20cm	812	23.33	189.54
S.Em. ±	107.04	1.22	22.45
C.D. (P=0.05)	314	NS	65.84
Fertility levels (F)			
F ₁ : 75% RDF	808	20.99	168.77
F ₂ : 100% RDF	1044	23.13	241.10
F ₃ : 125% RDF	1081	24.17	260.27
S.Em. ±	92.70	1.05	19.44
C.D. (P=0.05)	272	3.093	57.02
Interaction (S × F)			
S.Em. ±	61.80	0.70	12.96
C.D. (P=0.05)	NS	NS	NS
C.V.%	10.95	5.35	10.05



Fig. 2. 30 cm x 10 cm + 125 % RDF




Fig. 3. Effect of spacings and fertility levels on available N, P₂O₅, and K₂O content in soil after harvest of summer greengram

3.1.2 Available NPK in soil after harvest

The results graphically illustrated in Fig. 3 showed that available N, P₂O₅ and K₂O in soil after harvest did not differ significantly. Though marginally higher available N, P₂O₅ and K₂O (167.56, 41.19 and 261.90 kg/ha) was recorded under 45 x 20 cm (S₄) spacing. This might be due to less competition for nutrient in wider spacing which ultimately lead more nutrient left over in soil. The lowest available nitrogen of 161.19 kg/ha was observed under the spacing of 30 x 20 cm while available P₂O₅ and K₂O less under spacing 30 x 10 cm. Similar findings were reported by Patel [1] and Ezung et al. [11].

3.2 Effect of Fertility Levels on Quality Parameters and Nutrient Studies

3.2.1 Protein content and protein yield

A perusal of data presented in Table 1 indicated that application of 125% RDF (F₃) registered

significantly higher protein content (24.17 %) and protein yield (260.27 kg/ha) in seed of greengram but it was statistically at par with 100% RDF (F₂). Protein content in mungbean seed increased with an increase in the availability of N. This was mainly due to the structural role of N in building up amino acids. Meena et al. [12]. The lowest protein content (20.99 %) and protein yield of 168.77 kg/ha registered with application of 75% RDF (F₁). These results are also close conformity with the finding of Singh [13] and Meena et al. [12].

3.2.2 Available NPK in soil after harvest

The beneficial effect of varying fertility levels on available N and P₂O₅ in soil after harvest have been graphically presented in Fig. 3. Application of 125% RDF (F₃) resulted Significantly higher available N and P₂O₅ in soil 179.97 and 42.15 kg/ha) and was statistically at par with 100% RDF (F₂) Because higher rate of fertilizer application and nitrogen fixation by plants, while

non-significant effect of varying fertility levels on available K₂O in soil after harvest of greengram. Similar results were also observed by khavith et al. [14] in cowpea and Jat et al. [15] in greengram.

4. CONCLUSION

On the basis of finding of the present investigation, it can be concluded that significantly higher seed yield, stover yield, protein yield and higher available N, P₂O₅ in soil after harvest of summer greengram should be sown at 30 cm × 10 cm spacing and fertilized with 100% RDF (20-40-00 N-P₂O₅-K₂O kg/ha) under loamy sand soil.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Patel BR. Performance of summer greengram (*Vigna radiata*) cultivars to plant densities under north Gujarat conditions. Unpublished M. Sc. (Agri.) Thesis submitted to Sardarkrushinagar Dantiwada Agricultural University, Sardarkrushinagar (Gujarat); 2017.
- Ali S. Weed control in *kharif* greengram (*Vigna radiata* L. wilczek). Unpublished M.Sc. (Agri.) Thesis submitted to Sardarkrushinagar Dantiwada Agricultural University, Sardarkrushinagar (Gujarat); 2010.
- Patel PT. Response of summer greengram (*Vigna radiata* L.) varieties to row spacing and foliar sprays under south Gujarat conditions. Unpublished M.Sc. (Agri.) Thesis submitted to Navsari Agricultural University, Navsari (Gujarat);2020.
- Ihsanullah FH, Akbar H, Basir A, Ullah N. Effect of row spacing on agronomic traits and yield of mungbean (*Vigna radiata* L. Wilczek). Asian Journal of Plant Sciences. 2002;1(4):328-329.
- Jain VK, Chauhan YS. Performance of greengram cultivars under different row spacings. Indian Journal of Agronomy. 1988;33(3):300-302.
- Shelke AV. Varietal response to different levels of fertility and bio-fertilizers on yield and quality of summer greengram. Unpublished M.Sc (Agri) Thesis submitted to Anand Agricultural University, Anand (Gujarat); 2011.
- Chaudhari PR, Patel KI, Talaviya PL, Patel PD, Patel PM. Effect of different spacings under varying fertility levels on summer greengram (*Vigna radiata* L.). The Pharma Innovation Journal. 2023;12(2): 1027-1032.
- Panchal CC. Response of summer greengram (*Vigna radiata* L.) to different row spacing and weed management practices under south Gujarat conditions. Unpublished M.Sc. (Agri.) Thesis submitted to Navsari Agricultural University, Navsari (Gujarat); 2015.
- Patel DK. Influence of irrigation based on IW/CPE ratio and spacing on growth, yield attributes and yield of summer greengram (*Vigna radiata* L.) cv. Meha under middle Gujarat condition. Unpublished M.Sc (Agri) Thesis submitted to Anand Agricultural University, Anand (Gujarat); 2012.
- Patel BJ, Patel HH, Gavit VC. Effect of spacing and weed management on yield, quality and nutrient uptake of moth bean (*Vigna aconitifolia*). Journal of Pharmacognosy and Phytochemistry. 2019;8(5):948-952.
- Ezung NK, Yanthan M, Rajkhowa DJ, Jamir T. Performance of greengram as influenced by different levels of vermicompost + rhizobium and spacing on growth, yield and economics under upland terrace condition of north east India. Research Journal of Agricultural Sciences. 2020;11(2):349-354.
- Meena RS, Verma T, Verma S, Singh A, Kumar S, Gurjar DS. Influence of organic and inorganic sources of nutrients on growth, yield and quality of mungbean (*Vigna radiata*). Indian Journal of Agricultural Sciences. 2020; 90(11): 2233-2236.
- Singh. Effect of organic and inorganic fertilizers on growth, yield and quality of different varieties of soybean (*Glycine max*); 2012. Accessed 10 August 2015. Available:<https://www.researchgate.net/publication/280829958>
- Kavitha MP, Balakumbahan R, Prabukumar G. Effect of foliar spray and fertilizer levels on growth and yield of vegetable cowpea [*Vigna unguiculata* (L.) Walp.]. Indian Journal

- Agricultural Research. 2019; 53(6):745-748.
15. Jat RA, Arvadia MK, Tndel B, Patel TU, Mehta RS. Response of saline water irrigated greengram (*Vigna radiata* L.) to land configuration, fertilizer and farm yard manure in Tapi command area of south Gujarat. Indian Journal of Agronomy. 2012; 57(3): 270-274.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/113719>