



Effect of FYM and Inorganic Fertilizers on Growth, Yield of Wheat (*Triticum aestivum* L.) and Soil Properties of Red Alfisol

Seema Hodkashia ^{a,b}, Parmod Kumar Sharma ^a,
Ashish Marotrao Latore ^a, Saroj Choudhary ^c, Bharti Yadav ^a
and Sarvendra Kumar ^{b*}

^a Department of Soil Science and Agricultural Chemistry, Institute of Agricultural Science, Banaras Hindu University, Varanasi (Uttar Pradesh), India.

^b Division of Soil Science and Agricultural Chemistry, ICAR-Indian Agricultural Research Institute, New Delhi-110012, India.

^c ICAR- Agricultural Technology Application Research Institute, Jodhpur (Rajasthan), India.

Authors' contributions

This work was carried out in collaboration among all authors. Author SH performed methodology and did the analysis of soil and plant samples. Authors PKS and AML supervised the study. Author AML did design of the experiment, monitoring of experiments and laboratory analysis. Authors SC, SH and BY analysis of data. Authors SK and PKS reviewed and edited the manuscript. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajsspn/2024/v10i4415>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/124681>

Original Research Article

Received: 16/08/2024

Accepted: 18/10/2024

Published: 12/11/2024

*Corresponding author: Email: sarv.grandee@gmail.com;

Cite as: Hodkashia, Seema, Parmod Kumar Sharma, Ashish Marotrao Latore, Saroj Choudhary, Bharti Yadav, and Sarvendra Kumar. 2024. "Effect of FYM and Inorganic Fertilizers on Growth, Yield of Wheat (*Triticum Aestivum* L.) and Soil Properties of Red Alfisol". *Asian Journal of Soil Science and Plant Nutrition* 10 (4):403-13. <https://doi.org/10.9734/ajsspn/2024/v10i4415>.

ABSTRACT

The soil multi-nutrient deficiency is not only a problem for soil quality, but also for crop productivity. In problematic soils, in particular acid soils, where low nutrient availability is a concern for general soil use, organic fertilizers are important. The importance of FYM is not limited to their role as accessibility, cost-effectiveness, soil nutrient reservoirs, moisture and ameliorating soil properties that determine soil fertility and productivity status. A pot experiment was conducted on wheat crop at Banaras Hindu University, India, to investigate the effect of FYM and inorganic fertilizers on growth, yield of wheat (*Triticum aestivum* L.) and soil properties of red Alfisol. The results revealed that maximum number of tiller pot⁻¹ (17.0), plant height (78.0 cm), grain yield (27.4 g pot⁻¹), straw yield (9.98 g pot⁻¹), biological yield (67.5 g pot⁻¹), harvest index (40.6%), were recorded in 1.5RF+FYM1 applied pots. The correlation study revealed a significant positive correlation of soil organic carbon with biological yield, and soil available N, P and K. The integration of FYM along with 1.5 times RF significantly enhanced wheat productivity, quality, and soil nutrient availability, indicating the potential of this integrated approach for restoring soil fertility in degraded soils to sustainable crop production in red Alfisol.

Keywords: *Integrated nutrient management; crop productivity; soil fertility; degraded land; super optimal dose.*

1. INTRODUCTION

Red soils (Alfisol in the US soil taxonomy system), are particularly prevalent in India and cover approximately 70 million hectares or 28% of the total geographical area [1]. With the high temperature and rainfall, these regions are important for agriculture. However, because of high weathering, poorly developed soil structure and some unsuitable management practices, produce significant multi-nutritional deficiency including both primary (Nitrogen, phosphorous and potassium), and secondary nutrients (calcium, magnesium and sulfur) [2]. Therefore, effective reclamation strategies are required for these soils such as incorporation of organic matter, irrigation management and the use of appropriate soil amendments to improve soil structure and fertility. As such, inorganic fertilizers play an important role in crop production and the growth of the agricultural economy [3]. The unsatisfied effect of the sole application of chemical fertilizer on soil fertility and crop productivity under degraded land, need for better planning for resource and nutrient management besides intensification of cropping. The use of chemical fertilizers increases in an unbalanced manner and has produced problems of multi-nutrient deficiency, mainly micronutrients, reducing soil fertility and unsustainable crop productivity [4]. The application of inorganic fertilizers with organic manures seems to be a good solution for these problems [5].

Wheat (*Triticum aestivum* L.) is the most important cereal crop grown worldwide and a staple of about 2.5 billion of the world population

[6]. The nutrient-rich cereal is grown in various environmental conditions; globally wheat established around 217 million hectares holding the position of highest acreage among all other crops with annual production up to 731 million tonnes [7]. It requires a high amount of nitrogen (N) fertilizers to enable grain protein production which is necessary for backing and processing quality [8]. The recommended dose of fertilizers was not able to maintain soil fertility, productivity and quality of grain under degraded soil conditions, thus we used integrated nutrient management with various rates of NPK doses. Hence the use of fertilizers either alone or in combination with organic fertilizers is important for sustaining soil fertility and crop productivity [9] because organic manures supply a good amount of plant nutrients [10].

The organic sources might include green manure, organic manure, crop residues, rural wastes, vermicompost and biofertilizers [11]. Among many kinds of organic sources, farmyard manure (FYM) is a commonly used organic fertilizer in developing countries because it is easily available and cheap in rate (Kei and Palanichamy, 2013). The use of well-decomposed farmyard manure (FYM) in soil management practices is a good practice for increasing crop yield, improving SOM, enhancing microbial activity, provide eco-friendly soil environmental management [12-14] and increasing plant available macro and micronutrients in soil [15]. Moreover, organic fertilizers act as slow-release fertilizers and provide nutrients over a long period [16]. However, to date, little attention has focused on

how the FYM with various NPK rates influences the management of soil fertility.

The utilization of inorganic nutrient sources with organic is a feasible approach for higher crop productivity and maintaining soil health [17,18]. FYM is a good nutrient source that has the potential to positively affect soil physical, chemical and biological properties [19-21] and thus, help in improve nutrient availability and crop yield [22]. Integrated nutrient management (INM) are essential for better plant growth, water utilization, soil and land management which is more important for sustaining agriculture productivity [23,24]. Schoebitz et al. [25] revealed that INM increases applied nutrient use efficiency while reducing nutrient losses. A better soil physical environment helped in better uptake of water and nutrients due to nutrient sufficiency in FYM and inorganic fertilizer application together [26,27]. Therefore, the purpose of this study was to compare the various doses of inorganic fertilizer management with or without FYM to improve low nutrient availability, soil properties, crop growth and productivity in degraded red alfisol using a pot experiment.

2. MATERIALS AND METHODS

2.1 Experimental Site and Experimental Methods

The pot experiment of wheat crop was conducted at the Rajiv Gandhi South Campus, Banaras

Hindu University, Barkachha, Mirzapur (at 25.07° N latitude and 82.59° E longitudes and altitude of 427 meters above mean sea level) during the rabi season of 2019-2020. The experimental soil was collected from an agricultural research farm, Barkachha, Mirzapur. The soil was air-dried, gently powdered to pass through a 2 mm sieve and homogenized and placed in an earthen pot. The pots had a diameter of 15 inches and height 17 inches. We used twenty-seven pots with 10 kg of prepared soil per pot. The treatment comprised a combination of FYM (0, 5, 10 t ha⁻¹) and inorganic fertilizers i.e. NPK (0%, RF, 1.5RF and soil test based) (Table 2). There were total nine treatments, replicated thrice in a completely randomized design viz., T₁: control, T₂: RF, T₃: 1.5 RF, T₄: FYM1 (5 t ha⁻¹), T₅: FYM2 (10 t ha⁻¹), T₆: RF+FYM1 (5 t ha⁻¹), T₇: 1.5RF+FYM1 (5 t ha⁻¹), T₈: Soil test based fertilizer, T₉: Soil test based fertilizer + FYM1 (5 t ha⁻¹). According to the recommended doses of fertilizers (RF) for wheat, (120 N: 60 P₂ O₅: 60 K₂ O kg ha⁻¹), we calculated doses of N, P, and K for each of the experimental treatments (Table 2).

The dose of nitrogen(N) and full dose of phosphorous(P) and potassium (K) were applied as basal dose at the time of sowing. The remaining dose of N was top dressed in two equal splits at 30 and 60 days after sowing of wheat. The source of N, P and K were urea, di ammonium phosphate (DAP) and murate of potash (MOP), respectively. After application of NPK fertilizers and FYM, the soil in each pot was

Table1. Physio-chemical properties of soil and farm yard manure used in the experiment

Properties	Soil	FYM
Soil texture	Sandy loam	-
pH	6.6	6.56
EC (dSm ⁻¹)	0.204	3.52
Organic Carbon (%)	0.36	9.82
Available nitrogen	132(kg ha ⁻¹)	0.46%
Available phosphours	7.69 (kg ha ⁻¹)	0.23%
Available potassium	168 (kg ha ⁻¹)	0.44%

Table 2. Description Treatment detail

Treatments	Treatments detail	
	Per ha	Per pot
T ₁ -CK	-	-
T ₂ -RF	120 kg N+ 60kg P ₂ O ₅ +60 kg K ₂ O ha ⁻¹	0.53 g N + 0.26 g P ₂ O ₅ + 0.26 g K ₂ O pot ⁻¹
T ₃ - 1.5 RF	180 kg N + 90 kg P ₂ O ₅ + 90 kg K ₂ O ha ⁻¹	0.79 g N+ 0.39g P ₂ O ₅ + 0.39g K ₂ O pot ⁻¹
T ₄ - FYM1 (5t ha ⁻¹)	23 kg N+11.5 kg P ₂ O ₅ + 22 kg K ₂ O ha ⁻¹	0.102 g N + 0.051 g P ₂ O ₅ + 0.098 g K ₂ O pot ⁻¹
T ₅ - FYM2 (10 t ha ⁻¹)	46 kg N + 23 kg P ₂ O ₅ + 44 kg K ₂ O ha ⁻¹	0.204 g N + 0.102 g P ₂ O ₅ + 0.196 g K ₂ O pot ⁻¹
T ₆ -RF+FYM1	143 kg N + 71.5 kg P ₂ O ₅ + 82 kg K ₂ O ha ⁻¹	0.632 g N + 0.311 g P ₂ O ₅ + 0.358 g K ₂ O pot ⁻¹
T ₇ -1.5 RF+FYM1	203 kg N + 101.5 kg P ₂ O ₅ + 112 kg K ₂ O ha ⁻¹	0.892 g N+ 0.441 g P ₂ O ₅ + 0.488 g K ₂ O pot ⁻¹
T ₈ -Soil test-based (STF)	150 kg N + 75 kg P ₂ O ₅ + 40 kg K ₂ O ha ⁻¹	0.66 g N + 0.33 g P ₂ O ₅ + 0.17 g K ₂ O pot ⁻¹
T ₉ -STF + FYM1	173 kg N + 86.5 kg P ₂ O ₅ + 62 kg K ₂ O ha ⁻¹	0.762 g N+ 0.381 g P ₂ O ₅ + 0.268 g K ₂ O pot ⁻¹

CK- Control (unfertilized); RF- recommended dose of inorganic NPK by inorganic fertilizers; FYM-farm yard manure

thoroughly remixed and equilibrated for 1 week before sowing. The seeds of wheat (var. Malviya234) were sown in mid-november during the year as a test crop. Ten seeds of wheat were sown in the mixture filled pots. Seven days after seedling emergence the seedlings were thinned to 5 per pot. The moisture contain in all pots was maintained and irrigated with tap water. During the experiment, the maximum mean air temperature was 34.7°C and the minimum 16°C. The growth parameters of wheat plant viz. plant height and number of tillers were recorded on each of the three replicate pots. The plant heights were measured at 30,60,90 days after sowing and harvest of the crop. Plant height was recorded by a 150 cm long scale with 0.5 cm interval markings and from the base level to the tip of the middle trifoliate leaf. The tiller number per plant was recorded at 45 and 90 days after the sowing of wheat. The number of tillers per plant was counted manually. The crop was harvested at physiological maturity, washed with dilute HCl, soap solution and deionized water. Soil samples were also collected for chemical analysis. Soil samples were packed in plastic bags and labelled accordingly. The soil properties like pH and electrical conductivity (EC), organic carbon (OC), available nitrogen, phosphorous and potassium were analyzed. Laboratory procedures were followed to determine various soil and FYM properties. The pH and electrical conductivity (EC) were determined in soil {soil to water ratio of 1:2.5 (weight/volume)} and FYM {FYM to water ratio of 1:5} by using pH meter and EC meter, respectively [28,29]. The organic carbon content in soil was determined using the wet digestion-based Walkley-black method (1934). Available nitrogen in soil was measured by applying the Alkaline potassium permanganate method (0.32% KMnO_4) (Subbiah and Asija, 1956) [30]. The available phosphorous (P_2O_5) in soil was measured by extracting 0.5 M NaHCO_3 with the help of an ascorbic acid method using calorimetry at 730nm (Olsen and Dean. 1965) [31]. Soil-available potassium was extracted with neutral ammonium acetate (pH, 7.0) and then determined with the help of flame photometrically [28,29]. General properties of soil and FYM prior to the experiment are provided in Table 1.

2.2 Statistical Analysis

Statistical analysis was used to establish the significance between the treatment means and draw valid results. Analysis of variance (ANOVA) method was adopted as the statistical analysis

tool for the raw data recorded during the whole experiment. The difference of the treatment means was evaluated using the least significant difference (LSD) at 5% level of probability as per Gomez and Gomez [32] following the complete Randomized Design (CRD). Statistical analysis of data was carried out with the help of SPSS statistical package (version 23.0, SPSS Inc.) for windows.

3. RESULTS AND DISCUSSION

3.1 Effect of FYM with Inorganic Fertilizer on Growth Indices of Wheat

The plant height of wheat had an increasing trend from 30 days after sowing to harvest of the crop, and significant changes among the treatments were also noticed (Table 3). Plant height ranged from 7.0 to 18.2 cm at 30 DAS, 19.3 to 35.6 cm at 60 DAS, 49.6 to 65.8 cm at 90 DAS, and 59.6 to 78.0 cm at harvest of wheat. The highest plant height of wheat was recorded in treatment T_7 (1.5 NPK + FYM1) followed by T_3 (1.5 NPK) and minimum was in treatment T_1 irrespective of growth stage of crop. Secondly, the combined application of inorganic fertilizers with FYM favoured the growth of plant and recorded higher plant height than only inorganic fertilizers applied plants. The plants grown in treatment T_7 and T_3 had 159% and 143% at higher plant height at 30 DAS, respectively. However, the intensity of increment of plant height was decreased 31% and 28% at harvest over control, respectively. It might be due to the N uptake at the initial stage of growth being more. Plant height increased availability of nutrients in soil due to higher fertilizer application may have increased meristematic activity (multiplication and elongation of cells) leading to enhanced plant height [29].

The range of tiller number was 9.0 to 17.0 (Table 3). It was revealed that application of 1.5 NPK+FYM1 (T_7) recorded significantly the highest numbers of tiller plant⁻¹ over the rest of treatments followed by 1.5RF (T_3) and minimum number tiller was observed in T_1 (9.0). Treatment T_6 was found at par with respect to T_8 . The plant growth had 88.8% and 84.4% higher numbers of tillers in T_7 and T_3 over control, respectively. It may be due to FYM with NPK application help in improving physical properties of the soil, enhancing soil fertility and availability of many nutrients elements to plant uptake which increases plant growth. Apriyani et al., [30] observed that increasing supply of NPK and

Table 3. Effect of FYM and inorganic fertilizer on plant growth rate and tiller numbers

Treatment	Plant height				Tiller pot ⁻¹
	30DAS	60DAS	90DAS	At harvest	
T ₁	7.02	19.3	49.6	59.6	9.0
T ₂	14.2	30.1	58.1	68.5	12.6
T ₃	17.0	34.0	65.3	76.0	16.6
T ₄	14.1	22.3	56.0	66.9	9.3
T ₅	14.2	28.1	56.5	67.8	10.0
T ₆	15.0	31.0	60.1	70.0	13.6
T ₇	18.2	35.6	65.8	78.0	17.0
T ₈	16.0	32.1	64.2	73.9	13.6
T ₉	16.4	33.5	65.0	75.0	14.3
LSD (P = 0.05)	1.0	1.8	2.8	2.0	1.0

CK- Control (unfertilized); RF- recommended dose of inorganic NPK by inorganic fertilizers; FYM-farm yard manure
 {T₁: CK, T₂: RF, T₃: 1.5 RF, T₄: FYM1 (5t ha⁻¹), T₅: FYM2 (10t ha⁻¹), T₆: RF+FYM1, T₇: 1.5RF+FYM1, T₈: Soil test based fertilizer(STF), T₉: STF+ FYM (5t ha⁻¹), LSD = Least significant difference}.

micronutrients from the FYM offers more balanced nutrition for plants. Inorganic fertilizers improve the quick availability of nutrients compared to the gradual release of nutrients from organic fertilizer sources [31].

3.2 Effect of FYM with Inorganic Fertilizer on Growth Yield Attributes and Yield of Wheat

The crop under integrated fertilization recorded a significantly higher yield of wheat compared to inorganic fertilizers alone. The grain yield of wheat observed range from 10.5 to 27.4 g pot⁻¹. The maximum grain yield of wheat was in 1.5NPK+FYM1 (27.4 g pot⁻¹) followed by 1.5 RF (25.6 g pot⁻¹). The lowest grain yield was recorded in control (T₁) (10.5 g pot⁻¹). It might be due to integrated fertilizer application increasing soil quality and availability of nutrients to sustain greater yield compared to inorganic fertilizer alone. The grain yield of wheat was also increased with the rate of FYM application increased. It was recorded that 12.5 g pot⁻¹ to 15.2 g pot⁻¹ with application of 5t ha⁻¹ (FYM1) to 10 t ha⁻¹ (FYM2), respectively. Organic manure increases total soil porosity, soil water and nutrient holding capacity and concentration of essential plant nutrients (macro and micro) [33] in soil ultimately improving wheat grain yield. Sharma and Biswas. (2017) revealed that microbiological activity increases with the application of organic manures, providing better nutrient mobilization from applied nutrient manures to higher root density that helps in adequate absorption of water and nutrients from the soil.

The straw yield of wheat ranged from 15.7 to 39.9 (g pot⁻¹). The highest straw yield (39.9 g pot⁻¹) was revealed in 1.5NPK+FYM1 (T₇) followed by 1.5NPK (T₃) (37.5 g pot⁻¹). It was significantly

higher over rest of the treatments. The straw yield was 6.35% higher in T₇ over T₃. Adding NPK with organic fertilizers supplied not only additional quantities of NPK directly, but also additional secondary and micronutrients that were limiting in the soil [34]. According to Meena et al. [35,36] increased uptake of N in leaf, stem and grain in NPK +FYM incorporated treatment might be associated with the mineralization of FYM throughout the growing season that ensured its availability to wheat crop and help to increase crop yield.

The biological yield (g pot⁻¹) of wheat significantly increased as the application of NPK with or without FYM increased. The highest biological yield was observed with the application of 1.5NPK+FYM1 (T₇) (67.5 g pot⁻¹) followed by 1.5 RF (63.2 g pot⁻¹) (Table 4). The minimum yield was recorded in T₁ (26.2 g pot⁻¹). It was revealed that 31.2 to 37.8 g pot⁻¹ in T₄ and T₅, respectively. It might be due to the FYM application increasing nutrient availability that helps in the improvement of yield. Increased availability of nutrients would have helped in the synthesis and translocation of carbohydrates from source to sink (grain) resulting in higher yield production [37]. Harvest index of wheat varied from 40.6 to 40.0% (Table 4). The highest harvest index was recorded with the application of 1.5NPK+FYM1 (T₇) among all treatments and minimum (40.0%) with control (T₁). The integrated application of fertilizer increases the harvest index more compared to the sole application. The same results were also observed by Ahmad et al., [38], Kumar et al., [39] [40].

3.3 Effect of FYM with Inorganic Fertilizer Soil Properties and Nutrients Status

Soil pH generally decreased after the addition of manure and inorganic fertilizers. The highest soil

pH was observed in CK (T₁) over the remaining treatments. same results were also found by Sharma et al. [41]. However, soil pH and EC both were non- significantly affected with the application of various treatments. As expected, integrated nutrient management significantly improve soil organic carbon values (Table 5). We observed maximum soil organic carbon value in 1.5RF+FYM1 (T₇) treatment followed by 1.5 RF (T₃). The enhancement of organic carbon in T₇ was 11.11% with respect to CK (T₁). This might be due to the direct incorporation of organic matter, and better root-growth of plants (Krishnamurthy et al. 2020) [42,43].

There was a significant difference among treatments with respect to the available N content of soil. The available N content of soils under various treatments ranged from 116 kg ha⁻¹ to 171 kg ha⁻¹(Table 5). The maximum available N was recorded with the application of 1.5RF+FYM1 (T₇) (171 kg ha⁻¹) followed by 1.5RF (T₃) (168 kg ha⁻¹) and minimum in CK (T₁) (116 kg ha⁻¹). It was 2.04% higher in T₇ as compared to 1.5NPK (T₃). It might be due to an additional supply of N through chemical fertilizers and mineralization of N from the

decomposition of FYM. Sharma et al., (2014) also reported greater available N where FYM applied along with 100%NP than in 100%NPK alone. The beneficial soil conditions like organic carbon, porosity, water holding capacity and greater multiplication of soil microbes help in the improvement of soil available nitrogen [44,45].

The available phosphorous ranged from 7.5 kg ha⁻¹ to 12.1 kg ha⁻¹ (Table 5). The maximum amount of available P was significantly higher in 1.5RF+FYM1 (T₇) over the rest of the treatments followed by 1.5RF (T₃). Available P level in RF +FYM1 (T₆) was found at par with respect to soil test based (T₈). The enhancement in available phosphorous with the application of NPK fertilizers in combination with organic manure might be due to organic acids secretion during decomposition which, improves the release of native phosphorous in soil [44]. Sharma et al. [46] found that FYM reduces P fixation by the inactivation of iron, aluminium, and hydroxyl-aluminium ions. Besides, the addition of FYM in soil increased available P in soil by mineralization or solubilization of native P reserves [47].

Table 4. Effect of FYM and inorganic fertilizer on yield of wheat crop

Treatment	Grain yield (g pot ⁻¹)	Stover yield (g pot ⁻¹)	Biological yield (g pot ⁻¹)	Harvest index(%)
T ₁	10.5	15.7	26.2	40.0
T ₂	21.4	31.8	53.3	40.2
T ₃	25.6	37.5	63.2	40.5
T ₄	12.5	18.7	31.2	40.0
T ₅	15.2	22.6	37.8	40.1
T ₆	22.4	33.1	55.5	40.3
T ₇	27.4	39.9	67.5	40.6
T ₈	23.5	34.6	58.1	40.3
T ₉	24.9	36.5	61.5	40.5
LSD (P = 0.05)	1.6	2.7	3.5	NS

CK- Control (unfertilized); RF- recommended dose of inorganic NPK by inorganic fertilizers; FYM-farm yard manure {T₁: CK, T₂: RF, T₃: 1.5 RF, T₄: FYM1 (5t ha⁻¹), T₅: FYM2 (10t ha⁻¹), T₆: RF+FYM1, T₇: 1.5RF+FYM1, T₈: Soil test based fertilizer(STF), T₉: STF+ FYM (5t ha⁻¹), LSD = Least significant difference}.

Table 5. Effect of FYM and inorganic fertilizer on soil pH, EC, SOC and available nitrogen, phosphorous and potassium.

Treatment	Soil chemical properties			Nitrogen	Phosphorous (kg ha ⁻¹)	Potassium
	pH	EC (dSm ⁻¹)	SOC (g kg ⁻¹)			
T ₁	5.91	0.10	3.60	116.5	7.5	160.5
T ₂	5.90	0.10	3.67	142.1	9.5	182.9
T ₃	5.85	0.11	3.90	168.0	11.2	205.4
T ₄	5.78	0.07	3.63	133.8	8.0	171.7
T ₅	5.73	0.06	3.65	137.9	8.2	175.4
T ₆	5.82	0.06	3.68	158.8	10.3	195.0
T ₇	5.81	0.10	4.0	171.4	12.1	212.8
T ₈	5.87	0.10	3.70	161.0	10.6	197.8
T ₉	5.85	0.08	3.80	163.0	10.8	198.8
LSD (P = 0.05)	NS	NS	0.17	12.1	1.5	17.2

{T₁: CK, T₂: RF, T₃: 1.5 RF, T₄: FYM1 (5t ha⁻¹), T₅: FYM2 (10t ha⁻¹), T₆: RF+FYM1, T₇: 1.5RF+FYM1, T₈: Soil test based fertilizer (STF), T₉: STF+ FYM (5t ha⁻¹), LSD = Least significant difference, SEM± = Standard error of mean}.

Table 6. Correlation between soil and crop characteristics

	Tiller pot-1	Biological yield (g pot-1)	pH	EC	OC	N	P	K
Tiller pot-1	1							
Biological yield (g pot-1)	0.96***	1						
pH	0.13	0.12	1					
EC	0.44	0.34	0.71*	1				
OC	0.85**	0.74*	0.006	0.51	1			
N	0.94***	0.95***	-0.081	0.19	0.76*	1		
P	0.98***	0.97***	0.101	0.37	0.82**	0.96***	1	
K	0.97***	0.97***	-0.01	0.29	0.81**	0.99***	0.99***	1

***Correlation is significant at 0.001 level

**Correlation is significant at 0.01 level

*Correlation is significant at 0.05 level

Study revealed that available potassium ranged from 160 to 212 kg ha⁻¹ (Table 5). The higher content of available K in 1.5RF+FYM1 (T₇) followed 1.5 RF and minimum in CK (T₁). The K content was observed 3.60%, 7.55%, 16.33% and 32.56% higher in T₇ over 1.5RF (T₃), soil test based(T₈), RF (T₂) and control (T₁), respectively. It might be due to an additional supply of K through the decomposition of FYM. Sharma and Walia (2018) [32,48] was found mineralization of organic fertilizers and solubilization from native sources during their decomposition increases available K content.

3.4 Correlation Study

Results on simple correlation coefficient revealed that most of the soil and plant parameters significantly ($p < 0.001$) correlated with each other as depicted (Table 6). Some typing error was here. Organic carbon was positively correlated with tiller number ($r = 0.85$), total N uptake ($r = 0.83$), available phosphorous ($r = 0.82$), and available potassium ($r = 0.81$) and these parameters were found non-significant at $p < 0.05$ but significant at $p < 0.01$. Available N ($r = 0.76$) and total K uptake ($r = 0.77$) were found significantly correlated to soil organic carbon at $p < 0.05$ indicating that the availability of N and total K uptake increased as organic carbon content increased in soil. A non-significant positive correlation was also observed between wheat biological yield and total uptake of N ($r = 0.98$), P ($r = 0.90$) and K ($r = 0.99$) by the crop. On the other hand, pH showed a negative correlation between protein content, available N and available K, indicating decreased availability with increased pH. The protein content significantly correlated with biological yield ($r = 0.89$, $p < 0.01$). This indicated that FYM is an important source of maintaining soil fertility, nutrient availability and crop productivity. Katkar et al. [49] observed the same positive correlation

after applying organic fertilizer alone or with mineral fertilizers [50-55].

4. CONCLUSION

The result from the present study indicated that the addition of FYM with superimposed inorganic fertilizers had a maximum restoration effect on soil properties and crop productivity, nutrient uptake, quality of grains over inorganic fertilizers alone, organic fertilizers alone and control treatments. The highest wheat production and soil quality parameters were observed in 1.5RF+FYM1 treated pots. This justifies that the application of integrated nutrient management with a super optimal dose can improve the soil's nutritional status to get good plant growth, higher yields and better produce quality in degraded land. The addition of FYM is one of the best options for sustaining soil health and crop productivity with an affordable cost by reclamation of alfisol, which can significantly improve soil structure and fertility.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI tools were used for data analysis, or writing materials during the editing of this manuscript.

ACKNOWLEDGEMENT

Seema Hodkashia is highly grateful to the Institute of Agricultural Science, Rajiv Gandhi South Campus, BHU, India for providing facilities to carry out this research work. The authors are thankful to Ms. Thaiseen Tabasum for her technical support during the experiment.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Singh R, Dass A, Singh VK. Sustaining crop production in rainfed areas in India: issues and strategies. *Current Journal of Applied Science and Technology*. 2020;39(25):154-173.
2. Zhang H, Gan Y, Huang G, Zhao W, Li F. Postharvest residual soil nutrients and yield of spring wheat under water deficit in arid northwest China. *Agricultural Water Management*. 2009;96(6):1045-1051.
3. Wang J, Cramer GL, Wailes EJ. Production efficiency of Chinese agriculture: Evidence from rural household survey data. *Agricultural economics*. 1996;15(1):17-28.
4. Moharana PC, Sharma BM, Biswas DR. Changes in the soil properties and availability of micronutrients after six-year application of organic and chemical fertilizers using STCR-based targeted yield equations under pearl millet-wheat cropping system. *Journal of Plant Nutrition*. 2017;40:165-176.
DOI: 10.1080/01904167.2016.1201504
5. Abbas G, Khatkhat JZK, Mir A, Ishaque M, Hussain M, Wahedi HM, Ullah A. Effect of organic manures with recommended dose of NPK on the performance of wheat (*Triticum aestivum L.*). *J. Anim. Plant. Sci*. 2012;22:683-687.
6. Ramadas S, Kumar TK, Singh GP. Wheat production in India: Trends and prospects. In *Recent advances in grain crops research*. Intech Open; 2019.
DOI: 10.5772/intechopen.86341
7. USDA-FAS. United States Department of Agriculture/Foreign Agricultural Service. Accessed July 1; 2018.
Available: <https://www.fas.usda.gov/commodities/cotton>
8. Zörb C, Ludewig U, Hawkesford MJ. Perspective on wheat yield and quality with reduced nitrogen supply. *Trends in Plant Science*. 2018;23:1029-1037.
DOI: 10.1016/j.tplants.2018.08.012
9. Shambhavi S, Kumar R, Sharma SP, Verma G, Sharma RP, Sharma SK. Long-term effect of inorganic fertilizers and amendments on productivity and root dynamics under maize-wheat intensive cropping in an acid Alfisol. *Journal of Applied and Natural Science*. 2017;9:2004-2012.
DOI: 10.31018/jans.v9i4.1480
10. Subehia SK, Sepehya S, Rana SS, Negi SC, Sharma SK. Long-term effect of organic and inorganic fertilizers on rice (*Oryza sativa L.*)–wheat (*Triticum aestivum L.*) yield, and chemical properties of an acidic soil in the western Himalayas. *Experimental Agriculture*. 2013;49:382-394.
DOI: 10.1017/S0014479713000173
11. Kant S, Sharma PK, Kumar V, Kumar A. Chelating compounds influence the chemical properties of post-harvest chromium contaminated soil after maize and mustard. *International Journal of Chemical Studies*. 2018;6:1672–80.
12. Blair N, Faulkner RD, Till AR, Poulton PR. Long-term management impacts on soil C, N and physical fertility: Part I: Broad balk experiment. *Soil and Tillage Research*. 2006;91:30-38.
DOI: 10.1016/j.still.2005.11.002
13. Kundu S, Bhattacharyya R, Prakash V, Ghosh BN, Gupta HS. Carbon sequestration and relationship between carbon addition and storage under rainfed soybean–wheat rotation in a sandy loam soil of the Indian Himalayas. *Soil and Tillage Research*. 2007;92:87-95.
DOI: 10.1016/j.still.2006.01.009
14. Rathod DD, Rathod PH, Patel KP, Patel KC. Integrated use of organic and inorganic inputs in wheat-fodder maize cropping sequence to improve crop yields and soil properties. *Archives of Agronomy and Soil Science*. 2013;59:1439-1455.
DOI: 10.1080/03650340.2012.724171
15. Sandhu PS, Walia SS, Gill RS, Dheri GS. Thirty-one years study of integrated nutrient management on Physico-chemical properties of soil under rice–wheat cropping system. *Communications in Soil Science and Plant Analysis*. 2020;51:1641-1657.
DOI: 10.1080/00103624.2020.1791156
16. Shaji H, Chandran V, Mathew L. Organic fertilizers as a route to controlled release of nutrients. In *Controlled release fertilizers for sustainable agriculture*. Academic Press. 2021;231-245.
DOI: 10.1016/B978-0-12-819555-0.00013-3
17. Kumar B, Dhar S, Paul S, Paramesh V, Dass A, Upadhyay PK, Abdelbacki AM. Microbial biomass carbon, activity of soil enzymes, nutrient availability, root growth, and total biomass production in wheat cultivars under variable irrigation and

- nutrient management. *Agronomy*. 2021;11: 669.
DOI: 10.3390/agronomy11040669
18. Kakraliya SK, Jat RD, Kumar S, Choudhary KK, Prakash J, Singh LK. Integrated nutrient management for improving, fertilizer use efficiency, soil biodiversity and productivity of wheat in irrigated rice wheat cropping system in Indo-Gangatic plains of India; 2017.
DOI: 10.20546/ijcmas.2017.603.017
 19. Chandra P, Gill SC, Prajapat K, Barman A, Chhokar RS, Tripathi SC, Singh GP. Response of wheat cultivars to organic and inorganic nutrition: Effect on the yield and soil biological properties. *Sustainability*. 2022;14:15-9578.
DOI: 10.3390/su14159578
 20. Wolie AW, Admassu MA. Effects of integrated nutrient management on rice (*Oryza sativa* L) yield and yield Attributes, nutrient uptake and some Physico-chemical properties of soil: A review. *Journal of Biology, Agriculture and Healthcare*. 2016;6:20-26.
 21. Khayat M. Evaluation Effect of Farmyard Manure (FYM) to Improve Cereal Crop Yield. *Journal of Crop Nutrition Science*. 2021;7:59-67.
 22. Islam MR, Sikder S, Bahadur MM, Hafiz MHR. Effect of different fertilizer management on soil properties and yield of fine rice cultivar. *Journal of Environmental Science and Natural Resources*. 2012;5:239-242.
DOI: 10.3329/jesnr.v5i1.11588
 23. Shah SA, Shah SM, Mohammad W, Shafi M, Nawaz H, Shehzadi S, Amir M. Effect of integrated use of organic and inorganic nitrogen sources on wheat yield. *Sarhad J. Agric*. 2010;26:559-563.
 24. Singh RJ, Ghosh BN, Sharma NK, Patra S, Dadhwal KS, Meena VS, Mishra PK. Effect of seven years of nutrient supplementation through organic and inorganic sources on productivity, soil and water conservation, and soil fertility changes of maize-wheat rotation in North-Western Indian Himalayas. *Agriculture, Ecosystems and Environment*. 2017;249 :177-186.
DOI: 10.1016/j.agee.2017.08.024
 25. Schoebitz M, Vidal G. Microbial consortium and pig slurry to improve chemical properties of degraded soil and nutrient plant uptake. *Journal of Soil Science and Plant Nutrition*. 2016;16:226-236.
DOI:10.4067/S0718-95162016005000018
 26. Rasool R, Kukal SS, Hira GS. Soil organic carbon and physical properties as affected by long-term application of FYM and inorganic fertilizers in maize-wheat system. *Soil and Tillage Research*. 2008; 101:31-36.
DOI: 10.1016/j.still.2008.05.015
 27. Singh DK, Pandey PC, Nanda G, Gupta S. Long-term effects of inorganic fertilizer and farmyard manure application on productivity, sustainability and profitability of rice-wheat system in Mollisols. *Archives of Agronomy and Soil Science*. 2019; 65:139-151.
DOI: 10.1080/03650340.2018.1491032
 28. Jackson ML. *Soil chemical analysis*. New Delhi: Prentice Hall of India Private Limited; 1973.
 29. Bahuguna A, Singh DK, Hodkashia S, Padhan SR, Verma P. Effect of different fertility levels on growth and production potential of rice genotypes. *Int. J. Plant Soil Sci*. 2023;35:539-548.
DOI: 10.9734/IJPSS/2023/v35i173243
 30. Apriyani S, Wahyuni S, Harsanti ES, Zu'Amah H, Kartikawati R, Sutriadi MT. Effect of inorganic fertilizer and farmyard manure to available P, growth and rice yield in rainfed lowland Central Java. In *IOP Conference Series: Earth and Environmental Science*. 2021;648:1-012190.
DOI: 10.1088/1755-1315/648/1/012190
 31. Abera T, Tufa T, Midega T, Kumbi H, Tola B. Effect of integrated inorganic and organic fertilizers on yield and yield components of Barley in Liben Jawi District. *International Journal of Agronomy*. 2018;1-7.
DOI: 10.1155/2018/2973286
 32. Gomez KA, Gomez AA. *Statistical procedures for agricultural research*. John Wiley and Sons; 1984.
 33. Ahmad S, Ghaffar A, Rahman MHU, Hussain I, Iqbal R, Haider G, Bashi MS. Effect of application of biochar, poultry and farmyard manures in combination with synthetic fertilizers on soil fertility and cotton productivity under arid environment. *Communications in Soil Science and Plant Analysis*. 2021;52:2018-2031.
DOI: 10.1080/00103624.2021.1908324
 34. Bhatt B, Chandra R, Ram S, Pareek N. Long-term effects of fertilization and manuring on productivity and soil biological properties under rice (*Oryza sativa*)-wheat (*Triticum aestivum*) sequence in Mollisols.

- Archives of Agronomy and Soil Science. 2016;62: 1109-1122.
DOI: 10.1080/03650340.2015.1125471
35. Meena KB, Alam MS, Singh H, Bhat MA, Singh AK, Mishra AK, Thomas T. Influence of farmyard manure and fertilizers on soil properties and yield and nutrient uptake of wheat. *International Journal of Chemical Studies*. 2018;6:386-390.
 36. Gill JS, Walia SS. Influence of FYM, brown manuring and nitrogen levels on direct seeded and transplanted rice (*Oryza sativa* L.) A review. *Research Journal of Agriculture and Environmental Management*. 2014;3:417-426.
 37. Aasif M, Chinnamani I, Kumar NS, Hemalatha M, Suresh S. Influence of integrated nutrient management practices on yield and nutrient uptake of rice under system of rice intensification. *International Journal of Advances in Agricultural Science and Technology*. 2018;5:10-6.
 38. Ahmad W, Shah Z, Khan F, Ali S, Malik W. Maize yield and soil properties as influenced by integrated use of organic, inorganic and bio-fertilizers in a low fertility soil. *Soil and Environment*. 2013;32:121-129.
 39. Kumar A, Bandyopadhyay KK, Prasad S, Kumar SN, Singh R, Kaur R, Shrivastava M. Impacts on various management practices on crops yield and soil biology in maize-wheat cropping system. *Asian Journal of Soil Science and Plant Nutrition*. 2024;10(2):445-454.
 40. Patel A, Singh S, Babu A, Verma S, Singh SK. Effect of monthly rainfall distribution on physico-chemical properties and availability of nutrients in upland red soil of Mirzapur. *Journal of Pharmacognosy and Phytochemistry*. 2018;7(1):424-429.
 41. Sharma U, Paliyal SS, Sharma SP, Sharma GD. Effects of continuous use of chemical fertilizers and manure on soil fertility and productivity of maize-wheat under rainfed conditions of the Western Himalayas. *Communications in Soil Science and Plant Analysis*. 2014;45:2647-2659.
DOI: 10.1080/00103624.2014.941854
 42. Ram S, Singh V, Sirari P. Effects of 41 years of application of inorganic fertilizers and Farm yard manure on crop yields, soil quality, and sustainable yield index under a rice-wheat cropping system on Mollisols of North India. *Communications in Soil Science and Plant Analysis*. 2016;47:179-193.
DOI: 10.1080/00103624.2015.1109653
 43. Masood S, Naz T, Javed MT, Ahmed I, Ullah H, Iqbal M. Effect of short-term supply of farmyard manure on maize growth and soil parameters in pot culture. *Archives of Agronomy and Soil Science*. 2014;60:337-347.
DOI: 10.1080/03650340.2013.792990
 44. Dhiman D, Sharma R, Sankhyan NK, Sepehya S, Sharma SK, Kumar R. Effect of regular application of fertilizers, manure and lime on soil health and productivity of wheat in an acid Alfisol. *Journal of Plant Nutrition*. 2019;42:2507-2521.
DOI: 10.1080/01904167.2019.1659317
 45. Gosal SK, Gill GK, Sharma S, Walia SS. Soil nutrient status and yield of rice as affected by long-term integrated use of organic and inorganic fertilizers. *Journal of Plant Nutrition*. 2018;41:539-544.
DOI: 10.1080/01904167.2017.1392570
DOI: 10.1080/1343943X.2019.1657777
 46. Sharma J, Goyal V, Dahiya R, Kumar M, Dey P. Response of long-term application of fertilizers and manure on P Pools in Inceptisols. *Communications in Soil Science and Plant Analysis*. 2023;54:1042-1061.
 47. Chauhan N, Sankhyan NK, Sharma RP, Singh J, Gourav. Effect of long-term application of inorganic fertilizers, farm yard manure and lime on wheat (*Triticum aestivum* L.) productivity, quality and nutrient content in an acid alfisol. *Journal of Plant Nutrition*. 2020;43:2569-2578.
DOI: 10.1080/01904167.2020.1783298
 48. Kannan RL, Dhivya M, Abinaya D, Krishna RL, Krishnakumar S. Effect of integrated nutrient management on soil fertility and productivity in maize. *Bulletin of Environment, Pharmacology and Life Sciences*. 2013;2:61-67.
 49. Katkar RN, Sonune BA, Kadu PR. Long-term effect of fertilization on soil chemical and biological characteristics and productivity under sorghum (*Sorghum bicolor*)-wheat (*Triticum aestivum*) system in Vertisol. *Indian Journal of Agricultural Sciences*. 2011;1:81- 734.
 50. Bayu W, Rethman NFG, Hammes PS, Alemu G. Effects of farmyard manure and inorganic fertilizers on sorghum growth, yield, and nitrogen use in a semi-arid area of Ethiopia. *Journal of Plant Nutrition*. 2006;29:391-407.

- DOI: 10.1080/01904160500320962
51. DV P, Krishnamurthy R, Naveen DV. Long-term effect of integrated nutrient management on soil nutrient status, content and uptake by finger millet crop in a typic kandiuustalf of eastern dry zone of Karnataka. *Communications in Soil Science and Plant Analysis*. 2020;51:161-174.
DOI: 10.1080/00103624.2019.1695829
52. Singh N, Pal N, Mahajan G, Singh S, Shevkani K. Rice grain and starch properties: Effects of nitrogen fertilizer application. *Carbohydrate Polymers*. 2011; 86:219-225.
- DOI: 10.1016/j.carbpol.2011.04.039
53. Kajisa K, Palanichamy NV. Chemical fertilizer, organic fertilizer, and cereal yields in India. *An African Green Revolution: Finding Ways to Boost Productivity on Small Farms*. 2013;225-241.
54. Olsen SR, Cole CV, Watanabe FS, Dean LA. Estimation of available phosphorus in soil by extraction with sodium bicarbonate US Dept. Agric. Circ. 1954;939.
55. Wagh NS, Katkar RN, Kharche VK. Effect of tillage and nutrient management on soil properties, growth and seed cotton yield; 2016.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/124681>